

KORESPONDENSI
WITH: RESEARCH ON CROPS (Q3) ISSN: 0972-3226

Title:
Effect of the bioherbicide application timing and plant spacing on weed growth in rice (Oryza sativa L.) cultivation


Link Web Journal: [HTTPS://GAURAVPUBLICATIONS.COM/JOURNALS/RESEARCH-ON-CROPS](https://GAURAVPUBLICATIONS.COM/JOURNALS/RESEARCH-ON-CROPS)

The screenshot shows the homepage of the 'Research on Crops' journal website. The header includes the Gaurav Publications logo (founded in 1968) and navigation links for Journals, Services, and Policies. A search bar and a 'Submit Manuscript' button are also present. The main content area features a navigation menu with links to Journal Home, Aims and Scope, Editorial Board, Current Issue, Archive, Author Guidelines, Online Published, and Publication Ethics. Below this, the journal's details are listed: Chief Editor: Dr. Vedpal Singh, Print ISSN: 0972-3226, Online ISSN: 2348-7542, Print Frequency: Quarterly, Scientific Journal Ranking (SJR): 0.22, and National Academy of Agricultural Sciences (NAAS): 4.96. A description of the journal is provided, along with a cover image of the journal.

H Index Journal: [HTTPS://WWW.SCOPUS.COM/SOURCEID/19900191751](https://www.SCOPUS.COM/SOURCEID/19900191751)

The screenshot shows the Scopus Preview page for the 'Research on Crops' journal. The page displays the journal's source details, including Scopus coverage years (from 2008 to 2023), publisher (Gaurav Publications), ISSN (0972-3226), and E-ISSN (2348-7542). The subject area is listed as Agricultural and Biological Sciences: Agronomy and Crop Science, and Agricultural and Biological Sciences: Soil Science. The source type is 'Journal'. Key performance indicators are shown: CiteScore 2022 (1.7), SJR 2022 (0.216), and SNIP 2022 (0.428). A CiteScoreTracker 2023 is also displayed, showing a CiteScore of 1.5 based on 765 citations and 527 documents to date. A notification box indicates that the CiteScore 2022 counts citations received in 2019-2022 to articles, reviews, conference papers, book chapters and data papers published in 2019-2022, and divides this by the number of publications published in 2019-2022.

Tanggal Submit: 3 Februari 2024



Dear PAIMAN,

Thank you for submitting your manuscript **Effect of the bioherbicide application timing and plant spacing on weed growth in rice (*Oryza sativa L.*) cultivation** to **Research on Crops** with the online journal management system that we are using, your manuscript is forwarded to the processing department now. You will receive official acknowledgement with the reference number within 5 working days.

You are requested to quote reference number in all your future correspondence with regard to the same. If you have any questions or query, please contact us or editor office on given emails on Contact Us Page. Thank you for considering this journal as a venue for your work.

Best Regards
Gaurav Publications



Cover letter:

COVER LETTER FOR SUBMISSION

To Research on Crops

Dear Editor,

I would like to send a research article entitled: “**Effect of the bioherbicide application timing and plant spacing on weed growth in rice (*Oryza sativa L.*) cultivation**” for Research on Crops to consider. I confirm that this work is genuine and has not been published elsewhere, nor is it considered for publication elsewhere. We believe and hope that this manuscript is worthy of publication by Research on Crops. We are interested in publishing articles in this journal because it has an excellent reputation, so it is a matter of pride if published in Research on Crops. Here I attach the manuscript.

Thank you
Best regards,

Paiman
Universitas PGRI Yogyakarta, Indonesia

Manuscript that was submitted:

**Effect of the bioherbicide application timing and plant spacing on weed growth in rice
(*Oryza sativa* L.) cultivation**

PAIMAN^{1,*}, ZULKAISI NURUL AMAR², IKHSAN FIRMANSYAH³, HAGENG
RAHMATULLOH PRIYO PAMBUDI⁴, ANGGUNILA⁵, CICILIA TRI KUSUMASTUTI⁶,
AND ARDIYANTA⁷

^{1,2,3,4,5,6,7}Department of Agrotechnology, Faculty of Agriculture, Universitas PGRI Yogyakarta,
Yogyakarta 55182, Indonesia.

¹Address: Jl. PGRI I Sonosewu No. 117, Yogyakarta, Indonesia.

Author email:

¹PAIMAN@UPY.AC.ID

²ZULKAISICUNG100@GMAIL.COM

³IKHSANDOLKENS@GMAIL.COM

⁴HAGENGRAHMATULLOH468@GMAIL.COM

⁵ANGGUNILA0811@GMAIL.COM

⁷ASTYABADY@YAHOO.COM

⁸ir.ardiyanta@upy.ac.id

*Corresponding author: Telp: +6282134391616, fax: +62274(376808), email: PAIMAN@UPY.AC.ID

ABSTRACT

Weeds around rice plants must be controlled. Cogon grass extract can be used as a bioherbicide to control weeds. Also, the use of plant spacing can suppress weed growth. Therefore, this research aimed to know the effect of application timing of cogon grass extract and plant spacing on weed growth in rice cultivation. The study was conducted from October 2022 to February 2023 in the Special Region of Yogyakarta, Indonesia. The study was 3 × 2 factorial and arranged in a randomized completely block design (RCBD). The first factor was the bioherbicide application timing: control (no application), three days before planting, and at the time of planting. The second factor was plant spacing: 15 × 15 and 25 × 25 cm. In total, there were six treatment combinations

and three replications (replications as blocks). The research results indicated the application timing of bioherbicide did not significant effect on growth of weed and rice, as well as the grain dry weight. *Fimbristylis miliacea* was obtained as the dominant weed species. Bioherbicide application at the time of planting could alter the composition of weed species compared to the control in plant spacing of 15 × 15 cm. Furthermore, the plant spacing of 25 × 25 cm resulted in higher grain and weed dry weight (19.40 g/clump and 45.00 g/0.25 m²) compared to 15 × 15 cm (12.92 g/clump and 23.04 g/0.25 m²). On the contrary, the grain dry weight per hectare was higher in 15 × 15 cm (5.74 tons/ha) than in 25 × 25 cm (3.10 tons/ha). The research findings indicate that closer plant spacing can suppress weed growth and increase the grain dry weight per hectare compared to wider plant spacing. We recommend that the use of cogon grass extract dosage needs to be increased above 50 L/ha.

Keywords: Bioherbicide, cogon grass extract, plant spacing, rice, weed

Running headline: Weed control in rice cultivation

INTRODUCTION

Rice plants are the main producers of rice to meet the food needs of the majority of Indonesians. The use of superior varieties and short-lived is the right choice to fulfill national food needs. As the population increases, the demand for rice continues to rise. However, every rice cultivation is always accompanied by the emergence of weeds around the plants. The presence of weeds significantly disrupts and harms rice growth and yield. So it must be controlled.

Many methods of weed control with chemicals have been recommended, but biological control with cogon grass (*Imperata cylindrica* (L.) Beauv.) extract is not widely known. The cogon grass extract as a bioherbicide contains allelopathic compounds. These compounds can inhibit the germination of weed seeds in the soil by hindering enzyme activity through the degradation of food reserves in the seeds. As a result, the energy produced is low due to a decrease in germination power. In addition, weed growth control can also be achieved through plant spacing.

Closer plant spacing is a method of weed control through technical cultivation. The plant canopy plays a role in reducing the distribution of sunlight to the soil surface. As a result, weed growth is inhibited due to a lack of light. The suitable plant spacing can enhance rice yield per hectare and suppress weed growth. In fertile land, a more spacious plant spacing is advisable, while

in less fertile land, a closer spacing is recommended. Short-lived varieties are better suited for close spacing, whereas long-lived varieties are suitable for wider spacing. The use of superior rice varieties requires a looser plant spacing. Increasing plant spacing provides weeds with the opportunity to grow freely and subsequently interfere with the crop. Therefore, understanding the suitable plant spacing is essential.

Weeds are troublesome plants that adversely affect the growth and yield of plants. Extracts from certain weeds have the potential as bioherbicides because they can produce allelopathy. Allelopathy is capable of reducing seed germination and slowing down the germination process. Allelopathic compounds can inhibit the activity of enzymes that degrade food reserves in seeds, resulting in very low energy production and decreasing germination potential.

In general, plant-derived bioherbicides contain phytotoxic allelochemicals or specific disease-carrying microbes (Hasan *et al.*, 2021). The advantages of using bioherbicides are that they contain no toxic organic secondary metabolites, that they are easy to acquire from the field, that they have no secondary metabolite mechanisms found in synthetic herbicides, that they have more than one secondary metabolite compound, and that they do not cause plant poisoning (Sihombing *et al.*, 2018). When bioherbicides are absorbed by weed seeds or roots, they disrupt the integrity of cell membranes and other essential biochemical processes. It should be noted that the cytotoxic effect on weeds is reflected in low levels of root cell division, low ability to absorb nutrients, growth hormones, and pigment synthesis, as well as low growth of reactive oxygen species (ROS), stress-related hormones, and abnormal antioxidant activity (Hasan *et al.*, 2021).

One type of weed with potential as a bioherbicide is *Imperata cylindrica* (Sari *et al.*, 2017). The roots of cogon grass can be used as a pre-emergence bioherbicide. Cogon grass root extract contains allelochemical compounds, specifically flavonoids. A concentration of 1% root extract of cogon grass is capable of inhibiting the growth of weed seeds located below the ground (Sari *et al.*, 2017). Furthermore, a concentration of 14% extract from cogon grass rhizomes can effectively control weeds such as *Ageratum conyzoides*, *Eleusine indica*, and *Cyperus rotundus* (Lau *et al.*, 2021).

Extract cogon grass contains terpenoid, phenolic, and steroid compounds (Erida *et al.*, 2019). The extracts, leachates, root exudates, or growing medium of cogon grass also contained several allelochemicals, including phenolic acids, phenolic aldehydes, phenylpropanoids, flavonoids, quinones, fatty acids, terpenoids, simple phenolics, benzoic acids, and alkaloids (Kato-Noguchi,

2022). The application of cogon grass extract at the time of planting with a concentration of 30% could increase the number of panicles and the weight of dry rice grains in rice cultivation in polybags. It had a greater potential as a bioherbicide to suppress weed growth and increase rice yield (Paiman *et al.*, 2022).

The percentage of germination, germination rate, and seedling length of weed seeds such as *Amaranthus spinosus*, *Bidens biternata*, and *Tridax procumbens* could all be greatly suppressed by using cogon grass extract. *Tridax procumbens* had the highest level of sensitivity among weed seeds to the extract from the roots and rhizomes of cogon grass, followed by *Bidens biternata* and *Amaranthus spinosus*. Extracts from the roots and rhizomes of cogon grass exhibit significant potential as bioherbicides to inhibit the germination of weed seeds (Pujiwati, 2011). On the other hand, using close plant spacing can also suppress weed growth and increase yield per unit area. But, grain yield per clump decreases. The appropriate plant spacing can provide plants with the opportunity to grow well without competition from neighboring plants for water, nutrients, and sunlight. The suitable plant spacing can enhance rice yield. Also, the use of plant spacing depends on the cultivated rice variety.

Rice plants with curved panicles were used for planting density was not too tight and not too loose, but rice plants with erect panicles were better with planting density quite tight (Yena *et al.*, 2018). Wide inter-row width and spacing in narrow rows will be conducive to building a high-yield population to increase yields (Li *et al.*, 2016). If the spacing between rows was fixed at 30 cm, then the spacing in rows of 11-12 cm could increase plant population, and grain yield, and was the optimal plant spacing for rice cultivation (Yun *et al.*, 2020). Grain yield was maintained in a large number of plant populations, primarily through the effects of more tillers per plant and more grain per panicle in lower plant populations (Dunn *et al.*, 2020).

Plant spacing significantly influenced plant height, tiller number, and rice panicle length. A plant spacing of 15 × 15 cm provided the highest panicle count and good yield per square meter (Marie-Noel *et al.*, 2021). The optimal Jajar Legowo system involved skipping one row after every three rows at plant spacing of 15 × 15 cm, resulting in the highest grain yield (Rautaray, 2016). Consistent with subsequent research, the highest grain dry weight was found at a plant spacing of 15 × 15 cm compared to 20 × 20 cm and 25 × 25 cm (Rautaray, 2016). In line with further studies, plant density with a plant spacing of 15 × 15 cm resulted in the highest grain yield (4.71 tons/ha), while the 25 × 25 cm treatment yielded a lower amount (4.09 tons/ha) (Alogaidi

et al., 2019). Also supported by Faisul-ur-Rasool *et al.* (2012), closer spacing (15×15 cm) produced the maximum tillers number, leaf area index, dry matter accumulation, and rice yield. Plant spacing of 15 × 15 cm resulted in a yield of 8.97% greater than 20 × 20 cm because of better absorption of photosynthetically active light.

Plant population had a significant effect on morphological and physiological characteristics as well as crop yield. The leaf area per clump was broader with a looser plant spacing. Additionally, wider plant spacing results in lower competition among plants, allowing them to acquire more nutrients. As a result, there were more tillers, a larger leaf area, and greater dry weight per cluster, as opposed to dense plant spacing (Mondal *et al.*, 2013). There was limited information available on the optimal row width needed to maximize grain yield potential (Dunn *et al.*, 2020). Superior rice varieties prefer a looser plant spacing for their growth compared to local varieties, but yield per unit area needs to be maximized using closer plant spacing.

Based on the literature review above, cogon grass extract as a bioherbicide has proven to be quite effective in biologically controlling weeds, particularly at a concentration of 30%. However, this bioherbicide has not yet been directly applied in rice fields, and the appropriate application timing in rice cultivation remains unknown. Additionally, it has been established from several previous studies that the use of close plant spacing (15 × 15 cm) in rice cultivation can effectively suppress weed growth and increase yield. However, the use of plant spacing has not been investigated in the cultivation of the short-lived rice of the Padjajaran Agritan variety.

Research about the application timing of cogon grass extract as a bioherbicide and the plant spacing to suppress weed growth and increase rice yield needs to be conducted. This study is expected to contribute insights into environmentally friendly weed control through biological and technical cultural approaches. Therefore, this research aimed to know the effect of application timing of cogon grass extract and plant spacing on weed growth in rice cultivation of the Padjajaran Agritan variety.

MATERIALS AND METHODS

Study Site

The research was conducted from October 2022 to February 2023 during the rainy season. The study was located in Minggir Sub-district, Sleman Regency, Special Region of Yogyakarta, Indonesia. The elevation of the location was 110 meters above sea level. Geographically, Sleman

Regency is situated between 110°33' 00" - 110°13' 00" East Longitude and 7°34'51" - 7°47'30" South Latitude.

Experimental Design

The study was 3×2 factorial and arranged in a randomized completely block design (RCBD). The first factor was the application timing of bioherbicide consisting of three methods: control (no application), three days before planting, and at the time of planting. The second factor was the use of plant spacing, consisting of two kinds: 15×15 cm (close plant spacing) and 25×25 cm (wide plant spacing). In total, there were six treatment combinations. Each treatment combination was repeated three times (replications as blocks), resulting in a total of 18 treatment plots. Treatment plots of plant spacings of 15×15 cm and 25×25 cm can be seen in Figure 1.

Research Procedures

In this study, seeds of the Padjajaran Agritan variety were used. Before sowing, the seeds were soaked in water for three hours to ensure uniform imbibition. Subsequently, the seeds were lifted from the soaking medium, and the excess water was drained. After one day and one night, roots emerged from the base of the seeds. Plastic seedling trays measuring 50×60 cm with a height of 10 cm were used for seed germination. The germination medium consisted of a mixture of organic fertilizer and soil in a 1:1 ratio. The seeds were then sown on the surface of the germination medium at field capacity. Four days later, shoots emerged above the soil surface, and the seedlings were ready to be transplanted into the rice field at the age of 18 days after sowing.

Soil processing was conducted in two stages. The first stage of soil processing involved using a rotary plow to overturn and crush the soil. The second stage of processing utilized a harrow to level the soil. Subsequently, treatment plots were established with dimensions of 3×3 m, totaling 18 treatment plots. Drainage channels with a width of 0.5 m were created between the treatment plots. Treatment randomization was performed per block for all blocks.

In this research, cogon grass extract was used as a bioherbicide. The application timing of bioherbicide corresponds to the randomized treatment results. The control plot was without bioherbicide treatment, while the treatment plots involved the use of a 30% concentration of bioherbicide with a dosage of 45 ml/3 m² (50 L/ha). The application timing of bioherbicide was divided into two methods: three days before planting and at the time of planting. Bioherbicide was

applied only once for each treatment in research. During the three days following the treatment, measures were taken to ensure that no external water entered the treatment plots.

The seedlings number that planted in each treatment plot depends on the plant spacing. For a plant spacing of 15×15 cm, 20 seedlings were planted in rows and 20 seedlings between rows, requiring a total of 400 seedlings per plot (Figure 1a). Meanwhile, for a plant spacing of 25×25 cm, 12 seedlings were planted in rows and 12 seedlings between rows, requiring a total of 144 seedlings per plot (Figure 1b). Only one rice seedling was planted per planting hole.




Figure 1. Plot treatments of plant spacing of 15×15 cm (a) and 25×25 cm (b)

Recommended fertilizer usage was 225 kg/ha of urea and 225 kg/ha of NPK Phonska 15-15-15 (BPPP, 2014). Fertilizer application was divided into two stages. In the first stage, 40% of the recommended dosage was applied 15 days after planting (DAP). The second stage involved applying 60% of the recommended dosage at 35 DAP. Irrigation was carried out every four days after the treatment. Pest and disease control were implemented during the rice flowering stage to minimize the impact of brown planthoppers. Harvesting was conducted when the rice plants reached 104 DAP.

Parameters

Observations on weed growth were conducted at 60 days after transplanting (DAT) and included the number of weeds (individual/ 0.25 m^2), weed dry weight ($\text{g}/0.25 \text{ m}^2$), weed dominance, and weed community coefficient (%). Observations on rice plants were performed on

growth and yield components, including the tillers number (stalks/clump) observed at 60 DAT, plant dry weight (g/clump), panicles number (strands/clump), grain dry weight (g/clump), and hectare (ton/ha), all observed at 104 DAT.

Statistical Analysis

Observational data were analyzed by analysis of variance (ANOVA) at $P = 0.05$ probability levels with the software IBM SPSS Statistic 23. In addition, Duncan's new multiple range test (DMRT) at $P = 0.05$ probability levels was used to test the difference between treatment averages. To determine the dominant weed species and weed community sift, the summed dominance ratio (SDR) and coefficient of weed community were used.

RESULTS AND DISCUSSION

Number and Dry Weight of Weeds

The timing of the bioherbicide application did not have a significant effect on the number and dry weight of weeds. However, only the plant spacing had an impact on the weed dry weight. The average number and dry weight of weeds can be seen in Table 1.

Table 1. Effect of bioherbicide application and plant spacing on the weed growth per plot

Treatments	Parameter of observations	
	Weed number (individual/0.25 m ²)	Weed dry weight (g/0.25 m ²)
<u>Timing of bioherbicide application</u>		
Control	58.17 a	32.08 a
Three days before planting	41.67 a	28.97 a
At the time of planting	44.67 a	41.01 a
<u>Plant spacing (cm)</u>		
15 × 15	41.22 p	23.04 q
25 × 25	55.11 p	45.00 p

Remarks: According to DMRT at $P = 0.05$ probability levels, there is no significant difference between the numbers in the same column followed by the same characters.

Table 1 shows that a plant spacing of 25 × 25 cm resulted in a higher weed dry weight compared to 15 × 15 cm. Wider plant spacing contributed to a heavier weed dry weight. In wider plant spacing, sunlight penetration can reach the soil surface, and then support weed growth. The weed dry weight at a plant spacing of 15 × 15 cm was lower than that at 25 × 25 cm. It indicates that closer plant spacing is more effective in suppressing weed growth because the plant canopy quickly covers the soil surface. Sunlight penetration to the soil surface is lower, so inhibits weed growth. In line with Fahad *et al.* (2015), less weed growth was seen at narrow row spacing (11 cm) compared to wider row spacing (15 and 23 cm) in wheat crops. In line with the research result by Chauhan and Johnson (2010) narrowing the row spacing could decrease weed growth and seed production, as well as increase the grain yield of rice.

Dominance of Weed Species

In this study, three types of weeds were identified based on their morphology, namely sedges, grasses, and broad-leaved weeds, with a total of nine weed species overall. The effect of the combination of bioherbicide application timing and plant spacing on the SDR of each weed species can be seen in Table 2.

Table 2. SDR (%) of each weed species

No.	Weed species	SDR (%)					
		J ₁ B ₀	J ₁ B ₁	J ₁ B ₂	J ₂ B ₀	J ₂ B ₁	J ₂ B ₂
1	<i>Cyperus brevifolius</i> (Rottb.) Rassk.	0.00	2.56	0.00	0.00	0.00	0.00
2	<i>Cyperus rotundus</i> L.	0.00	0.00	13.58	10.86	5.26	4.05
3	<i>Scirpus mucronatus</i> L.	10.72	6.29	4.06	6.35	0.00	3.24
4	<i>Cynodon dactylon</i> (L.) Pers.	4.92	7.57	10.51	12.45	5.50	8.10
5	<i>Echinochloa colona</i> (L.) Link.	5.55	9.70	16.75	6.73	4.27	5.94
6	<i>Echinochloa crus-galli</i> (L.) P. Beauv.	18.90	8.49	0.00	12.37	27.1	9.42
7	<i>Fimbristylis miliacea</i> (L.) Vahl.	47.48	50.54	44.34	51.21	50.3	52.26
8	<i>Limnocharis flava</i> (L.) Buch	0.00	7.87	2.91	0.00	0.00	6.66
9	<i>Monochoria vaginalis</i> (Burm.f.) C.Presl.	12.39	6.95	7.82	0.00	7.49	10.29

Remarks: J₁ = Plant spacing of 15 × 15 cm, J₂ = Plant spacing of 25 × 25 cm, B₀ = Control, B₁ = Three days before planting, and B₂ = At the time of planting.

Table 2 indicates that the identified sedges included *Cyperus brevifolius*, *Cyperus rotundus*, and *Scorpus mucronatus*. The grass weeds identified in this study were *Cynodon dactylon*, *Echinochloa colona*, *Echinochloa crus-galli*, and *Fimbristylis miliacea*. Several types of broad-leaved weeds found in this research were *Limnocharis flava* and *Monochoria vaginalis*. *Fimbristylis miliacea* was the dominant weed species followed by *Echinochloa crus-galli* in this study. It can be explained that grass weeds more dominate in rice cultivation than other types of weeds. It was supported by Kurniadie *et al.* (2019), *Fimbristylis miliacea* was a dominant weed species in conventional and organic rice cultivation.

Weed Community Shift

Based on Table 2, the weed community coefficient (C%) can be calculated between weed communities, both among the application timing of bioherbicide and plant spacing treatments. The results of the weed community coefficient are presented in Tables 3 and 4.

Table 3. Weed community coefficient between application timing of bioherbicide in each plant spacing

Plant spacing (cm)	The comparison between application timing of bioherbicide.	Weed community coefficient (C%)	Information
15 × 15	1. Control and three days before planting	79.68	Homogeny
	2. Control and at the time of planting	66.69	No homogeny
	3. Three days before planting and at the time of planting	75.53	Homogeny
25 × 25	1. Control and three days before planting	77.70	Homogeny
	2. Control and at the time of planting	81.96	Homogeny
	3. Three days before planting and at the time of planting	81.03	Homogeny

Table 4. Weed community coefficient between plant spacing in application time of bioherbicide

Application timing of bioherbicide	The comparison between plant spacing	Weed community coefficient (C%)	Information
Control	15 × 15 and 25 × 25 cm	76.67	Homogeny
Three days before planting	15 × 15 and 25 × 25 cm	75.51	Homogeny
At the time of planting	15 × 15 and 25 × 25 cm	76.40	Homogeny

Table 3 indicates that the coefficient of weed community in the comparison between the control and application timing of bioherbicide at the time of planting was 66.69% in a plant spacing of 15 × 15 cm. It shows a shift in the weed species composition due to the application timing of bioherbicide at the time of planting. In other words, there is a difference in the weed species that grow in these two communities. Meanwhile, for the other communities, there is no shift in the weed species community because their community coefficients were > 75%.

Growth and Yield of Rice

Observations on the growth and yield of rice included the tillers number, plant dry weight, panicles number, grain dry weight per clump, and hectare. The application timing of bioherbicide did not have a significant effect on all the observed parameters. However, the plant spacing significantly influenced all the observed parameters. The average tillers number, plant dry weight, panicles number, grain dry weight per clump, and hectare can be seen in Table 5.

Table 5. The effect of application timing of bioherbicide and plant spacing on the growth and yield of rice

Treatments	Parameters			
	Tillers number (stem/clump)	Plant dry weight (g/clump)	Panicles number (strands/clump)	Grain dry weight (g/clump) (tons/ha)
<u>Application timing of bioherbicide</u>				
Control	10.57 a	9.97 a	9.06 a	14.90 a 4.15 a

Three days before planting	11.37 a	13.93 a	9.13 a	16.58 a	4.32 a
At the time of planting	10.73 a	12.94 a	9.50 a	17.00 a	4.80 a
<u>Plant spacing (cm)</u>					
15 × 15	9.27 q	10.75 q	7.75 q	12.92 q	5.74 p
25 × 25	12.51 p	14.91 p	10.71 p	19.40 p	3.10 q

Remarks: According to DMRT at P = 0.05 probability levels, there is no significant difference between the numbers in the same column followed by the same characters.

Table 4 indicates that a plant spacing of 25 × 25 cm resulted in tillers number (12.51 stem), plant dry weight (14.91 g), panicles number (10.71 strands), and grain dry weight (19.40 g) per clump, respectively. Plant spacing of 25 × 25 cm was given growth and yield component per clump higher than in 15 × 15 cm. The higher tillers number, plant dry weight, and panicles number at wider plant spacing correlated with the grain dry weight per clump. The result of this study was supported by Dass *et al.* (2017), wider plant spacing significantly increased rice yield compared with closer plant spacing. In line with Anwari *et al.* (2019), yield components like tillers number and grain per panicle were found in plant spacing 25 × 25 cm compared to 10 × 10 cm, 15 × 15 cm, or 20 × 20 cm.

A plant spacing of 25 × 25 cm could yield higher grain dry weight per clump because of a better microclimate compared to closer plant spacing. It caused lower competition among plants for nutrients, improved water access, and smoother air circulation. Rice plants could grow well so that the size of the habitus was larger. Additionally, sunlight distribution was more even across the entire plant body, leading to enhanced photosynthesis. These conditions resulted in a more efficient carbohydrate-filling process for all seeds in the panicle. With a greater seed number per clump and a heavier weight per seed, the total grain dry weight was higher. On the other hand, sunlight penetration to the soil surface could benefit weeds to support their growth.

However, a plant spacing of 15 × 15 cm could provide a higher grain dry weight per hectare compared to 25 × 25 cm. In closer plant spacing, the population density per hectare was higher, leading to a higher yield per hectare. The highest radiation use efficiency was obtained at close plant spacing which resulted in a higher amount of biomass for energy production. Following the opinion Ya-jie *et al.* (2016), higher density was applied to increase panicles and grains number for

higher yield per unit area. Supported by Marie-Noel *et al.* (2021), the highest panicle number/m² was found in plant spacing of 15 × 15 cm but not obtained in 20 × 20 cm, 25 × 25 cm, or 30 × 30 cm. In contrast with the study results by Asghar *et al.* (2021), the highest grain yield of 6.75 tons/ha was obtained in plant spacing of 23 × 23 cm followed by 5.46 tons/ha from 18 × 18 cm. No in line with the study results by Neupane (2022), loose plant spacing (20 × 20 cm) was given superior morphophysiological and yield components and resulted in a grain yield of 8.53 tons/ha higher than 6.47 tons/ha in tight plant spacing (20 × 10 cm).

The use of a plant spacing of 15 × 15 cm resulted in plant populations higher than 25 × 25 cm, with 444,444 plants/ha higher than 160,000 plants/ha. Total grain yield per hectare could be increased by close plant spacing. The plant population at a plant spacing of 15 × 15 cm was almost three times higher than 25 × 25 cm. The panicles number per hectare was found to be higher at a plant spacing of 15 × 15 cm which was 3,444,441 panicles greater than 1,713,600 panicles in 25 × 25 cm. It means that a greater panicle number could support the total grain dry weight per hectare, although the total grain dry weight per panicle was lower. Deng *et al.* (2022) explained that increasing light utilization by the canopy was a major way to increase photosynthesis and rice productivity.

In this study, the Padjajaran Agritan variety was used, which has short-lived growth, fewer tillers, and a smaller plant size. Therefore, using a closer plant spacing was the best solution to achieve a higher yield per hectare by intercepting more solar radiation. It can be concluded that using a closer plant spacing for rice varieties with a compact habitus is more advantageous because it can capture and harness more sunlight penetration. Additionally, the growth of weeds under the plant canopy can be inhibited.

CONCLUSIONS

In conclusion, the the application timing of bioherbicide did not significantly affect on growth of weed and rice, as well as the grain dry weight. *Fimbristylis miliacea* was obtained as the dominant weed species. Bioherbicide application at the time of planting could alter the composition of weed species compared to the control in plant spacing of 15 × 15 cm. Furthermore, the plant spacing of 25 × 25 cm resulted in higher grain and weed dry weight (19.40 g/clump and 45.00 g/0.25 m²) compared to 15 × 15 cm (12.92 g/clump and 23.04 g/0.25 m²). On the contrary, the grain dry weight per hectare was higher in 15 × 15 cm (5.74 tons/ha) than in 25 × 25 cm (3.10

tons/ha). The research findings indicate that closer plant spacing can suppress weed growth and increase the grain dry weight per hectare compared to wider plant spacing. We recommend that the use of cogon grass extract dosage needs to be increased above 50 L/ha.

CONFLICT OF INTEREST

The authors declare no conflict of interest, financial or otherwise

FUNDING STATEMENT

Not applicable.

ACKNOWLEDGEMENTS

We would like to thank the Institute for Research and Community Service at Universitas PGRI Yogyakarta for funding and supporting this research. Furthermore, we would like to thank the Faculty of Agriculture, Universitas PGRI, which has provided financing for greenhouse facilities and equipment needed for research.

REFERENCES


- Alogaidi, F. F., Al-shugeairy, Z. K., Hadi, B. H. and Hassan, W. A. (2019). Effect of planting distance on yield and yield components of four effect of planting distance on yield and yield components of four introduced upland rice varieties. *Plant Arch.* **19**: 699–707. [HTTPS://PLANTARCHIVES.ORG/PDF%2019-1/699-707%20\(4827\).PDF](https://plantarchives.org/pdf%2019-1/699-707%20(4827).pdf)
- Anwari, G., Moussa, A. A., Wahidi, A. B., Mandozai, A., Nasar, J. and El-Rahim, M. G. M. A. (2019). Effects of planting distance on yield and agro-morphological characteristics of local rice (Bara variety) in Northeast Afghanistan. *Curr. Agric. Res. J.* **7**: 350–7. [HTTPS://DOI.ORG/10.12944/CARJ.7.3.11](https://doi.org/10.12944/CARJ.7.3.11)
- Asghar, M., Hassan, T., Arshad, M., Aziz, A., Latif, M. T. and Sabir, A. M. (2021). Effect of plant spacing on incidence of rice planthoppers in transplanted rice crop. *Int. J. Trop. Insect. Sci.* **41**: 575–85. [HTTPS://DOI.ORG/10.1007/s42690-020-00242-4](https://doi.org/10.1007/s42690-020-00242-4)
- Chauhan, B. S. and Johnson, D. E. (2010). Implications of narrow crop row spacing and delayed *Echinochloa colona* and *Echinochloa crus-galli* emergence for weed growth and crop yield loss in aerobic rice. *F. Crop. Res.* **117**: 177–82. [HTTPS://DOI.ORG/10.1016/J.FCR.2010.02.014](https://doi.org/10.1016/j.fcr.2010.02.014)
- Dass, A., Chandra, S., Uphoff, N., Choudhary, A. K., Bhattacharyya, R. and Rana, K. S. (2017). Agronomic fortification of rice grains with secondary and micronutrients under differing crop


- management and soil moisture regimes in the North Indian Plains. *Paddy Water Environ.* **15**: 745–60. [HTTPS://LINK.SPRINGER.COM/ARTICLE/10.1007/S10333-017-0588-9](https://link.springer.com/article/10.1007/s10333-017-0588-9)
- Deng, F., Li, B., Yuan, Y., He, C., Zhou, X., Li, Q., Zhu, Y., Huang, X., He, Y., Ai, X., Tao, Y., Zhou, W., Wang, L., Cheng, H., Chen, Y., Wang, M. and Ren, W. (2022). Increasing the number of seedlings per hill with reduced number of hills improves rice grain quality by optimizing canopy structure and light utilization under shading stress. *F. Crop. Res.* **287**: 108668. [HTTPS://DOI.ORG/10.1016/J.FCR.2022.108668](https://doi.org/10.1016/j.fcr.2022.108668)
- Dunn, B. W., Dunn, T. S., Mitchell, J. H. and Brinkhoff, J. (2020). Effects of plant population and row spacing on grain yield of aerial-sown and drill-sown rice. *Crop. Pasture Sci.* **71**: 219–28. [HTTPS://DOI.ORG/10.1071/CP19421](https://doi.org/10.1071/CP19421)
- Erida, G., Saidi, N., Hasanuddin, and Syafruddin. (2019). Allelopathic screening of several weed species as potential bioherbicides. The 3rd CSAFS. *IOP Conf. Series: Earth Environ. Sci.* **334**: 012034. [HTTPS://DOI.ORG/10.1088/1755-1315/334/1/012034](https://doi.org/10.1088/1755-1315/334/1/012034)
- Fahad, S., Hussain, S., Chauhan, B. S., Saud, S., Wu, C., Hassan, S., Tanveer, M., Jan, A. and Huang, J. (2015). Weed growth and crop yield loss in wheat as influenced by row spacing and weed emergence times. *Crop. Prot.* **71**: 101–8. [HTTPS://DOI.ORG/10.1016/J.CROPRO.2015.02.005](https://doi.org/10.1016/j.cropro.2015.02.005)
- Faisul-ur-Rasool, Habib, R. and Bhat, M. I. (2012). Evaluation of plant spacing and seedlings per hill on rice (*Oryza sativa* L.) productivity under temperate conditions. *Pakistan J. Agric. Res.* **49**: 169–72.
- Hasan, M., Ahmad-hamdani, M. S., Rosli, A. M. and Hamdan, H. (2021). Bioherbicides: An eco-friendly tool for sustainable weed management. *Plants.* **10**: 1212. [HTTPS://DOI.ORG/10.3390/PLANTS10061212](https://doi.org/10.3390/plants10061212)
- Kato-Noguchi, H. (2022). Allelopathy and allelochemicals of *Imperata cylindrica* as an invasive plant species. *Plants.* **11**: 2551. [HTTPS://DOI.ORG/10.3390/PLANTS11192551](https://doi.org/10.3390/plants11192551)
- Kurniadie, D., Irda, M., Umiyati, U., Widayat, D. and Nasahi, C. (2019). Weeds diversity of lowland rice (*Oryza sativa* L.) with different farming system in Purwakarta Regency Indonesia. *J. Agron.* **18**: 21–6.
- Lau, D. F. W., Sofian, and Mirza, A. (2021). Ekstrak rimpang alang-alang (*Imperata cylindrica* L.) sebagai herbisida nabati untuk mengendalikan gulma. *J. Agroekoteknologi Trop. Lembab.* **4**: 29–34. (in Indonesian)
- Li, G., Zhang, J., Yang, C., Liu, Z., Wang, S. and Ding, Y. (2016). Population characteristics of

- high-yielding rice under different densities. *Agron. J.* **108**: 1415–23.
[HTTPS://DOI.ORG/10.2134/AGRONJ2015.0214](https://doi.org/10.2134/AGRONJ2015.0214)
- Marie-Noel, M. F., Joseph, B. M., Liliane, T., Felix, F., Dorothy, M. and Salvador, D. P. (2021). Effect of plant spacing on the growth and yield of rainfed rice (*Oryza sativa*) in the bimodal rain forest zone of Cameroon. *J. Agric. Crop.* **7**: 48–59. [HTTPS://DOI.ORG/10.32861/JAC.72.48.59](https://doi.org/10.32861/JAC.72.48.59)
- Mondal, M. M. A., Puteh, A. B., Ismail, M. R. and Rafii, M. Y. (2013). Optimizing plant spacing for modern rice varieties. *Int. J. Agric. Biol.* **15**: 175–8.
- Neupane, P. (2022). Effect of plant spacing and genotype on growth, yield, and yield-related traits of rice varieties. *Russ. J. Agric. Socio-Economic Sci.* **3**: 89–98.
[HTTPS://DOI.ORG/10.18551/RJOAS.2022-03.10](https://doi.org/10.18551/RJOAS.2022-03.10)
- Paiman, Hidayat, A. K., Shobirin, S. S. and Khasanah, S. I. (2022). Efficacy of weed extract as a bioherbicide in rice (*Oryza sativa* L.) cultivation. *Res. Crop.* **23**: 488–96.
- Pujiwati, I. (2011). Land use potential through reseds (*Imperata cylindrica*) as bioherbicide. *Gea.* **11**: 226–34.
- Rautaray, S. K. (2016). Effect of planting geometry and fertilizer dose on grain yield and disease incidence in some mid duration rice (*Oryza sativa* L.) varieties. *Arch. Agron. Soil Sci.* **53**: 327–33. [HTTPS://DOI.ORG/10.1080/03650340701223320](https://doi.org/10.1080/03650340701223320)
- Sari, V. I., Nanda, S. and Sinuraya, R. (2017). Bioherbisida pra-tumbuh alang-alang (*Imperata cylindrica*) untuk pengendalian gulma di perkebunan kelapa sawit. *J. Citra Widya Edukasi.* **9**: 301–8. (in Indonesian)
- Sihombing, N., Purba, Z., Samosir, S. and Karim, S. (2018). Potensi metabolit sekunder gulma sebagai pestisida nabati di Indonesia (Potency of secondary metabolite from weeds as natural pesticides in Indonesia). *J. Kultiv.* **17**: 683–93. (in Indonesian)
- Ya-jie, H., Hai-jun, Q., Wei-wei, C., Zhi-peng, X., Hong-cheng, Z., Qi-gen, D., Zhong-yang, H., Ke, X., Hai-yan, W. and Bao-wei, G. (2016). Effect of different mechanical transplantation methods and density on yield and its components of different panicle-typed ricechinese. *Chinese J. Rice Sci.* **2016**: 493–506. [HTTPS://DOI.ORG/10.16819/J.1001-7216.2016.6015](https://doi.org/10.16819/J.1001-7216.2016.6015)
- Yena, W., Xiang, Z., Guotao, Y., Xuechun, W., Youlin, P., Yao, H. and Yungao, H. (2018). Effects of planting density on the microclimate of rice population with different panicle-types and its yield components. *Chinese J. Appl. Environ. Biol.* **2018**: 813–23.
[HTTPS://DOI.ORG/10.19675/J.CNKI.1006-687X.2017.12049](https://doi.org/10.19675/J.CNKI.1006-687X.2017.12049)
- Yun, C., Yajun, Z., Honglu, Z., An, Z., Jian, H., Hao, Z., Junfei, G., Lijun, L., & Jianchang, Y.


(2020). Effects of plant spacing on grain yield and population quality in mechanically-transplanted rice with good tasting quality. *Chinese J. Rice Sci.* **2020**: 550–60. [HTTPS://DOI.ORG/10.16819/J.1001-7216.2020.0205](https://doi.org/10.16819/j.1001-7216.2020.0205)

Balasan Email dari Pihak Jurnal: 3 Februari 2024





Perbaiki Manuskript: 22 Maret 2024



mail.google.com/mail/u/0/#inbox/FMfcgzGxSRHmrPkCQRJgmhRTSgJLqXvc

Search in mail

Active

2 of 5,423

Comments of Reviewer on the manuscript " ROC-1057 "

External Inbox

Gaurav Publications
to me, paimanupy

2:45 PM (3 hours ago)

Dear PAIMAN,

It is with reference to your Manuscript No. Gaurav Publications/ROC-1057 entitled **Effect of the bioherbicide application timing and plant spacing on weed growth in rice (*Oryza sativa* L.) cultivation** which you have sent for processing in "Research on Crops", comments by reviewer are attached, you are requested to revise the article carefully as per comments and follow guidelines to authors given on the website www.gauravpublications.com. Submit the corrected complete article including all author names so that further action can be taken.

Give point wise reply to reviewer comments in Word File .

Attached is one Sample Manuscript as well. You are also advised to revise the article and references according to the Format of the journal. Click here for author guidelines and Sample Paper - <http://gauravpublications.com/author-guidelines>

Guidelines for Authors:-

1. Full-length articles (not exceeding 3000-3500 words including references, tables and figures).
2. Review Articles (not exceeding 6000 words)
3. Kindly mention the name of the Institute where work was carried out in Materials and Methods. (Mandatory*)
5. Give year of experiment with research period in Materials and Methods. (Mandatory)

Note:- No further authors will be added after receiving the revised article.

Resubmit your revised article as the reply of this email.

mail.google.com/mail/u/0/#inbox/FMfcgzGxSRHmrPkCQRJgmhRTSgJLqXvc

Search in mail

Active

2 of 5,423

Resubmit your revised article as the reply of this email.

Best Regards
Managing Editor
Gaurav Publications |
1314 (GF), Housing Board, Sector-15A, | Hisar - 125001 | Haryana (INDIA) |
Web: www.gauravpublications.com | E-mail: info@gauravpublications.com

2 Attachments • Scanned by Gmail

ROC-1057 - Revis...
ROC-1057 Refere...

Dr. Ir. Paiman, M.P. - UPY
to Gaurav, paimanupy

6:47 PM (0 minutes ago)

Ok, I will do it.

Reply Reply all Forward

RESEARCH ON CROPS
Gaurav Publications (Regd.), Hisar, India

Ref. No. GP-ROC/2023/ROC/1057

Date: 06/03/2024


REFEREE'S COMMENTS

Title of the Article/note: Optimizing Bioherbicide Application Timing and Plant Spacing to Manage Weed Growth in Rice (*Oryza sativa* L.) Cultivation

Please tick (√) in the brackets of appropriate word (s) along with comments if any.

- | | | | |
|--|-------|--|-------------------|
| 1. Title | | | |
| Suitable | (√) | | Title revised |
| Need modification | () | | |
| 2. Abstract | | | |
| Adequate | (√) | | Revised |
| Inadequate | () | | |
| Should be rewritten | () | | |
| 3. Introduction | | | |
| Adequate | () | | Citations missing |
| Inadequate | (√) | | |
| Should be rewritten | () | | |
| 4. Materials and Methods | | | |
| Adequate | (√) | | Well written |
| Inadequate | () | | |
| Should be rewritten | () | | |
| 5. Statistical analysis | | | |
| Adequate | (√) | | Sufficient |
| Inadequate | () | | |
| Erroneous | () | | |
| Not analysed | () | | |
| 6. Interpretation of results and discussion | | | |
| Adequate | (√) | | Sufficient |
| Not supported by data | () | | |
| Suffers from omissions | () | | |
| Vague and generalized | () | | |
| Too brief for clarity | () | | |
| Too comprehensive, must be condensed | () | | |
| Result and Discussion need to be merged | () | | |
| 7. Conclusion | | | |
| Adequate | (√) | | Sufficient |
| Need modification | () | | |
| To be rewritten | () | | |

Submit Perbaikan Manuscript: 22 Maret 2024



Manuscript yang telah diperbaiki (Revisi judul):

Weed control in rice cultivation

Optimizing Bioherbicide Application Timing and Plant Spacing to Manage Weed Growth in Rice (*Oryza sativa* L.) Cultivation

PAIMAN^{1,*}, ZULKAISI NURUL AMAR¹, IKHSAN FIRMANSYAH¹, HAGENG RAHMATULLOH PRIYO PAMBUDI¹, ANGGUNILA¹, CICILIA TRI KUSUMASTUTI¹ AND ARDIYANTA¹

¹*Department of Agrotechnology, Faculty of Agriculture
Universitas PGRI Yogyakarta, Yogyakarta 55182, Indonesia*

**(e-mail: PAIMAN@UPY.AC.ID)*

ABSTRACT

Weed management is crucial for successful rice cultivation. Bioherbicides derived from cogon grass extract offer a promising solution for weed control. Additionally, optimizing plant

spacing has been recognized as an effective method to suppress weed growth. Thus, this study aims to investigate the impact of cogon grass extract application timing and plant spacing on weed growth in rice cultivation. The study was conducted from October 2022 to February 2023 in the Special Region of Yogyakarta, Indonesia. The study was 3×2 factorial and arranged in a randomized completely block design (RCBD). The first factor was the bioherbicide application timing: control (no application), three days before planting, and at the time of planting. The second factor was plant spacing: 15×15 and 25×25 cm. In total, there were six treatment combinations and three replications (replications as blocks). The research results indicated the application timing of bioherbicide did not significant effect on the growth of weed and rice, as well as the grain dry weight. *Fimbristylis miliacea* was obtained as the dominant weed species. Bioherbicide application at the time of planting could alter the composition of weed species compared to the control in plant spacing of 15×15 cm. Furthermore, the plant spacing of 25×25 cm resulted in higher grain and weed dry weight (19.40 g/clump and 45.00 g/0.25 m²) compared to 15×15 cm (12.92 g/clump and 23.04 g/0.25 m²). On the contrary, the grain dry weight per hectare was higher in 15×15 cm (5.74 t/ha) than in 25×25 cm (3.10 t/ha). The research findings indicate that closer plant spacing can suppress weed growth and increase the grain dry weight per hectare compared to wider plant spacing. We recommend that the use of cogon grass extract dosage needs to be increased above 50 L/ha.

Key words: Bioherbicide, cogon grass extract, plant spacing, rice, weed

INTRODUCTION

Rice plants are the main producers of rice to meet the food needs of the majority of Indonesians. The use of superior varieties and short-lived ones is the right choice to fulfill national food needs. As the population increases, the demand for rice continues to rise. However, every rice cultivation is always accompanied by the emergence of weeds around the plants. The presence of weeds significantly disrupts and harms rice growth and yield. So it must be controlled.

Many methods of weed control with chemicals have been recommended, but biological control with cogon grass (*Imperata cylindrica* (L.) Beauv.) extract is not widely known. The cogon grass extract as a bioherbicide contains allelopathic compounds (Kato-Noguchi, 2022). These compounds can inhibit the germination of weed seeds in the soil by hindering enzyme activity through the degradation of food reserves in the seeds. As a result, the energy produced is low due

to a decrease in germination power. In addition, weed growth control can also be achieved through plant spacing.

Closer plant spacing is a method of weed control through technical cultivation. The plant canopy plays a role in reducing the distribution of sunlight to the soil surface. As a result, weed growth is inhibited due to a lack of light (Listyowati *et al.*, 2022). The suitable plant spacing can enhance rice yield per hectare and suppress weed growth. In fertile land, a more spacious plant spacing is advisable, while in less fertile land, a closer spacing is recommended. Short-lived varieties are better suited for close spacing, whereas long-lived varieties are suitable for wider spacing. The use of superior rice varieties requires a looser plant spacing. Increasing plant spacing provides weeds with the opportunity to grow freely and subsequently interfere with the crop. Therefore, understanding the suitable plant spacing is essential.

Weeds are troublesome plants that adversely affect the growth and yield of plants. Extracts from certain weeds have the potential as bioherbicides because they can produce allelopathy (Erida *et al.*, 2019). Allelopathy is capable of reducing seed germination and slowing down the germination process. Allelopathic compounds can inhibit the activity of enzymes that degrade food reserves in seeds, resulting in very low energy production and decreasing germination potential.

In general, plant-derived bioherbicides contain phytotoxic allelochemicals or specific disease-carrying microbes (Hasan *et al.*, 2021). The advantages of using bioherbicides are that they contain no toxic organic secondary metabolites, that they are easy to acquire from the field, that they have no secondary metabolite mechanisms found in synthetic herbicides, that they have more than one secondary metabolite compound, and that they do not cause plant poisoning (Sihombing *et al.*, 2018). When bioherbicides are absorbed by weed seeds or roots, they disrupt the integrity of cell membranes and other essential biochemical processes. It should be noted that the cytotoxic effect on weeds is reflected in low levels of root cell division, low ability to absorb nutrients, growth hormones, and pigment synthesis, as well as low growth of reactive oxygen species (ROS), stress-related hormones, and abnormal antioxidant activity (Hasan *et al.*, 2021).

One type of weed with potential as a bioherbicide is *Imperata cylindrica* (Sari *et al.*, 2017). The roots of cogon grass can be used as a pre-emergence bioherbicide. Cogon grass root extract contains allelochemical compounds, specifically flavonoids. A concentration of 1% root extract of cogon grass is capable of inhibiting the growth of weed seeds located below the ground (Sari *et al.*, 2017). Furthermore, a concentration of 14% extract from cogon grass rhizomes can effectively

control weeds such as *Ageratum conyzoides*, *Eleusine indica*, and *Cyperus rotundus* (Lau *et al.*, 2021).

Extract cogon grass contains terpenoid, phenolic, and steroid compounds (Erida *et al.*, 2019). The extracts, leachates, root exudates, or growing medium of cogon grass also contained several allelochemicals, including phenolic acids, phenolic aldehydes, phenylpropanoids, flavonoids, quinones, fatty acids, terpenoids, simple phenolics, benzoic acids, and alkaloids (Kato-Noguchi, 2022). The application of cogon grass extract at the time of planting with a concentration of 30% could increase the number of panicles and the weight of dry rice grains in rice cultivation in polybags. It had a greater potential as a bioherbicide to suppress weed growth and increase rice yield (Paiman *et al.*, 2022).

The percentage of germination, germination rate, and seedling length of weed seeds such as *Amaranthus spinosus*, *Bidens biternata*, and *Tridax procumbens* could all be greatly suppressed by using cogon grass extract. *Tridax procumbens* had the highest level of sensitivity among weed seeds to the extract from the roots and rhizomes of cogon grass, followed by *Bidens biternata* and *Amaranthus spinosus*. Extracts from the roots and rhizomes of cogon grass exhibit significant potential as bioherbicides to inhibit the germination of weed seeds (Pujiwati, 2011). On the other hand, using close plant spacing can also suppress weed growth and increase yield per unit area. But, grain yield per clump decreases. The appropriate plant spacing can provide plants with the opportunity to grow well without competition from neighbouring plants for water, nutrients, and sunlight. The suitable plant spacing can enhance rice yield. Also, the use of plant spacing depends on the cultivated rice variety.

Rice plants with curved panicles were used for planting density was not too tight and not too loose, but rice plants with erect panicles were better with planting density quite tight (Yena *et al.*, 2018). Wide inter-row width and spacing in narrow rows will be conducive to building a high-yield population to increase yields (Li *et al.*, 2016). If the spacing between rows was fixed at 30 cm, then the spacing in rows of 11-12 cm could increase plant population, and grain yield, and was the optimal plant spacing for rice cultivation (Yun *et al.*, 2020). Grain yield was maintained in a large number of plant populations, primarily through the effects of more tillers per plant and more grain per panicle in lower plant populations (Dunn *et al.*, 2020).

Plant spacing significantly influenced plant height, tiller number, and rice panicle length. A plant spacing of 15 × 15 cm provided the highest panicle count and good yield per square meter

(Marie-Noel *et al.*, 2021). The optimal Jajar Legowo system involved skipping one row after every three rows at plant spacing of 15 × 15 cm, resulting in the highest grain yield (Rautaray, 2016). Consistent with subsequent research, the highest grain dry weight was found at a plant spacing of 15 × 15 cm compared to 20 × 20 cm and 25 × 25 cm (Rautaray, 2016). In line with further studies, plant density with a plant spacing of 15 × 15 cm resulted in the highest grain yield (4.71 t/ha), while the 25 × 25 cm treatment yielded a lower amount (4.09 t/ha) (Alogaidi *et al.*, 2019). Also supported by Faisul-ur-Rasool *et al.* (2012), closer spacing (15×15 cm) produced the maximum tillers number, leaf area index, dry matter accumulation, and rice yield. Plant spacing of 15 × 15 cm resulted in a yield of 8.97% greater than 20 × 20 cm because of better absorption of photosynthetically active light.

Plant population had a significant effect on morphological and physiological characteristics as well as crop yield. The leaf area per clump was broader with a looser plant spacing. Additionally, wider plant spacing results in lower competition among plants, allowing them to acquire more nutrients. As a result, there were more tillers, a larger leaf area, and greater dry weight per cluster, as opposed to dense plant spacing (Mondal *et al.*, 2013). There was limited information available on the optimal row width needed to maximize grain yield potential (Dunn *et al.*, 2020). Superior rice varieties prefer a looser plant spacing for their growth compared to local varieties, but yield per unit area needs to be maximized using closer plant spacing.

Based on the literature review above, cogon grass extract as a bioherbicide has proven to be quite effective in biologically controlling weeds, particularly at a concentration of 30%. However, this bioherbicide has not yet been directly applied in rice fields, and the appropriate application timing in rice cultivation remains unknown. Additionally, it has been established from several previous studies that the use of close plant spacing (15 × 15 cm) in rice cultivation can effectively suppress weed growth and increase yield. However, the use of plant spacing has not been investigated in the cultivation of the short-lived rice of the Padjajaran Agritan variety.

Research about the application timing of cogon grass extract as a bioherbicide and the plant spacing to suppress weed growth and increase rice yield needs to be conducted. This study is expected to contribute insights into environmentally friendly weed control through biological and technical cultural approaches. Therefore, this research aimed to know the effect of application timing of cogon grass extract and plant spacing on weed growth in rice cultivation of the Padjajaran Agritan variety.

MATERIALS AND METHODS

Study Site

The research was conducted from October 2022 to February 2023 during the rainy season. The study was located in Minggir Sub-district, Sleman Regency, Special Region of Yogyakarta, Indonesia. The elevation of the location was 110 m above sea level. Geographically, Sleman Regency is situated between 110°33' 00" - 110°13' 00" East Longitude and 7°34'51" - 7°47'30" South Latitude.

Experimental Design

The study was 3 × 2 factorial and arranged in a randomized completely block design (RCBD). The first factor was the application timing of bioherbicide consisting of three methods: control (no application), three days before planting, and at the time of planting. The second factor was the use of plant spacing, consisting of two kinds: 15 × 15 cm (close plant spacing) and 25 × 25 cm (wide plant spacing). In total, there were six treatment combinations. Each treatment combination was repeated three times (replications as blocks), resulting in a total of 18 treatment plots. Treatment plots of plant spacing of 15 × 15 cm and 25 × 25 cm can be seen in Fig. 1.

Research Procedures

In this study, seeds of the Padjajaran Agritan variety were used. Before sowing, the seeds were soaked in water for three hours to ensure uniform imbibition. Subsequently, the seeds were lifted from the soaking medium, and the excess water was drained. After one day and one night, roots emerged from the base of the seeds. Plastic seedling trays measuring 50 × 60 cm with a height of 10 cm were used for seed germination. The germination medium consisted of a mixture of organic fertilizer and soil in a 1:1 ratio. The seeds were then sown on the surface of the germination medium at field capacity. Four days later, shoots emerged above the soil surface, and the seedlings were ready to be transplanted into the rice field at the age of 18 days after sowing.

Soil processing was conducted in two stages. The first stage of soil processing involved using a rotary plow to overturn and crush the soil. The second stage of processing utilized a harrow to level the soil. Subsequently, treatment plots were established with dimensions of 3 × 3 m, totaling 18 treatment plots. Drainage channels with a width of 0.5 m were created between the treatment plots. Treatment randomization was performed per block for all blocks.

In this research, cogon grass extract was used as a bioherbicide. The application timing of bioherbicide corresponds to the randomized treatment results. The control plot was without

bioherbicide treatment, while the treatment plots involved the use of a 30% concentration of bioherbicide with a dosage of 45 ml/3 m² (50 L/ha). The application timing of bioherbicide was divided into two methods: three days before planting and at the time of planting. Bioherbicide was applied only once for each treatment in research. During the three days following the treatment, measures were taken to ensure that no external water entered the treatment plots.

The seedlings number that planted in each treatment plot depends on the plant spacing. For a plant spacing of 15 × 15 cm, 20 seedlings were planted in rows and 20 seedlings between rows, requiring a total of 400 seedlings per plot (Figure 1a). Meanwhile, for a plant spacing of 25 × 25 cm, 12 seedlings were planted in rows and 12 seedlings between rows, requiring a total of 144 seedlings per plot (Figure 1b). Only one rice seedling was planted per planting hole.

Recommended fertilizer usage was 225 kg/ha of urea and 225 kg/ha of NPK Phonska 15-15-15 (BPPP, 2014). Fertilizer application was divided into two stages. In the first stage, 40% of the recommended dosage was applied 15 days after planting (DAP). The second stage involved applying 60% of the recommended dosage at 35 DAP. Irrigation was carried out every four days after the treatment. Pest and disease control were implemented during the rice flowering stage to minimize the impact of brown planthoppers. Harvesting was conducted when the rice plants reached 104 DAP.

Parameters

Observations on weed growth were conducted at 60 days after transplanting (DAT) and included the number of weeds (individual/0.25 m²), weed dry weight (g/0.25 m²), weed dominance, and weed community coefficient (%). Observations on rice plants were performed on growth and yield components, including the tillers number (stalks/clump) observed at 60 DAT, plant dry weight (g/clump), panicles number (strands/clump), grain dry weight (g/clump), and hectare (ton/ha), all observed at 104 DAT.

Statistical Analysis

Observational data were analyzed by analysis of variance (ANOVA) at P = 0.05 probability levels with the software IBM SPSS Statistic 23. In addition, Duncan's new multiple range test (DMRT) at P = 0.05 probability levels was used to test the difference between treatment averages. To determine the dominant weed species and weed community sift, the summed dominance ratio (SDR) and coefficient of weed community were used.

RESULTS AND DISCUSSION

Number and Dry Weight of Weeds

The timing of the bioherbicide application did not have a significant effect on the number and dry weight of weeds. However, only the plant spacing had an impact on the weed dry weight. The average number and dry weight of weeds can be seen in Table 1.

Table 1 shows that a plant spacing of 25×25 cm resulted in a higher weed dry weight compared to 15×15 cm. Wider plant spacing contributed to a heavier weed dry weight. In wider plant spacing, sunlight penetration can reach the soil surface, and then support weed growth. The weed dry weight at a plant spacing of 15×15 cm was lower than that at 25×25 cm. It indicates that closer plant spacing is more effective in suppressing weed growth because the plant canopy quickly covers the soil surface. Sunlight penetration to the soil surface is lower, so inhibits weed growth. In line with Fahad *et al.* (2015), less weed growth was seen at narrow row spacing (11 cm) compared to wider row spacing (15 and 23 cm) in wheat crops. In line with the research result by Chauhan and Johnson (2010), narrowing the row spacing could decrease weed growth and seed production, as well as increase the grain yield of rice.

Dominance of Weed Species

In this study, three types of weeds were identified based on their morphology, namely sedges, grasses, and broad-leaved weeds, with a total of nine weed species overall. The effect of the combination of bioherbicide application timing and plant spacing on the SDR of each weed species can be seen in Table 2.

Table 2 indicates that the identified sedges included *Cyperus brevifolius*, *Cyperus rotundus*, and *Scorpus mucronatus*. The grass weeds identified in this study were *Cynodon dactylon*, *Echinochloa colona*, *Echinochloa crus-galli*, and *Fimbristylis miliacea*. Several types of broad-leaved weeds found in this research were *Limnocharis flava* and *Monochoria vaginalis*. *Fimbristylis miliacea* was the dominant weed species followed by *Echinochloa crus-galli* in this study. It can be explained that grass weeds more dominate in rice cultivation than other types of weeds. It was supported by Kurniadie *et al.* (2019), *Fimbristylis miliacea* was a dominant weed species in conventional and organic rice cultivation.

Weed Community Shift

Based on Table 2, the weed community coefficient (C%) can be calculated between weed communities, both among the application timing of bioherbicide and plant spacing treatments. The results of the weed community coefficient are presented in Tables 3 and 4.

Table 3 indicates that the coefficient of weed community in the comparison between the control and application timing of bioherbicide at the time of planting was 66.69% in a plant spacing of 15×15 cm. It shows a shift in the weed species composition due to the application timing of bioherbicide at the time of planting. In other words, there is a difference in the weed species that grow in these two communities. Meanwhile, for the other communities, there is no shift in the weed species community because their community coefficients were $>75\%$.

Growth and Yield of Rice

Observations on the growth and yield of rice included the tillers number, plant dry weight, panicles number, grain dry weight per clump, and hectare. The application timing of bioherbicide did not have a significant effect on all the observed parameters. However, the plant spacing significantly influenced all the observed parameters. The average tillers number, plant dry weight, panicles number, grain dry weight per clump, and hectare can be seen in Table 5.

Table 4 indicates that a plant spacing of 25×25 cm resulted in tillers number (12.51 stem), plant dry weight (14.91 g), panicles number (10.71 strands), and grain dry weight (19.40 g) per clump, respectively. Plant spacing of 25×25 cm was given growth and yield component per clump higher than in 15×15 cm. The higher tillers number, plant dry weight, and panicles number at wider plant spacing correlated with the grain dry weight per clump. The result of this study was supported by Dass *et al.* (2017), wider plant spacing significantly increased rice yield compared with closer plant spacing. In line with Anwari *et al.* (2019), yield components like tillers number and grain per panicle were found in plant spacing 25×25 cm compared to 10×10 cm, 15×15 cm, or 20×20 cm.

A plant spacing of 25×25 cm could yield higher grain dry weight per clump because of a better microclimate compared to closer plant spacing. It caused lower competition among plants for nutrients, improved water access, and smoother air circulation. Rice plants could grow well so that the size of the habitus was larger. Additionally, sunlight distribution was more even across the entire plant body, leading to enhanced photosynthesis. These conditions resulted in a more efficient carbohydrate-filling process for all seeds in the panicle. With a greater seed number per clump and

a heavier weight per seed, the total grain dry weight was higher. On the other hand, sunlight penetration to the soil surface could benefit weeds to support their growth.

However, a plant spacing of 15×15 cm could provide a higher grain dry weight per hectare compared to 25×25 cm. In closer plant spacing, the population density per hectare was higher, leading to a higher yield per hectare. The highest radiation use efficiency was obtained at close plant spacing which resulted in a higher amount of biomass for energy production. Following the opinion of Ya-Jie *et al.* (2016), higher density was applied to increase panicles and grains number for higher yield per unit area. Supported by Marie-Noel *et al.* (2021), the highest panicle number/m² was found in plant spacing of 15×15 cm but not obtained in 20×20 cm, 25×25 cm, or 30×30 cm. In contrast with the study results by Asghar *et al.* (2021), the highest grain yield of 6.75 t/ha was obtained in plant spacing of 23×23 cm followed by 5.46 t/ha from 18×18 cm. No in line with the study results by Neupane (2022), loose plant spacing (20×20 cm) was given superior morphophysiological and yield components and resulted in a grain yield of 8.53 t/ha higher than 6.47 t/ha in tight plant spacing (20×10 cm).

The use of a plant spacing of 15×15 cm resulted in plant populations higher than 25×25 cm, with 444,444 plants/ha higher than 160,000 plants/ha. Total grain yield per hectare could be increased by close plant spacing. The plant population at a plant spacing of 15×15 cm was almost three times higher than 25×25 cm. The panicles number per hectare was found to be higher at a plant spacing of 15×15 cm which was 3,444,441 panicles greater than 1,713,600 panicles in 25×25 cm. It means that a greater panicle number could support the total grain dry weight per hectare, although the total grain dry weight per panicle was lower. Deng *et al.* (2022) explained that increasing light utilization by the canopy was a major way to increase photosynthesis and rice productivity.

In this study, the Padjajaran Agritan variety was used, which has short-lived growth, fewer tillers, and a smaller plant size. Therefore, using a closer plant spacing was the best solution to achieve a higher yield per hectare by intercepting more solar radiation. It can be concluded that using a closer plant spacing for rice varieties with a compact habitus is more advantageous because it can capture and harness more sunlight penetration. Additionally, the growth of weeds under the plant canopy can be inhibited.

CONCLUSION

In conclusion, the the application timing of bioherbicide did not significantly effect on growth of weed and rice, as well as the grain dry weight. *Fimbristylis miliacea* was obtained as the dominant weed species. Bioherbicide application at the time of planting could alter the composition of weed species compared to the control in plant spacing of 15 × 15 cm. Furthermore, the plant spacing of 25 × 25 cm resulted in higher grain and weed dry weight (19.40 g/clump and 45.00 g/0.25 m²) compared to 15 × 15 cm (12.92 g/clump and 23.04 g/0.25 m²). On the contrary, the grain dry weight per hectare was higher in 15 × 15 cm (5.74 t/ha) than in 25 × 25 cm (3.10 t/ha). The research findings indicate that closer plant spacing can suppress weed growth and increase the grain dry weight per hectare compared to wider plant spacing. We recommend that the use of cogon grass extract dosage needs to be increased above 50 L/ha.

ACKNOWLEDGEMENTS

We would like to thank the Institute for Research and Community Service at Universitas PGRI Yogyakarta for funding and supporting this research. Furthermore, we would like to thank the Faculty of Agriculture, Universitas PGRI, which has provided financing for greenhouse facilities and equipment needed for research.

REFERENCES

- Alogaidi, F. F., Al-shugeairy, Z. K., Hadi, B. H. and Hassan, W. A. (2019). Effect of planting distance on yield and yield components of four effect of planting distance on yield and yield components of four introduced upland rice varieties. *Plant Arch.* **19**: 699–707. [HTTPS://PLANTARCHIVES.ORG/PDF%2019-1/699-707%20\(4827\).PDF](https://plantarchives.org/pdf%2019-1/699-707%20(4827).pdf)
- Anwari, G., Moussa, A. A., Wahidi, A. B., Mandozai, A., Nasar, J. and El-Rahim, M. G. M. A. (2019). Effects of planting distance on yield and agro-morphological characteristics of local rice (Bara variety) in Northeast Afghanistan. *Curr. Agric. Res. J.* **7**: 350–7. [HTTPS://DOI.ORG/10.12944/CARJ.7.3.11](https://doi.org/10.12944/CARJ.7.3.11)
- Asghar, M., Hassan, T., Arshad, M., Aziz, A., Latif, M. T. and Sabir, A. M. (2021). Effect of plant spacing on incidence of rice planthoppers in transplanted rice crop. *Int. J. Trop. Insect. Sci.* **41**: 575–85. [HTTPS://DOI.ORG/10.1007/s42690-020-00242-4](https://doi.org/10.1007/s42690-020-00242-4)
- Chauhan, B. S. and Johnson, D. E. (2010). Implications of narrow crop row spacing and delayed *Echinochloa colona* and *Echinochloa crus-galli* emergence for weed growth and crop yield loss in aerobic rice. *F. Crop. Res.* **117**: 177–82. [HTTPS://DOI.ORG/10.1016/J.FCR.2010.02.014](https://doi.org/10.1016/j.fcr.2010.02.014)

- Dass, A., Chandra, S., Uphoff, N., Choudhary, A. K., Bhattacharyya, R. and Rana, K. S. (2017). Agronomic fortification of rice grains with secondary and micronutrients under differing crop management and soil moisture regimes in the North Indian Plains. *Paddy Water Environ.* **15**: 745–60. [HTTPS://LINK.SPRINGER.COM/ARTICLE/10.1007/S10333-017-0588-9](https://link.springer.com/article/10.1007/s10333-017-0588-9)
- Deng, F., Li, B., Yuan, Y., He, C., Zhou, X., Li, Q., Zhu, Y., Huang, X., He, Y., Ai, X., Tao, Y., Zhou, W., Wang, L., Cheng, H., Chen, Y., Wang, M. and Ren, W. (2022). Increasing the number of seedlings per hill with reduced number of hills improves rice grain quality by optimizing canopy structure and light utilization under shading stress. *F. Crop. Res.* **287**: 108668. [HTTPS://DOI.ORG/10.1016/J.FCR.2022.108668](https://doi.org/10.1016/j.fcr.2022.108668)
- Dunn, B. W., Dunn, T. S., Mitchell, J. H. and Brinkhoff, J. (2020). Effects of plant population and row spacing on grain yield of aerial-sown and drill-sown rice. *Crop. Pasture Sci.* **71**: 219–28. [HTTPS://DOI.ORG/10.1071/CP19421](https://doi.org/10.1071/CP19421)
- Erida, G., Saidi, N., Hasanuddin, and Syafruddin. (2019). Allelopathic screening of several weed species as potential bioherbicides. The 3rd CSAFS. *IOP Conf. Series: Earth Environ. Sci.* **334**: 012034. [HTTPS://DOI.ORG/10.1088/1755-1315/334/1/012034](https://doi.org/10.1088/1755-1315/334/1/012034)
- Fahad, S., Hussain, S., Chauhan, B. S., Saud, S., Wu, C., Hassan, S., Tanveer, M., Jan, A. and Huang, J. (2015). Weed growth and crop yield loss in wheat as influenced by row spacing and weed emergence times. *Crop. Prot.* **71**: 101–8. [HTTPS://DOI.ORG/10.1016/J.CROPRO.2015.02.005](https://doi.org/10.1016/j.cropro.2015.02.005)
- Faisal-ur-Rasool, Habib, R. and Bhat, M. I. (2012). Evaluation of plant spacing and seedlings per hill on rice (*Oryza sativa* L.) productivity under temperate conditions. *Pakistan J. Agric. Res.* **49**: 169–72.
- Hasan, M., Ahmad-hamdani, M. S., Rosli, A. M. and Hamdan, H. (2021). Bioherbicides: An eco-friendly tool for sustainable weed management. *Plants.* **10**: 1212. [HTTPS://DOI.ORG/10.3390/PLANTS10061212](https://doi.org/10.3390/plants10061212)
- Kato-Noguchi, H. (2022). Allelopathy and allelochemicals of *Imperata cylindrica* as an invasive plant species. *Plants.* **11**: 2551. [HTTPS://DOI.ORG/10.3390/PLANTS11192551](https://doi.org/10.3390/plants11192551)
- Kurniadie, D., Irda, M., Umiyati, U., Widayat, D. and Nasahi, C. (2019). Weeds diversity of lowland rice (*Oryza sativa* L.) with different farming system in Purwakarta Regency Indonesia. *J. Agron.* **18**: 21–6.
- Lau, D. F. W., Sofian, and Mirza, A. (2021). Ekstrak rimpang alang-alang (*Imperata cylindrica* L.) sebagai herbisida nabati untuk mengendalikan gulma. *J. Agroekoteknologi Trop. Lembab.*

4: 29–34. (in Indonesian)

Li, G., Zhang, J., Yang, C., Liu, Z., Wang, S. and Ding, Y. (2016). Population characteristics of high-yielding rice under different densities. *Agron. J.* **108**: 1415–23. [HTTPS://DOI.ORG/10.2134/AGRONJ2015.0214](https://doi.org/10.2134/AGRONJ2015.0214)

Listiyowati, C., Indradewa, D. and Irwan, S. N. R. (2022). Study on weeds abundance on rice fields in mycorrhizal inoculation and different planting methods. *Caraka Tani*, **37**: 259–274. <https://doi.org/10.20961/carakatani.v37i2.53131>

Marie-Noel, M. F., Joseph, B. M., Liliane, T., Felix, F., Dorothy, M. and Salvador, D. P. (2021). Effect of plant spacing on the growth and yield of rainfed rice (*Oryza sativa*) in the bimodal rain forest zone of Cameroon. *J. Agric. Crop.* **7**: 48–59. [HTTPS://DOI.ORG/10.32861/JAC.72.48.59](https://doi.org/10.32861/JAC.72.48.59)

Mondal, M. M. A., Puteh, A. B., Ismail, M. R. and Rafii, M. Y. (2013). Optimizing plant spacing for modern rice varieties. *Int. J. Agric. Biol.* **15**: 175–8.

Neupane, P. (2022). Effect of plant spacing and genotype on growth, yield, and yield-related traits of rice varieties. *Russ. J. Agric. Socio-Economic Sci.* **3**: 89–98. [HTTPS://DOI.ORG/10.18551/RJOAS.2022-03.10](https://doi.org/10.18551/RJOAS.2022-03.10)

Paiman, Hidayat, A. K., Shobirin, S. S. and Khasanah, S. I. (2022). Efficacy of weed extract as a bioherbicide in rice (*Oryza sativa* L.) cultivation. *Res. Crop.* **23**: 488–96.

Pujiwati, I. (2011). Land use potential through resds (*Imperata cylindrica*) as bioherbicide. *Gea.* **11**: 226–34.

Rautaray, S. K. (2016). Effect of planting geometry and fertilizer dose on grain yield and disease incidence in some mid duration rice (*Oryza sativa* L.) varieties. *Arch. Agron. Soil Sci.* **53**: 327–33. [HTTPS://DOI.ORG/10.1080/03650340701223320](https://doi.org/10.1080/03650340701223320)

Sari, V. I., Nanda, S. and Sinuraya, R. (2017). Bioherbisida pra-tumbuh alang-alang (*Imperata cylindrica*) untuk pengendalian gulma di perkebunan kelapa sawit. *J. Citra Widya Edukasi.* **9**: 301–8. (in Indonesian)


Sihombing, N., Purba, Z., Samosir, S. and Karim, S. (2018). Potensi metabolit sekunder gulma sebagai pestisida nabati di Indonesia (Potency of secondary metabolite from weeds as natural pesticides in Indonesia). *J. Kultiv.* **17**: 683–93. (in Indonesian)

Ya-Jie, H., Hai-Jun, Q., Wei-Wei, C., Zhi-Peng, X., Hong-Cheng, Z., Qi-Gen, D., Zhong-Yang, H., Ke, X., Hai-Yan, W. and Bao-Wei, G. (2016). Effect of different mechanical transplantation methods and density on yield and its components of different panicle-typed


ricechinese. *Chinese J. Rice Sci.* **2016**: 493–506. [HTTPS://DOI.ORG/10.16819/J.1001-7216.2016.6015](https://doi.org/10.16819/j.1001-7216.2016.6015)

Yena, W., Xiang, Z., Guotao, Y., Xuechun, W., Youlin, P., Yao, H. and Yungao, H. (2018). Effects of planting density on the microclimate of rice population with different panicle-types and its yield components. *Chinese J. Appl. Environ. Biol.* **2018**: 813–23. [HTTPS://DOI.ORG/10.19675/J.CNKI.1006-687X.2017.12049](https://doi.org/10.19675/j.cnki.1006-687x.2017.12049)

Yun, C., Yajun, Z., Honglu, Z., An, Z., Jian, H., Hao, Z., Junfei, G., Lijun, L., & Jianchang, Y. (2020). Effects of plant spacing on grain yield and population quality in mechanically-transplanted rice with good tasting quality. *Chinese J. Rice Sci.* **2020**: 550–60. [HTTPS://DOI.ORG/10.16819/J.1001-7216.2020.0205](https://doi.org/10.16819/j.1001-7216.2020.0205).



c. Plant spacing of 15 × 15 cm (400 seedlings)



d. Plant spacing of 25 × 25 cm (144 seedlings)

Fig. 1. Plot treatments of plant spacing of 15 × 15 cm (a) and 25 × 25 cm (b).

Table 1. Effect of bioherbicide application and plant spacing on the weed growth per plot.

Treatments	Parameter of observations	
	Weed number (individual/0.25 m ²)	Weed dry weight (g/0.25 m ²)
Timing of bioherbicide application:		
Control	58.17 a	32.08 a
Three days before planting	41.67 a	28.97 a
At the time of planting	44.67 a	41.01 a
Plant spacing (cm):		
15 × 15	41.22 p	23.04 q
25 × 25	55.11 p	45.00 p

Remarks: According to DMRT at P = 0.05 probability levels, there is no significant difference between the numbers in the same column followed by the same characters.

Table 2. Summed dominance ratio (SDR) (%) of each weed species.

No.	Weed species	SDR (%)					
		J ₁ B ₀	J ₁ B ₁	J ₁ B ₂	J ₂ B ₀	J ₂ B ₁	J ₂ B ₂
1	<i>Cyperus brevifolius</i> (Rottb.) Rassk.	0.00	2.56	0.00	0.00	0.00	0.00
2	<i>Cyperus rotundus</i> L.	0.00	0.00	13.58	10.86	5.26	4.05
3	<i>Scirpus mucronatus</i> L.	10.72	6.29	4.06	6.35	0.00	3.24
4	<i>Cynodon dactylon</i> (L.) Pers.	4.92	7.57	10.51	12.45	5.50	8.10
5	<i>Echinochloa colona</i> (L.) Link.	5.55	9.70	16.75	6.73	4.27	5.94
6	<i>Echinochloa crus-galli</i> (L.) P. Beauv.	18.90	8.49	0.00	12.37	27.1	9.42
7	<i>Fimbristylis miliacea</i> (L.) Vahl.	47.48	50.54	44.34	51.21	50.3	52.26
8	<i>Limnocharis flava</i> (L.) Buch	0.00	7.87	2.91	0.00	0.00	6.66
9	<i>Monochoria vaginalis</i> (Burm.f.) C.Presl.	12.39	6.95	7.82	0.00	7.49	10.29

Remarks: J₁ = Plant spacing of 15 × 15 cm, J₂ = Plant spacing of 25 × 25 cm, B₀ = Control, B₁ = Three days before planting, and B₂ = At the time of planting.

Table 3. Weed community coefficient between application timing of bioherbicide in each plant spacing.

Plant spacing (cm)	Comparison between application timing of bioherbicide	Weed community coefficient (C%)	Information
15 × 15	4. Control and three days before planting	79.68	Homogeny
	5. Control and at the time of planting	66.69	No homogeny
	6. Three days before planting and at the time of planting	75.53	Homogeny
25 × 25	4. Control and three days before planting	77.70	Homogeny
	5. Control and at the time of planting	81.96	Homogeny
	6. Three days before planting and at the time of planting	81.03	Homogeny

Table 4. Weed community coefficient between plant spacing in application time of bioherbicide



Application timing of bioherbicide	The comparison between plant spacing	Weed community coefficient (C%)	Information
Control	15 × 15 and 25 × 25 cm	76.67	Homogeny
Three days before planting	15 × 15 and 25 × 25 cm	75.51	Homogeny
At the time of planting	15 × 15 and 25 × 25 cm	76.40	Homogeny

Table 5. The effect of application timing of bioherbicide and plant spacing on the growth and yield of rice.



Treatments	Parameters				
	Tillers number (stem/clump)	Plant dry weight (g/clump)	Panicles number (strands/clump)	Grain dry weight (g/clump) (t/ha)	
<u>Application timing of bioherbicide</u>					
Control	10.57 a	9.97 a	9.06 a	14.90 a	4.15 a
Three days before planting	11.37 a	13.93 a	9.13 a	16.58 a	4.32 a
At the time of planting	10.73 a	12.94 a	9.50 a	17.00 a	4.80 a
<u>Plant spacing (cm)</u>					
15 × 15	9.27 q	10.75 q	7.75 q	12.92 q	5.74 p
25 × 25	12.51 p	14.91 p	10.71 p	19.40 p	3.10 q

Remarks: According to DMRT at P = 0.05 probability levels, there is no significant difference between the numbers in the same column followed by the same characters.

Tanggapan dari editor: 22 Maret 2024



Approved for Publication in Research on Crops: 9 April 2024



Konfirmasi dan Bukti Pembayaran: 20 April 2024

CONFIRMATION OF PAYMENT OF THE PUBLICATION FEE

Dear
Managing Editor

I already paid for this article publication fee with no. ROC-1057 entitled "**Optimizing herbicide application Timing and Plant Spacing to Manage Weed Growth in Rice (*Oryza sativa* L.) Cultivation** [Approved for publication in RESEARCH ON CROPS Vol. 25, No. 2 (June) 2024]". Sender on behalf of Ari Kusuma Wardana, Universitas PGRI Yogyakarta, Indonesia through Bank Name: BNI, Indonesia, Account Name: Ari Kusuma Wardana, Account number : 0723567241. Proof of payment transfer to State Bank of India, Patel Nagar, Hissar-125001, Haryana, India. Beneficiary account name: Gaurav Publications, and beneficiary account number: 37014428711. Here, I attached proof of payment.

Sincerely,
Paiman
Department of Agrotechnology
Faculty of Agriculture
Universitas PGRI Yogyakarta
Yogyakarta 55182, Indonesia

BNI Tanggal/Date: 10/04/2024 **Formulir Kiriman Uang**
Remittance Application

Perorangan/Personal Penduduk/Resident Perusahaan/Company Bukan Penduduk/Non Resident

Tunai/Cash Cek/BG No. Debit Rek./Debit Acc. No. RTGS Draft SWIFT

Jenis Pengiriman/
Type of Transfer

Sumber Dana/ Source of fund

Mata Uang/Currency: IDR USD

Jumlah Dana yang dikirim/Amount Transfer:

Jumlah/Amount	Kurs/Rate	Nilai/Total Amount
900		

Biaya/Charge

Nilai/Amount in Foreign Exchange	Kurs/Amount	Nilai/Total Amount

Komisi/Commission Pengiriman/Handling Bank Koresponden/Correspondent Bank

Jumlah Biaya/Amount Charge:

Total yang dibayarkan/Total Amount:

Terbilang/Amount in Words: Nine Hundred dollar

Tujuan Transaksi (Transaction Purpose): Article Payment

Berita (Message): Optimizing Herbicide Application

Biaya dari Bank koresponden dibebankan ke rekening/
Correspondent bank charges are for account of:

Penerima/Beneficiary Pengirim/Remitter Sharing

Pejabat Bank/Bank Officer: Teller

Pemohon/Applicant: Ari Kusuma Wardana

Saya menyetujui sepenuhnya syarat-syarat yang tercantum pada halaman belakang formulir ini / I unconditionally accept all the terms and conditions on the reverse form.

BANK INDONESIA (PERBANK) Tbk.

Lembar 1 & 2 : w/ Bank; Lembar 3 : w/ Nasabah

BNI 215 537013 021031 AFMT 19/04/2024 12:01:01
 140482010001 **Formulir Pemindahbukuan**
 : USD 25,00 NPD MISC. DE

Penerima
 Nama: *Pend. Restitusi biaya CBK USD*
 Nomor Rekening: *3840482010001*
 Nama Cabang: _____

Pengirim
 Nama: *Ari Kusuma Wardana*
 Nomor Rekening: *0920567241*
 Nama Cabang: _____

Jumlah: *25 USD*
 Terbilang: *dua puluh lima US dollar*
 Biaya: _____

Keterangan: *Biaya CBK 51076Y00070224*

Pejabat Bank _____ **Teller** _____ **Pemohon** _____

Tujuan Transaksi (Transaction Purpose): *Akuntabel*
 Berita (Message): *0920567241*
 Biaya dari Bank koresponden dibebankan ke rekening/
 Correspondent bank charges are for account of:
 Penerima/Beneficiary Pengirim/Remitter Sharing

Pejabat Bank/Bank Officer _____ **Teller** _____ **Pemohon/Applicant** _____

Sila sila ada catatan atau komentar atau laras mengenai form permohonan

Saya menyetujui sepenuhnya syarat-syarat yang tercantum pada seluruh halaman formulir ini / I unconditionally accept the terms and conditions on the reverse side.

Bank Indonesia (BI) is a member of the Financial Action Task Force (FATF) and is committed to the FATF Recommendations to prevent and combat money laundering. Bank Indonesia is also a member of the Global Financial Anti-Corruption Centre (GFACC) and is committed to the GFACC Recommendations to prevent and combat corruption.

REFRENSI : 51076Y00070224

NO. TRX : 83215 931161 96963 TRXN 19/04/2024 11:48:10
 NO. REK : 000000723567241 SDR ARI KUSUMA WARDANA
 JUMLAH : IDR 30,000 1568
 004 YOGYAKARTA

NO. TRX : 83215 931161 96963 TRXN 19/04/2024 11:48:10
 NO. REK : 004840200101001 PENEMPATAN PROFISI RU
 JUMLAH : IDR 30,000 1568
 004 YOGYAKARTA

NO. TRX : 81215 931161 96963 TRXN 19/04/2024 11:48:10
 NO. REK : 000000723567241 SDR ARI KUSUMA WARDANA
 JUMLAH : IDR 14,785,500 1568
 004 YOGYAKARTA

NO. TRX : 83215 931161 96963 TRXN 19/04/2024 11:48:10
 NO. REK : 004840200101001 RI YAKIN
 JUMLAH : USD 900 1568
 004 YOGYAKARTA

BNI
 BANK NEGARA INDONESIA (PERSERO) TBK
 KANTOR CENTRAL
 YOGYAKARTA

PT. BANK NEGARA INDONESIA (Persero), Tbk
KANTOR : YOGYAKARTA



Order ID
Date
Time


IBOC - Maintenance (SIO)

: 83215
: 19/04/2024
: 11:54:49


Sender's Reference:
:20:SIOYGY00049224
Bank Operation Code:
:23R:CRRD
Value Date/Currency/Interbank Settled Amount:
:32A:246419USD900,
Ordering Customer:
:50K:/0000000723567241
SDR ARI KUSUMA WARDANA
UNIVERSITAS PGRI
YOGYAKARTA
INDONESIA
Ordering Institution:
:52A:BNINIDJAXXX
Account With Institution:
:57D:STATE BANK OF INDIA
PATEL NAGAR INDIA
Beneficiary Customer:
:59:/37014428711
GNURAV PUBLICATIONS
HISAR 125001
HARYANA
INDIA
Remittance Information:
:70:ARTICLE PAYMENT OPTIMIZING
BIOHERBICIDE APPLICATION
PATMAN ROC 1057
Details Of Charges:
:71A:OUR
Sender to Receiver Information:
:72:SWIFTCODE SEININER100




Tanggapan pembayaran dari Jurnal: 20 April 2024




Galley Proof: 9 Mei 2024





Online Publication: 22-5-2024



Optimizing bioherbicide application timing and plant spacing to manage weed growth in rice (*Oryza sativa* L.) cultivation

DOI: 10.31830/2348-7542.2024.ROC-1057 | Article Id: ROC-1057

Authors and affiliations

PAIMAN, ZULKAIISI NURUL AMAR, IKHSAN FIRMANSYAH,
HAGENG RAHMATULLOH PRIYO PAMBUDI, ANGGUNILA,
CICILIA TRI KUSUMASTUTI AND ARDIYANTA
paiman@upy.ac.id

Address : Department of Agrotechnology, Faculty of
Agriculture, Universitas PGRI Yogyakarta, Yogyakarta 55182,
Indonesia

Submitted Date : 3-02-2024

Accepted Date : 20-04-2024

Online Published: 22-05-2024

Abstract

Weed management is crucial for successful rice cultivation. Bioherbicides derived from cogon grass extract offer a promising solution for weed control. Additionally, optimizing plant spacing has been recognized as an effective method to suppress weed growth. Thus, this study aims to investigate the impact of cogon grass extract application timing and plant spacing on weed growth in rice cultivation. The study was conducted from October 2022 to February 2023 in the Special Region of Yogyakarta, Indonesia. The study was 3 x 2 factorial and arranged in a randomized completely block design (RCBD). The first factor was the bioherbicide application timing: control (no application), three days before planting, and at the time of planting. The second factor was plant spacing: 15 x 15 and 25 x 25 cm. In total, there were six treatment combinations and three replications (replications as blocks). The research results indicated the application timing of bioherbicide did not significant effect on the growth of weed and rice, as well as the grain dry weight. *Fimbristylis miliacea* was obtained as the dominant weed species. Bioherbicide application at the time of planting could alter the composition of weed species compared to the control in plant spacing of 15 x 15 cm. Furthermore, the plant spacing of 25 x 25 cm resulted in higher grain and weed dry weight (19.40 g/clump



- Editorial Board
- Author guidelines
- Aims and Scope
- Role of Reviewers
- Editorial Policy
- Role of Authors
- Peer-Review Article Process

Publikasi Artikel: 7 Juni 2024

<https://mail.google.com/mail/u/0/#inbox/FMfcgzQVwwwMFsKFqRXkZzGZXwLQbwQf>

The screenshot shows a Gmail inbox with the following email list:

- Gaurav Publications** - Complete Publication of manuscript no. ROC-1057 entitled Optimizing bioherbicide application timing and pl... 7:52 PM. Attachments: roc.png, 1,207-215.pdf, Contents ROCV...
- Lembaga Layanan Pen.** - Permohonan Pengisian Survei Kepuasan Masyarakat LLDIKTI Wilayah V Triwulan II Tahun 2024 - Yth. Pimpina... Jun 6. Attachment: Permohonan Pe...
- eContent Pro** - You're Invited: How to Get Published in High Impact Journals Webinar - View Online | Forward to a Colleague ... Jun 6
- LinkedIn** - 1 person noticed you - See who is viewing your profile Jun 5
- Levi's® Red Tab™** - DISC 50% OFF for Banded Collar Shirt 🔥 - Suka gaya yang casual dengan sentuhan modern? Levi's® Men's ... Jun 5
- ResearchGate** - Paiman, we think we found some of your full-texts online - Paiman, we think we found some of your full-text... Jun 5
- Applied Ecology** - Extract of referee opinions for MS 15413 - Dear Author, We are sending you enclosed an extract of referee o... Jun 4
- APL Energy** - Open Access Research in Perovskites - Read the latest articles recently published on perovskites View in Bro... Jun 4
- Megan Dong - MDPI** - Call for Paper Reminder: [IJMS] (IF: 5.6, ISSN: 1422-0067) — Special Issue "Regulatory Mechanism and Netw... Jun 4
- Open Agriculture** - OPAG-D-24-00119 - Review Invitation Reminder - Ref.: Ms. No. OPAG-D-24-00119 Evaluating Agricultural Yiel... Jun 4
- Lembaga Layanan Pen.** - Informasi Usulan Kenaikan Pangkat Dosen Non-PNS (TMT Juli 2024) - Yth. Pimpinan Perguruan Tinggi Swast... Jun 4. Attachment: Informasi Usula...
- LinkedIn** - Dr. Ir. Paiman., follow Nasyella A. - Corporate Organization & Human Capital Development at ASTRA Infra - S... Jun 4
- eContent Pro** - The Weekly Scholarly Experts - View Online | Forward to a Colleague www.eContentPro.com What is Typese... Jun 4

The screenshot shows the full content of the email from Gaurav Publications:

Complete Publication of manuscript no. ROC-1057 entitled Optimizing bioherbicide application timing and plant spacing to manage weed growth in rice (*Oryza sativa* L.) cultivation (External) (Inbox x)

Gaurav Publications to me, paimanupy ▾ Thu, Jun 6, 7:52 PM (9 hours ago) ☆ ↶ ⋮

Dear Dr. PAIMAN,

We are glad to inform you that your Manuscript No. **ROC-1057** entitled **Optimizing bioherbicide application timing and plant spacing to manage weed growth in rice (*Oryza sativa* L.) cultivation** has been published in **VOLUME 25 ISSUE 2 (JUNE)**, page range 207-215 in **Research on Crops**. Enclosed is the Final article. You can find your manuscript online on <https://gauravpublications.com/journal/research-on-crops/ROC-1057>.

Congratulations to you once again on your manuscript publication in Gaurav Publications and we look forward to receiving more of your good submissions.

Best Regards
Managing Editor
Gaurav Publications |
17, Ground Floor I Om Shopping Complex I Old Area Udayapuram Street I
Behind Main Post Office | Hisar - 125001 | Haryana (INDIA) |
Web: www.gauravpublications.com | E-mail: info@gauravpublications.com

3 Attachments • Scanned by Gmail

- Research on Crops
- 1,207-215.pdf
- Contents ROCV...

CONTENTS

RESEARCH PAPERS

AGRONOMY & SOIL SCIENCE

- PAIMAN** 207 Optimizing bioherbicide application timing and plant spacing to manage weed growth in rice (*Oryza sativa* L.) cultivation
ZULKAISI NURUL AMAR
IKHSAN FIRMANSYAH
HAGENG RAHMATULLOH PRIYO-
PAMBUDI
ANGGUNILA
CICILIA TRI KUSUMASTUTI
ARDIYANTA
- T. D. H. TRUONG** 216 Enhancing the efficacy of biochar derived from crop residues on rice (*Oryza sativa* L.) dry matter, yield accumulation and soil organic carbon in Central Vietnam
T. D. TRAN
Q. C. NGUYEN
T. T. D. NGUYEN
T. T. H. HOANG
- T. X. P. TRAN** 222 Impact of rice straw biochar and reduced chemical fertilizer on the growth and yield of rice (*Oryza sativa* L.) in Central Vietnam
D. H. TRAN
- SUNITA MEHER** 228 Efficacy of broad-spectrum herbicide mixtures on weed flora in wet direct seeded rice (*Oryza sativa* L.) in the east coast plain region of India
SANJOY SAHA
NITISH TIWARI
ASHIRBACHAN MAHAPATRA
JAGADISH JENA
MINU MOHAN
- M. JAYARAJ** 235 Influence of establishment methods and seed coating techniques on the performance of direct seeded rice (*Oryza sativa* L.) under Periyar Vaigai command area of Tamil Nadu
R. DURAI SINGH
T. RAGAVAN
SYED ABUL HASSAN HUSSAINY
P. SUBRAMANIAN
B. BHAKIYATHU SALIHA
B. SIVASANKARI
- VICTORIA VLADIMIROVNA-** 241 Environmental surveillance of the rice massif in the Kuban-Protoka river region
USTINOVA
ALEXEY DMITRIEVICH TIMOSHKIN
- T. D. PHUNG** 247 Climate change impacts on rice cultivation paddies in the Plain of Reeds, Vietnam
- HAYDER ABID AL-RAZZAQ BAQIR** 253 Response of yield and its components of two wheat varieties to foliar spraying with stearic acid
NAJAT HUSSEIN ZEBOON
- NASEER FARAJ SHACHAI** 258 Investigating the influence of growth inhibitors on the growth and yield of oat (*Avena sativa*)
AHMED ALI HUSSEIN
BASHEER ABDALA ABRAHEEM

FATIMA E. ALDAINY 263 Assessment of maize leaf indicators under water
BASHEER A. ABRAHEEM reduction levels: Impact of polymer, charcoal, and
antiperspirant treatments

MEHRNOUSH AMINISARTESHNIZI 269 Analysing *Pandora* associated with *Brassica oleracea*
using 16S rDNA from Limpopo Province, South Africa

T. D. PHUNG 273 Assessing the impact of unsustainable exploitation and
climate change on soil properties alterations in the
mountainous regions, Vietnam

T. N. LE 280 Land cover change assessment in Thai Nguyen
D. D. NGUYEN Province, Vietnam using GIS and remote sensing
D. T. NGUYEN techniques

HORTICULTURAL & VEGETABLE CROPS

CUCU SUHERMAN 286 Optimizing oil palm farming: Soil quality, fertilization
JAJANG SUPRIATNA and agro-environmental performance
ANNE NURAINI
SYARIFUL MUBAROK

GAVIN GELDENHUYS 292 Evaluating the efficiency of Chelex 100 for DNA
extraction in orange (*Citrus sinensis*)

K. P. MALATJI 296 Response of selected nutrient elements in leaf tissues
P. W. MASHELA of sweet potato cv. 'Bophelo' to Nemafric-BL
K. M. POFU phytonematicide

RYAN WIDIANTO 302 Chemical desiccation of potato haulms to increase skin
DENNY KURNIADIE hardness and potato yield

GAVIN GELDENHUYS 308 Investigation on the population dynamics of
Helicotylenchus pseudorobustus and its relationship
with soil factors in tomato fields

GAVIN GELDENHUYS 312 Identification of molecular characters of tomato
seedlings (*Solanum lycopersicum*) from Limpopo
Province, South Africa, using 28S rDNA

BUSHRA SOLANKI 316 Unveiling the efficacy of a bacterial antagonist in the
MOHD SAGHIR KHAN management of tomato wilt disease caused by *Ralstonia*
solanacearum

GAVIN GELDENHUYS 324 Relationship between soil factors with *Helicotylenchus*
pseudorobustus isolated from tomato fields in Limpopo
Province, South Africa

SAGAR MAITRA 328 Growth, productivity and quality of colored capsicum
MASINA SAIRAM (*Capsicum annuum* L.) as influenced by hybrids and plant
D. T. SANTOSH growth regulators under controlled environment
DINKAR JAGANNATH GAIKWAD conditions
UPASANA SAHOO

MEDICINAL PLANTS

MAROPENG CHOSHI 336 Influence of growth stages on the nutritional value of
YVONNE MAILA *Momordica balsamina* leaf powder

MOSES TIMANA 343 *Sphenoptera* sp. (Buprestidae: Coleoptera) threatening
ZAKHELENI DUBE wild cancer bush (*Sutherlandia frutescens*) population
NCOBILE NKUNA
MMAGADIMA SEBATI
THOBILE MKHWANAZI
RIVONINGO UBISI

PLANT PROTECTION

MEHRNOUSH AMINISARTESHNIZI 350 Phylogenetic position of *Aceria* sp. (Trombidiformes:
Eriophyidae) based on 28S rDNA partial sequences
Limpopo Province, South Africa

MEHRNOUSH AMINISARTESHNIZI 353 Phylogenetic position of *Discolaimium* (Nematoda:
Dorylaimida) from South Africa using 28S rDNA

AGUS SUSANTO 356 Toxicity of several insecticides against fruit flies
WAHYU DARADJAT NATAWIGENA *Bactrocera dorsalis* (Hendel) and *Bactrocera carambolae*
YAYAN SUMEKAR (Drew & Hancock) (Diptera: Tephritidae) in the
ARIEF NASRULLAH HIDAYATH laboratory

