

BUKTI KORESPONDENSI PROSIDING

Judul :

The Multi-agents System for Job Recommendation

- 1. Bukti Pembayaran (30 Agustus 2019)**
- 2. Ucapan terimakasih atas partisipasi dari panitia (10 September 2019)
- Sertifikat**
- 3. Program Book**
- 4. Publish artikel (23 May 2020)**

- 1. Bukti Pembayaran (30 Agustus 2019)**



Meilany Nonsi Tentua <meilany@upy.ac.id>

ICDM 2019 : Payment Proof for Abstract ABS-52

1 message

ICDM 2019 <icdm2019.automail@centerconf.org>

Fri, Aug 30, 2019 at 11:21 AM

Reply-To: icdm@upgris.ac.id

To: meilany@upy.ac.id

Cc: icdm@upgris.ac.id

Dear Ms. Meilany Nonsi Tentua,

Payment proof for the following abstract has been confirmed.
Payment receipt can be printed directly from your account.

Abstract ID: ABS-52

Title: The Multi-agents System for Job Recommendation

Author: Meilany Nonsi Tentua(a,b)*, Azhari Azhari (b), Aina Musdholifah(b)

Presenter: Meilany Nonsi Tentua

Paid amount: IDR 3,000,000

Payment receipt can be printed directly from your account as follows:

- Login to your account
- Click "Abstract" on the top menu
- Click "Print Payment Receipt" button below your abstract

Thank you.

Best regards,

ICDM 2019 Organizing Committee

Homepage : <http://icdm.upgris.ac.id>Email : icdm@upgris.ac.id

<http://konfrenzi.com> | Web Systems for Scientific Conferences

**2. Ucapan terimakasih atas partisipasi dari panitia
(10 September 2019)
- Sertifikat**



Meilany Nonsi Tentua <meilany@upy.ac.id>

A thank you note from the committee

1 message

noreply.icdm upgris <noreply.icdm@upgris.ac.id>
To: meilany@upy.ac.id

Tue, Sep 10, 2019 at 10:12 AM

Dear Prof. Meilany Nonsi Tentua

We would like to express our most sincere gratitude for your enthusiastic participation on ICDM 2019. It was your solid and supportive cooperation that made the whole programs of the event ran well. We would also like to apologize for any inconvenience caused.

In this email several things are informed:

(1) E-certificate

We are sending the e-certificate of attendance as attached.

(2) Gallery

For your information, we posted pictures that our team had captured during the conference and now available on the website to download, <http://icdm.upgris.ac.id/gallery/>


(3) Full-paper submission reminder


The deadline for the full-paper submission is on 15 September 2019. Please upload your full paper through the submission system on the website and not by email. The paper format must follow IOP template.

(4) Evaluation form

At last but not least, we would really appreciate if you could give us your feedback through the following link https://docs.google.com/forms/d/e/1FAIpQLSf9Y3I7jlL8uvTG2A6Wx2RvxMoRHVpowO6zUa-c4UvzEUF4Gg/viewform?usp=sf_link , by Thursday 13 September 2019. Your participation in the conference evaluation is very important for us to improve our service at the next conference organization.

Best regards,
ICDM 2019 committee
icdm.upgris.ac.id
icdm@upgris.ac.id

 **ABS-52.pdf**
7173K



ICDM



UNIVERSITAS PGRI SEMARANG
UPGRIS
The Manning University

In the 7th International Conference on DV-X α Method
The Advances-Related Experiments and Theories on Material Science

Certificate

is awarded

Meilany Nonsi Tentua, Azhari Azhari, Aina Musdholifah

as Oral Presenter with the paper entitled

The Multi-agents System for Job Recommendation

In Universitas PGRI Semarang
2-4 September 2019, Semarang, Indonesia

Chairperson of the society,

Prof. Yoshiyuki KOWADA

Hyogo Univ.of teacher Educ., Japan

Rector,

Dr. Muhdi, S.H., M.Hum.

Universitas PGRI Semarang, Indonesia



Co-host



3. Program Book



PROGRAM BOOK

In the 7th International Conference on DV-X α Method

The Advances-Related Experiments and Theories on Material Science
Universitas PGRI Semarang, Central Java, Indonesia
2-4 September 2019

Co-host



Welcome Remarks: Conference Chairperson ICDM 2019



Mega Novita, S.Si., M.Si., M.Nat.Sc., Ph.D

It is a great honour and pleasure to welcome you at the 7th International Conference on DV-X α Method: The Advances-Related Experiments and Theories on Material Science (ICDM 2019). We are happy to have you from 2 - 4 September 2019 in Semarang. This event is organized by Universitas PGRI Semarang (UPGRIS - Indonesia) in cooperation with DV-X α Society (Japan), Kwansai Gakuin University (Japan), Kobe University (Japan), Universitas 17 Agustus 1945 Semarang (Indonesia), Universitas PGRI Madiun (Indonesia) and under the coordination of Centre of Science and Technology UPGRIS (Indonesia).

ICDM is an international event aimed at presenting contemporary research and accommodating discussion for scientists, scholars, engineers, students, research institutes, and industries in the field of advanced materials, science and information technology.

The interaction between the theoretical and practical aspects of integrating the innovative and advanced technologies would construct new fundamentals in the engineering field. Therefore, it would be one direction that ICDM should pursue, hence the theme of our symposium is "The Advances-related experiments and theories on material science".

We would like to express our gratitude to all keynote and invited speakers, and all participants who have travelled far away. Our appreciation also goes to the committee members who have worked hard to make this event possible. Thus, we wish all participants may engage in the fruitful and meaningful discussion in this conference.

Welcome Remarks:



Rector, Universitas PGRI Semarang**Dr. Muhdi, S.H., M.Hum**

I would like to warmly welcome you all to Semarang, Indonesia.

It is an honour for Universitas PGRI Semarang (UPGRIS-Indonesia) to be the host of the 7th International Conference on DV-X α Method: The Advances-Related Experiments and Theories on Material Science (ICDM 2019). We are pleased to have cooperation with DV-X α Society to hold this international event. As we know, the previous conferences related to this topic had been conducted six times: once in Hungary, three times in South Korea, two times in Japan and supported by some institutions as the hosts: The Institute of Nuclear Research of the Hungarian Academy of Sciences (ATOMKI-Hungary), Seoul National University (South Korea), Kunsan National University (South Korea), National Institute of Physical and Chemical Research (RIKEN-Japan), Korean Basic Science Institute (KBSI-South Korea), and University of Hyogo (Japan).

Therefore, we expect that the cooperation in running the event will be successful in the implementation. We also proudly invited Kwansai Gakuin University (KGU-Japan), Kobe University (Japan), Universitas PGRI Madiun (UNIPMA-Indonesia) and Universitas 17 Agustus 1945 (UNTAG-Indonesia) to be the co-hosts of the conference.

We would like to express our sincere gratitude and appreciation to all parties for any fatigue and support in organizing this event. Furthermore, we expect that this conference can facilitate researchers, engineers,

lecturers, academicians and practitioners to discuss and collaborate. We hope that this conference have positive impacts to strengthen and foster international networks and publications related to this field and

particularly introduce DV-X α method in Indonesia. Herewith, we are pleased to share the latest development of the powerful method to analyse advanced materials, science and IT through DV-X α method.

Eventually, I sincerely hope that this conference may bring benefits for all of us.

Thank you

TABLE OF CONTENTS

WELCOME REMARKS	0
TABLE OF CONTENTS	4
GENERAL CONFERENCE PROGRAM	5
LECTURE SESSIONS TIMETABLE	17
TABLE OF ABSTRACTS	18
GALA DINNER	107
GUIDELINES FOR PRESENTER(S)	108
EXCURSION	110

GENERAL CONFERENCE PROGRAM

Conference Day 1: September 2, 2019	
08:00-09:00	Registration
09:00-09:15	Welcoming Speech Mega Novita, S.Si., M.Si., M.Nat.Sc., Ph.D (Chairperson of ICDM 2019, Universitas PGRI Semarang, Indonesia)
09:15-11:00	Lecture Session 1 Moderator: Prof. Jun Onoe (Nagoya University, Japan)
09:15-10:00	ABS-5 (Keynote Speaker) <i>Supramolecular Rare Sugar</i> Prof. Tomohiko Ishii Kagawa University, Japan
10:00-10:15	Coffee Break
10:15-11:00	ABS-42 (Keynote Speaker) <i>Microscopic crystal field effects in impurity centers formed by the transition metal ions</i> Prof. Mikhail G. Brik University of Tartu, Estonia
11:00-12:00	Lecture Session 2 Moderator: Dr. Retno Ambarwati (Universitas 17 Agustus 1945 Semarang Indonesia)
11:00-11:20	ABS-3 (Invited Speaker) <i>Sorption characteristics of ferrocyanide nanoparticles to metal ions in nitric acid solution for development of vitrification processes of high level radioactive nuclear wastes</i> J. Onoe ¹ , S. Watanabe ¹ , M. Harigai ² , Y. Inaba ² , and K. Takeshita ² ¹ Nagoya University, ² Tokyo Inst. Technol., Japan

11:20-11:40	ABS-29 (Invited Speaker) <i>Weak Hydrogen Bonding of Polymers Studied by Terahertz and Low-frequency Raman Spectroscopy</i> Prof. Harumi Sato Kobe University, Japan
11:40-12:00	ABS-35 (Invited Speaker) <i>Recent applications of discrete variational multi-electron method</i> Prof. Kazuyoshi Ogasawara Kwansei Gakuin University, Japan
12:00-13:00	Lunch Break
13:00-14:45	Poster Session (3 Minutes Presentation) Moderator: Dr. Dyah Nugrahani (Universitas PGRI Semarang, Indonesia)
	ABS-4 <i>Crystal Size Effect in Polaritonic Luminescence from Atomic Cryocrystals</i> Alexander Ogurtsov, Olga Bliznjuk, Nataliia Masalitina National Technical University "KhPI" Ukraine
	ABS-9 <i>Prediction of $4f^2 \rightarrow 4f15d1$ transition energy of Pr^{3+} in fluorides based on first-principles calculations and machine learning</i> Hayato Obata, Kazuyoshi Ogasawara Kwansei Gakuin University, Japan
	ABS-10 <i>Optimization of first-principles calculation conditions of multiplet energies in Fe^{3+} and Co^{3+} in $\alpha-Al_2O_3$</i> Judo Fusamoto, Kazuyoshi Ogasawara Kwansei Gakuin University, Japan
	ABS-12 <i>Prediction of $4f^7 \rightarrow 4f65d1$ transition energies of Eu^{2+} in oxides based on first-principles calculations and machine learning</i> Hiroyuki Hori, Shota Takemura, Kazuyoshi Ogasawara Kwansei Gakuin University, Japan

	<p>ABS-18 Controlling of the Spin State by Jahn-Teller Distortion in Octahedral Hexa- Coordinate Metal Complex T. Oguma, Y. Huzikawa, D. Tsuneda, Y. Tsutsumi, M. Nakano, T. Ishii, G. Sakane, Kagawa University, Japan</p>
	<p>ABS-19 Single-Crystal X-Ray Structural Analysis and Electronic Structures Calculation of Rare Sugar Bui Quoc Huy, Tomohiko Ishii, Norito Taniguchi, Natsumi Nagayama, Genta Sakane Kagawa University, Japan</p>
	<p>ABS-22 <i>Prediction of emission energy of Cr³⁺ in oxides based on first-principles calculations and machine learning</i> Tsubasa Hori, Kazuyoshi Ogasawara Kwansei Gakuin University, Japan</p>
	<p>ABS-23 <i>Consideration of Super Atomic Hybrid Orbital in Super Atom M@Al₁₂ (M = Be, Mg, Ca, Sr)</i> Teruyuki Toba, Ishii Tomohiko, Syoya Yamasaki, Sakane Genta Kagawa University, Japan</p>
	<p>ABS-24 <i>Mechanism of Optical Rotation of Amino Acids Using Electronic State Calculation</i> Shuuichi Katumoto, Manami Mimura, Tomohiko Ishii, and Genta Sakane Kagawa University, Japan</p>
	<p>ABS-25 <i>Generation of Two-Dimensional Spectrochemical Series in Tetrahedral Complexes</i> Takuhiro Yamamoto, Takeyoshi Oguma, Momoe Nakano, Akira Tsuneda, Yuki Tsutsumi, Tomohiko Ishii Kagawa University, Japan</p>
	<p>ABS-26 <i>Effect of Molecular Distortion of Ligand Field Splitting in Five-Coordinated Metal Complex</i></p>

	Yohiki Fujikawa, Takeyoshi Oguma, Takuhiro Yamamoto, Yuki Tsutsumi, Dan Tsuneda, Momoe Nakano, Tomohiko Ishii, Genta Sakane Kagawa University, Japan
	ABS-27 <i>Single crystal X-ray structural analysis of sugar molecules and electronic theory</i> Norito TANIGUCHI, Tomohiko ISHII, Bui Quoc Huy, Natsumi NAGAYAMA, Genta SAKANE Kagawa University, Japan
	ABS-31 <i>Contribution of edge-carbon atoms and non-benzenoid rings in graphitic carbons to π^* peak profiles in CK-XANES</i> Yasuji Muramatsu and Yuma Hirai University of Hyogo, Japan
	ABS-48 <i>Designing Enterprise Architecture in Koperasi Karyawan using TOGAF Architecture Development Method</i> Noora Qotrun Nada, Setyoningsih Wibowo, Mega Novita Universitas PGRI Semarang, Indonesia
	ABS-49 <i>DECISION SUPPORT SYSTEM MUSEUM AMBASSADOR USING TOPSIS METHOD</i> Setyoningsih Wibowo, Maichifa Ayuning Tyas, Noora Qotrun Nada, Mega Novita Universitas PGRI Semarang, Indonesia
	ABS-50 <i>Design of Augmented Reality Game app "IntroME" using ADDIE Model</i> Rahmat Robi Waliyansyah, Irfan Nur Ridwan, Khoirul Huda, Febrian Murti Dewanto Universitas PGRI Semarang, Indonesia
	ABS-53 <i>DESIGNING ANDROID BASED EDUCATION GAME AKSARA JAWA USING SHUFFLE RANDOM ALGORITHM</i> Nugroho Dwi Saputro, Tri Ana Romadhani, Febrian Murti Dewanto Universitas PGRI Semarang, Indonesia
14:45-15:00	Coffee break

15:00-16:30	Lecture Session 3 Moderator: Prof. Yasuji Muramatsu (University of Hyogo, Japan)
15:00-15:45	ABS-6 (Keynote Speaker) <i>X-ray Absorption Spectroscopy and Its Simulation for Some Metal Oxides</i> Prof. Deok-Yong Cho Chonbuk National University, South Korea
15:45-16:30	ABS-46 (Keynote Speaker) <i>Carotenoid Dynamics Free and Bound to Pigment Protein Complexes: The Role of the 11Bu- State</i> Prof. Ferdy Samuel Rondonuwu Satya Wacana Christian University, Indonesia
16:30-17:30	Lecture Session 4 Moderator: Asst. Prof. Laily Nur Affini
16:30-16:50	ABS-30 (Invited Speaker) <i>Metal complexes with sulfur-containing ligands</i> Masahiro Mikuriya and Makoto Handa Kwansei Gakuin University, Japan
16:50-17:10	ABS-54 (Invited Speaker) <i>Electronic state of sulfide-based alkali-ion conducting solid-state electrolytes applied to all-solid-state secondary batteries</i> Yoshiyuki Kowada, Yasukazu Akaki, Akiotoshi Hayashi, and Masahiro Tatsumisago Hyogo University of Teacher Education, Japan
17:10-17:30	ABS-55 (Invited Speaker) <i>Geometry and Stability of Small Gold Cluster Ions by Graph Theory and Hückel model</i> Rika Sekine, Takashi Kasai, Eri Furuta, and Masaya Okakura Shizuoka University, Japan
17:30-19:00	Break
19:00-21:00	Gala dinner

Conference Day 2: September 3, 2019	
08:00-10:00	Parallel room 1 Moderator: Prof. Masataka Mizuno (Osaka University, Japan) Asst. Moderator : Asst. Prof. Laily Nur Afini (Universitas PGRI Semarang, Indonesia)
08:00-08:20	ABS-1 (Invited Speaker) <i>First-Principles Calculation of Laser Crystal Multiplet Levels via Hybridized Density Functional Theory and Configuration Interaction within the OLCAO Method</i> Dr. Benjamin Walker University of Missouri - Kansas City, USA
08:20-08:40	ABS-8 (Invited Speaker) <i>Systematic First-Principles Calculations of Charge Transfer Transitions of Transition Metal Ions (Sc^{3+}, Ti^{3+}, V^{3+}, Cr^{3+}, Mn^{3+}, Fe^{3+}) in $\alpha-Al_2O_3$ with Structural Optimization</i> Shota Takemura ¹ , Mega Novita ² , Kazuyoshi Ogasawara ¹ Kwansei Gakuin University, Japan
08:40-09:00	ABS-16 (Invited Speaker) <i>Study of LCAO-MO calculation by using completely numerical basis functions</i> Katsumi Nakagawa MO Basics Research
09:00-09:15	ABS-13 <i>Lattice Relaxation Effects on the Multiplet Energies of Ruby Under Pressure using One-Electron Calculations</i> Mega Novita, Nur Cholifah, Kazuyoshi Ogasawara Universitas PGRI Semarang, Indonesia, Kwansei Gakuin University, Japan
09:15-09:30	ABS-14 <i>Optical properties of Co^{3+} doped in $\alpha-Al_2O_3$ with Considering Lattice Relaxation Effect</i> Mega Novita, Duwi Nuvitalia, Nur Cholifah, Kazuyoshi Ogasawara Universitas PGRI Semarang, Indonesia, Kwansei Gakuin University, Japan
09:30-09:45	ABS-15 <i>Effects of Transition Metal Doping to Cerium-Oxides in Thermal Reduction Reaction</i>

	Takaki Nishimura, Yuki Tsutsumi, Tomohiko Ishii, Tatsuya Kodama, and Genta Sakane Kagawa University, Japan
09:45-10:00	ABS-61 <i>Extraction Sentiment Analysis Using Naive Bayes Algorithm and Reducing Noise Word Applied in Indonesian Language</i> Aris Tri Jaka Harjanta, Bambang Agus Herlambang Universitas PGRI Semarang, Indonesia
08:00-10:00	Parallel room 2 Moderator: Prof. Prompong Pienpinijtham (Chulalongkom University Thailand) Asst. Moderator: Dr. Muchamad Taufiq Anwar (Universitas Stikubank, Indonesia)
08:00-08:20	ABS-2 (Invited Speaker) <i>Soil decontamination by cesium-free mineralization with CaCl₂: effect of chlorine on phase transformation of a clay mineral</i> Iwao Shimoyama Japan Atomic Energy Agency, Japan
08:20-08:40	ABS-7 (Invited Speaker) <i>Study on The Higher-Order Structure and Hydrogen Bonding of Biodegradable Polymer by Low-Frequency Vibrational Spectroscopy</i> Dian Marlina, Harumi Sato Kobe University, Japan
08:40-09:00	ABS-21 (Invited Speaker) <i>An elaborate response in molecular random network - Approach for information processing functionality-</i> Megumi Akai-Kasaya Osaka University, Japan
09:00-09:15	ABS-11 <i>Cultivation of Oreochromis niloticus with an Environmentally friendly Bioflog System</i> Endah Rita Sulistya Dewi, Ary Susatyo Nugroho, Maria Ulfah Universitas PGRI Semarang, Indonesia

09:15-09:30	ABS-20 <i>Data Mining Application with Fuzzy Logic Method for Monitoring Vehicle Position Based on Android</i> Munaf Ismail, Sri Arttini Dwi Prasetyowati, Ida Widihastuti Universitas Islam Sultan Agung, Indonesia
09:30-09:45	ABS-33 <i>Smart Aquaponics Design Using Internet of Things Technology</i> R Hafid Hardyanto, Prahenusa Wahyu Ciptadi Universitas PGRI Yogyakarta, Indonesia
09:45-10:00	ABS-36 <i>Alternative Solutions For Determining Village Funds Using Weight Product Method</i> Tri Hastono, Firdiyan Syah Universitas PGRI Yogyakarta, Indonesia
10:00-10.15	Coffee Break
10:00-10:30	Preparation
10:30-12:30	Heading to Candi Borobudur
12:30-13:30	Lunch
13:30-18:00	Excursion
18:00-19:00	Dinner
19:00-21:00	Heading to Grand Candi Hotel, Semarang

Conference Day 3: September 4, 2019	
08:00-10:45	Parallel room 1 Moderator: Prof. Katsumi Nakagawa (MO Basic Research, Japan) Asst. Moderator: Dr. Dyah Nugrahani (Universitas PGRI Semarang, Indonesia)
08:00-08:20	ABS-34 (Invited Speaker) <i>Stability of carbon-vacancy complexes in α-Fe</i> Masataka Mizuno, Waka Yamada, Kazuki Sugita, Hideki Araki Osaka University, Japan
08:20-08:40	ABS-37 (Invited Speaker) <i>DESIGN OF ELECTRICITY ENERGY SOURCES IN SEMARANG PGRI UNIVERSITY USING OFF-GRID SOLAR PANEL SYSTEMS</i> Adhi Kusmantoro, Mega Novita, Th.Indriati Wardani Universitas PGRI Semarang, Indonesia
08:40-09:00	ABS-38 (Invited Speaker) <i>Novel shape analysis method for single bioparticles in aqueous solutions</i> Sou Ryuzaki, Makusu Tsutsui, Masateru Taniguchi Kyushu University, Japan
09:00-09:15	Coffee break
09:15-09:30	ABS-41 <i>BIOCONVERSION ON WASTEWATER OF SOYBEAN USING MICROBIAL FUEL CELL</i> Yohanes A Cahyono, Tilana Madurani, Widya F Azzahra and Retno A. S. Lestari Universitas 17 Agustus 1945 Semarang, Indonesia
09:30-09:45	ABS-44 <i>Potential of Ketapang Seed Oil (<i>Terminalia catappa</i> Linn) as Basic Material Mono-diglyceride Biodegradable Surfactant</i> Ery Fatarina Purwaningtyas, Khoir Fadilah Universitas 17 Agustus 1945 Semarang, Indonesia
09:45-10:00	ABS-45 <i>BIOPLASTIC FROM JACKFRUIT SEEDS AND RICE WASTE</i> Retno A.S. Lestari, Mega Kasmiyatun, Kevin Dermawan, Alfia N. Aini, Nur Riyati, Finka R. Putri

	Universitas 17 Agustus 1945 Semarang, Indonesia
10:00-10:15	ABS-52 <i>The Multi-agents System for Job Recommender System</i> Meilany Nonsi Tentua, Azhari SN, Aina Musdholifah Universitas PGRI Yogyakarta, Indonesia
10:15-10:30	ABS-64 <i>Color Based Segmentation Using Fuzzy C Means for Bamboo as Environmentally Friendly Material</i> Khoiriya Latifah, Joko Siswanto, Bambang Supriyadi, Abdul Rochim Universitas PGRI Semarang
10:30-10:45	ABS-58 <i>Interaction Analysis Application of Arduino Industrial Automation Trainer Based on Project to Improve Cognitive Ability and Bodily Kinesthetic Ability</i> Ir. Sulistyanning Kartikawati, MM, M.Pd., Sekreningsih Nita, S.Kom, MT. Universitas PGRI Madiun, Indonesia
08:00-10:45	Parallel room 2 Moderator: Dr. Benjamin Walker (University of Missouri - Kansas City) Asst. Moderator: Dr. Dian Marlina (Kobe University, Japan)
08:00-08:20	ABS-28 (Invited Speaker) <i>Structural changes in graphene oxide induced by silver nanoparticles – TERS study</i> Prompong Pienpinijtham, Sanpon Vantasin, Oraporn Wong-u-ra, Yasutaka Kitahama, Sanong Ekgasit and Yukihiro Ozaki Chulalongkom University Thailand
08:20-08:40	ABS-51 (Invited Speaker) <i>Phosphorus recovery through crystallization of struvite (MgNH₄PO₄.6H₂O): From basic research to applications</i> Stefanus Muryanto Universitas 17 Agustus 1945 Semarang, Indonesia
08:40-09:00	ABS-57 (Invited Speaker) <i>THE EFFECT OF CHLORIDE ACID (HCL) CONCENTRATION AND FERMENTATION ON</i>

	<p><i>BIOETHANOL LEVELS FROM BREADFRUIT (<i>Artocarpus Artilis</i>) SUBSTRATE</i></p> <p>Nasrul Rofiah Hidayati Universitas PGRI Madiun</p>
09:00-09:15	Coffee break
09:15-09:30	<p>ABS-39</p> <p><i>Design of Product Monitoring System using Internet of Things Technology for Smart Manufacturing</i></p> <p>Marti Widya Sari, Herianto, IGB Budi Dharma, Alva Edy Tontowi Universitas PGRI Yogyakarta, Indonesia</p>
09:30-09:45	<p>ABS-40</p> <p><i>A Novel and Fast Memory Perturbation Method to Increase Exploration in Particle Swarm Optimization Algorithm</i></p> <p>Eri Zuliarso, Muchamad Taufiq Anwar, Kristophorus Hadiono, Iswatun Chasanah Universitas Stikubank, Indonesia</p>
09:45-10:00	<p>ABS-43</p> <p><i>Internet of Things (IOT) with Cloud Computing and Machine to Machine (M2M) communication in Aquaponics Farming Systems</i></p> <p>Prahenusa Wahyu Cipadi, R.Hafid Hardyanto</p>
10:00-10:15	<p>ABS-47</p> <p><i>Detecting Hoaxes in Indonesian News Using TF-IDF and K Nearest Neighbor</i></p> <p>Muchamad Taufiq Anwar, Edy Winarno, Iswatun Chasanah Universitas Stikubank, Indonesia</p>
10:15-10:30	<p>ABS-59</p> <p><i>ANALYSIS OF THE IMPACT NARRATIVE ALGORITHM METHOD, PSEUDOCODE AND FLOWCHART TOWARDS STUDENTS UNDERSTANDING OF THE PROGRAMMING ALGORITHM COURSES</i></p> <p>Sekreningsih Nita, Sulistyaning Kartikawati Universitas PGRI Madiun, Indonesia</p>
10:30-10:45	<p>ABS-60</p> <p><i>Prediction of The Number Job Seeker Using Backpropagation Neural Network Method</i></p> <p>Khoiriya Latifah, Agung Handayanto, Anita Rahayu Universitas PGRI Semarang, Indonesia</p>

10:45-11:00	Coffee break
11:00-12:00	Closing Ceremony Moderator: Dr. Dyah Nugrahani (Universitas PGRI Semarang, Indonesia)
11:00-11:35	DV-X α meeting Prof. Yoshiyuki Kowada (Hyogo University of Teacher Education, Japan)
11:35-11:55	Awards Prof. Katsumi Nakagawa (MO Basic Research, Japan)
11:55-12:00	Closing statement Mega Novita, S.Si., M.Si., M.Nat.Sc., Ph.D (Chairperson of ICDM 2019, Universitas PGRI Semarang, Indonesia)
12:00-13:00	Lunch



INTERNATIONAL CONFERENCE ON DV-X α METHOD

LECTURE SESSIONS TIMETABLE
The 7th International Conference on DV-X α Method (ICDM 2019)
Grand Candi Hotel, Semarang, INDONESIA
2-4 September 2019

	2-Sep-2019		3-Sep-2019		4-Sep-2019	
Time	Room1	Time	Room1	Room2	Room1	Room2
08:00-09:00	Registration	08:00-08:20	ABS-1	ABS-2	ABS-34	ABS-28
09:00-09:15	Welcoming Speech	08:20-08:40	ABS-8	ABS-7	ABS-37	ABS-51
09:15-10:00	ABS-5	08:40-09:00	ABS-16	ABS-21	ABS-38	ABS-57
10:00-10:15	Coffee break	09:00-09:15	ABS-13	ABS-11	Coffee break	
10:15-11:00	ABS-42	09:15-09:30	ABS-14	ABS-20	ABS-41	ABS-39
11:00-11:20	ABS-3	09:30-09:45	ABS-15	ABS-33	ABS-44	ABS-40
11:20-11:40	ABS-29	09:45-10:00	ABS-61	ABS-36	ABS-45	ABS-43
11:40-12:00	ABS-35	10:00-10:15	Coffee break		ABS-52	ABS-47
12:00-13:00	Lunch Break	10:15-10:30			ABS-64	ABS-59
13:00-14:45	Poster session	10:30-10:45			ABS-58	ABS-60
14:45-15:00	Coffee break	10:45-11:00			Coffee break	
15:00-15:45	ABS-6	11:00-11:35			DV-X α general meeting	
15:45-16:30	ABS-46	11:35-11:55	Awards			
16:30-16:50	ABS-30	11:55-12:00	Closing statement			
16:50-17:10	ABS-54	12:00-13:00	Lunch			
17:10-17:30	ABS-55				ABS-	Keynote Speaker
17:30-19:00	Break				ABS-	Invited Speaker
19:00-21:00	Gala Dinner				ABS-	General Speaker



TABLE OF ABSTRACTS

Title	Page
[ABS-1] First-Principles Calculation of Laser Crystal Multiplet Levels via Hybridized Density Functional Theory and Configuration Interaction within the OLCAO Method Dr. Benjamin Walker	27
[ABS-2] Soil decontamination by cesium-free mineralization with CaCl₂: effect of chlorine on phase transformation of a clay mineral Iwao Shimoyama	28
[ABS-3] Sorption characteristics of ferrocyanide nanoparticles to metal ions in nitric acid solution for development of vitrification processes of high level radioactive nuclear wastes J. Onoe, S. Watanabe, M. Harigai, Y. Inaba, and K. Takeshita	29
[ABS-4] Crystal Size Effect in Polaritonic Luminescence from Atomic Cryocrystals Alexander Ogurtsov, Olga Bliznjuk, Nataliia Masalitina	31
[ABS-5] Supramolecular Rare Sugar Tomohiko Ishii	33
[ABS-6] X-ray Absorption Spectroscopy and Its Simulation for Some Metal Oxides Deok-Yong Cho	35

<p>[ABS-7] Study on The Higher-Order Structure and Hydrogen Bonding of Biodegradable Polymer by Low-Frequency Vibrational Spectroscopy Dian Marlina (a, b)*, Harumi Sato (a)</p>	36
<p>[ABS-8] Systematic First-Principles Calculations of Charge Transfer Transitions of Transition Metal Ions (Sc³⁺, Ti³⁺, V³⁺, Cr³⁺, Mn³⁺, Fe³⁺) in α-Al₂O₃ with Structural Optimization Shota Takemura¹, Mega Novita², Kazuyoshi Ogasawara¹</p>	38
<p>[ABS-9] Prediction of 4f²→4f¹5d¹ transition energy of Pr³⁺ in fluorides based on first-principles calculations and machine learning Hayato Obata</p>	39
<p>[ABS-10] Optimization of first-principles calculation conditions of multiplet energies in Fe³⁺ and Co³⁺ in α-Al₂O₃ Judo Fusamoto, Kazuyoshi Ogasawara</p>	40
<p>[ABS-11] Cultivation of Oreochromis niloticus with an Environmentally friendly Bioflog System Endah Rita Sulistya Dewi, Ary Susatyo Nugroho, Maria Ulfah</p>	41
<p>[ABS-12] Prediction of 4f⁷→4f⁶5d¹ transition energies of Eu²⁺ in oxides based on first-principles calculations and machine learning Hiroyuki Hori, Shota Takemura, Kazuyoshi Ogasawara</p>	42
<p>[ABS-13] Lattice Relaxation Effects on the Multiplet Energies of Ruby Under Pressure using One-Electron Calculations Mega Novita (a*), Nur Cholifah (b), Kazuyoshi Ogasawara (c)</p>	43

<p>[ABS-14] Optical properties of Co³⁺ doped in α-Al₂O₃ with Considering Lattice Relaxation Effect Mega Novita (a*), Duwi Nuvitalia (b), Nur Cholifah (c), Kazuyoshi Ogasawara (d)</p>	44
<p>[ABS-15] Effects of Transition Metal Doping to Cerium-Oxides in Thermal Reduction Reaction Takaki Nishimura¹, Yuki Tsutsumi², Tomohiko Ishii², Tatsuya Kodama³, and Genta Sakane⁴</p>	45
<p>[ABS-16] Study of LCAO-MO calculation by using completely numerical basis functions Katsumi Nakagawa</p>	47
<p>[ABS-18] Controlling of the Spin State by Jahn-Teller Distortion in Octahedral Hexa- Coordinate Metal Complex T. Oguma ,a Y. Huzikawa ,a D. Tsuneda, a Y. Tsutsumi, a M. Nakano, a T. Ishii, a G. Sakane, b</p>	49
<p>[ABS-19] Single-Crystal X-Ray Structural Analysis and Electronic Structures Calculation of Rare Sugar Bui Quoc Huy, Tomohiko Ishii, Norito Taniguchi, Natsumi Nagayama, Genta Sakane</p>	50
<p>[ABS-20] Data Mining Application with Fuzzy Logic Method for Monitoring Vehicle Position Based on Android Munaf Ismail, Sri Arttini Dwi Prasetyowati, Ida Widiastuti</p>	51
<p>[ABS-21] An elaborate response in molecular random network -Approach for information processing functionality- Megumi Akai-Kasaya</p>	52

<p>[ABS-22] Prediction of emission energy of Cr³⁺ in oxides based on first-principles calculations and machine learning Tsubasa Hori, Kazuyoshi Ogasawara</p>	54
<p>[ABS-23] Consideration of Super Atomic Hibrid Orbital in Super Atom M@Al₁₂ (M = Be, Mg, Ca, Sr) Teruyuki Toba 1, Ishii Tomohiko 1, Syoya Yamasaki 1, Sakane Genta 2</p>	55
<p>[ABS-24] Mechanism of Optical Rotation of Amino Acids Using Electronic State Calculation Shuuichi Katumoto,¹ Manami Mimura,¹ Tomohiko Ishii,¹ and Genta Sakane,²</p>	57
<p>[ABS-25] Generation of Two-Dimensional Spectrochemical Series in Tetrahedral Complexes Takuhiro Yamamoto, Takeyoshi Oguma, Momoe Nakano, Akira Tsuneda, Yuki Tsutsumi, Tomohiko Ishii</p>	59
<p>[ABS-26] Effect of Molecular Distortion of Ligand Field Splitting in Five-Coordinated Metal Complex Yohiki Fujikawa, Takeyoshi Oguma, Takuhiro Yamamoto, Yuki Tsutsumi, Dan Tsuneda, Momoe Nakano, Tomohiko Ishii, Genta Sakane</p>	60
<p>[ABS-27] Single crystal X-ray structural analysis of sugar molecules and electronic theory Norito TANIGUCHI(1), Tomohiko ISHII(1), Bui Quoc Huy(1), Natsumi NAGAYAMA(1), Genta SAKANE(2)</p>	62
<p>[ABS-28] Structural changes in graphene oxide induced by silver nanoparticles – TERS study Prompong Pienpinijtham (a,b,c,*), Sanpon Vantasin (a), Oraporn Wong-u-ra (b), Yasutaka Kitahama (a), Sanong Ekgasit (b,c) and Yukihiro Ozaki (a,*)</p>	63

<p>[ABS-29] Week Hydrogen Bonding of Polymers Studied by Terahertz and Low-frequency Raman Spectroscopy Harumi Sato</p>	65
<p>[ABS-30] Metal complexes with sulfur-containing ligands Masahiro Mikuriya and Makoto Handa</p>	66
<p>[ABS-31] Contribution of edge-carbon atoms and non-benzenoid rings in graphitic carbons to π^* peak profiles in CK-XANES Yasuji Muramatsu and Yuma Hirai</p>	67
<p>[ABS-33] Smart Aquaponics Design Using Internet of Things Technology R Hafid Hardyanto (1), Prahenusa Wahyu Ciptadi (2)</p>	68
<p>[ABS-34] Stability of carbon-vacancy complexes in α-Fe Masataka Mizuno, Waka Yamada, Kazuki Sugita, Hideki Araki</p>	69
<p>[ABS-35] Recent applications of discrete variational multi-electron method Kazuyoshi Ogasawara</p>	70
<p>[ABS-36] Alternative Solutions For Determining Village Funds Using Weight Product Method Tri Hastono, Firdiyan Syah</p>	71
<p>[ABS-37] DESIGN OF ELECTRICITY ENERGY SOURCES IN SEMARANG PGRI UNIVERSITY USING OFF-GRID SOLAR PANEL SYSTEMS Adhi Kusmantoro(a*), Mega Novita(b), Th.Indriati Wardani©</p>	72

<p>[ABS-38] Novel shape analysis method for single bioparticles in aqueous solutions Sou Ryuzaki(a,b*), Makusu Tsutsui(c), Masateru Taniguchi(c)</p>	<p>74</p>
<p>[ABS-39] Design of Product Monitoring System using Internet of Things Technology for Smart Manufacturing Marti Widya Sari (a*); Herianto (b); IGB Budi Dharma (c); Alva Edy Tontowi (d)</p>	<p>76</p>
<p>[ABS-40] A Novel and Fast Memory Perturbation Method to Increase Exploration in Particle Swarm Optimization Algorithm Muchamad Taufiq Anwar, Edy Winarno, Iswatun Chasanah, Saptono Nugrohadi</p>	<p>77</p>
<p>[ABS-41] BIOCONVERSION ON WASTEWATER OF SOYBEAN USING MICROBIAL FUEL CELL Yohanes A Cahyono, Tilana Madurani, Widya F Azzahra and Retno A. S. Lestari</p>	<p>78</p>
<p>[ABS-42] Microscopic crystal field effects in impurity centers formed by the transition metal ions Mikhail G. Brik</p>	<p>80</p>
<p>[ABS-43] Internet of Things (IOT) with Cloud Computing and Machine to Machine (M2M) communication in Aquaponics Farming Systems Prahenusa Wahyu Cipadi, R.Hafid Hardyanto</p>	<p>82</p>
<p>[ABS-44] Potential of Ketapang Seed Oil (Terminalia catappa Linn) as Basic Material Mono-diglyceride Biodegradable Surfactant Ery Fatarina Purwaningtyas (1*) ; Khoir Fadilah(2)</p>	<p>83</p>

<p>[ABS-45] BIOPLASTIC FROM JACKFRUIT SEEDS AND RICE WASTE Retno A.S. Lestari, Mega Kasmiyatun, Kevin Dermawan, Alfia N. Aini, Nur Riyati, Finka R. Putri</p>	85
<p>[ABS-46] Carotenoid Dynamics Free and Bound to Pigment Protein Complexes: The Role of the 11Bu- State Ferdy S. Rondonuwu</p>	88
<p>[ABS-47] Detecting Hoaxes in Indonesian News Using TF-IDF and K Nearest Neighbor Eri Zuliarso, Muchamad Taufiq Anwar, Kristophorus Hadiono, Iswatun Chasanah</p>	89
<p>[ABS-48] Designing Enterprise Architecture in Koperasi Karyawan using TOGAF Architecture Development Method Noora Qotrun Nada*, Setyoningsih Wibowo, Mega Novita</p>	90
<p>[ABS-49] DECISION SUPPORT SYSTEM MUSEUM AMBASSADOR USING TOPSIS METHOD Setyoningsih Wibowo*, Maichifa Ayuning Tyas, Noora Qotrun Nada, Mega Novita</p>	91
<p>[ABS-50] Design of Augmented Reality Game app "IntroME" using ADDIE Model Rahmat Robi Waliyansyah (a*), Irfan Nur Ridwan (a), Khoirul Huda (a), Febrian Murti Dewanto (a)</p>	92
<p>[ABS-51] Phosphorus recovery through crystallization of struvite (MgNH₄PO₄.6H₂O): From basic research to applications S.Muryanto</p>	94

<p>[ABS-52] The Multi-agents System for Job Recommendation Meilany Nonsi Tentua(a,b)*, Azhari Azhari (b), Aina Musdholifah(b)</p>	95
<p>[ABS-53] DESIGNING ANDROID BASED EDUCATION GAME AKSARA JAWA USING SHUFFLE RANDOM ALGORITHM Nugroho Dwi Saputro(a*), Tri Ana Romadhani(b), Febrian Murti Dewanto(b)</p>	96
<p>[ABS-54] Electronic state of sulfide-based alkali-ion conducting solid-state electrolytes applied to all-solid-state secondary batteries Yoshiyuki Kowada(a*), Yasukazu Akaki(a), Akiotoshi Hayashi(b), and Masahiro Tatsumisago(b)</p>	97
<p>[ABS-55] Geometry and Stability of Small Gold Cluster Ions by Graph Theory and Hückel model Rika Sekine, Takashi Kasai, Eri Furuta, and Masaya Okakura</p>	99
<p>[ABS-57] THE EFFECT OF CHLORIDE ACID (HCL) CONCENTRATION AND FERMENTATION ON BIOETHANOL LEVELS FROM BREADFRUIT (Artocarpus Artilis) SUBSTRATE Nasrul Rofiah Hidayati</p>	100
<p>[ABS-58] Interaction Analysis Application of Arduino Industrial Automation Trainer Based on Project to Improve Cognitive Ability and Bodily Kinesthetic Ability Ir. Sulistyning Kartikawati, MM, M.Pd., Sekreningsih Nita, S.Kom, MT.</p>	101
<p>[ABS-59] ANALYSIS OF THE IMPACT NARRATIVE ALGORITHM METHOD, PSEUDOCODE AND FLOWCHART TOWARDS STUDENTS UNDERSTANDING OF THE PROGRAMMING ALGORITHM COURSES Sekreningsih Nita, Sulistyning Kartikawati</p>	103

[ABS-60] Prediction Of The Number Job Seeker Using Backpropagation Neural Network Method Khoiriya Latifah, Agung Handayanto, Anita Rahayu	104
[ABS-61] Extraction Sentiment Analysis Using naive Bayes Algorithm and Reducing Noise Word applied in Indonesian Language Aris Tri Jaka Harjanta, Bambang Agus Herlambang	105
[ABS-64] Color Based Segmentation Using Fuzzy C Means for Bamboo as Environmentally Friendly Material Khoiriya Latifah, Joko Siswanto, Bambang Supriyadi, Abdul Rochim	106

ABS-1 **First-Principles Calculation of Laser Crystal Multiplet Levels via Hybridized Density Functional Theory and Configuration Interaction within the OLCAO Method**

Dr. Benjamin Walker

University of Missouri - Kansas City

Computation of highly-localized multiplet energy levels of transition metal dopants is essential to the design of materials such as laser host crystals. A purely first-principles density functional theory-configuration interaction (DFT-CI) hybrid computational method has been developed to accurately compute multiplet energy levels for single atoms of carbon, nitrogen, oxygen, sodium, aluminum, silicon, titanium, and chromium. The multiplet energy levels have been computed with close experimental agreement in terms of magnitude and degeneracy, and the method does not depend on empirical information (i.e. Racah parameters). The computed multiplet energy level results are distributed according to term symbols, which are then compared to experimentally-observed multiplet energy levels. The hybrid method consists of analytic computation of two-electron integrals via the DFT-based orthogonalized linear combination of atomic orbitals (OLCAO) method, which are subsequently used as input for the CI-based discrete variational multi-electron (DVME) method to obtain the multiplet energy values.

Keywords: exchange-correlation; electron repulsion integral; multiplet; DVME; OLCAO; density functional theory; configuration interaction

ABS-2 **Soil decontamination by cesium-free mineralization with CaCl₂: effect of chlorine on phase transformation of a clay mineral**

Iwao Shimoyama

Japan Atomic Energy Agency

As shown in the cases of nuclear accidents in Fukushima, a key issue of environmental remediation is management of contaminated soil. Soil decontamination is generally difficult because radiocesium is strongly fixed in micaceous clay minerals in soil. Cesium-free mineralization (CFM) has been proposed as a new soil decontamination method. While conventional heat treatments utilize physical melting of soil above 1000°C, CFM utilizes chemical reaction of clay minerals at lower temperatures and can convert clay minerals to other minerals which adapt for industrial use. Precident works imply a catalytic effect of Cl which facilitates phase transformation of clay minerals, however, the detail of the reaction is not clear. In this work, I used Cs-sorbed weathered biotite (WB) as a model soil and analyzed chemical bonding states of Cl in WB in the early stage of the reaction before phase transformation using near-edge X-ray absorption fine structure (NEXAFS) spectroscopy. After heat treatments with CaCl₂, Cl K-edge NEXAFS spectra of the samples were interpreted based on electronic structures of Cl sites in some model clusters using the DV-X α method indicating that high valent Cl ions which bond with O in the clay mineral were formed in the early stage of the reaction. Furthermore, we calculated total energy of model clusters, which mimics siloxane-ditrigonal cavity of micaceous clay minerals. Discussion on structural stability indicates that Cl in the reagent can destabilize the silica network by bonding with O in clay minerals and consequently plays a role to decrease reaction temperature for phase transformation.



ABS-3 **Sorption characteristics of ferrocyanide nanoparticles to metal ions in nitric acid solution for development of vitrification processes of high level radioactive nuclear wastes**

J. Onoe, S. Watanabe, M. Harigai, Y. Inaba, and K. Takeshita

Nagoya University, Tokyo Institute of Technology

High-level radioactive nuclear liquid wastes (HLLW: radioactive metal ions in 2 M nitric acid solution) are generated in the reprocessing of spent nuclear fuels or in the decommissioning of nuclear reactors. In the process of producing the vitrified objects containing the wastes by using a glass melter, platinum-group metals (PGMs) [especially, ruthenium (Ru), rhodium (Rh), and palladium (Pd)] and molybdenum (Mo) cause serious problems: PGMs tend to be accumulated on the sidewall surface of the melter, whereas Mo compounds, so called yellow phase, are formed in the vitrified object. These issues result in (i) degradation of vitrified objects due to heterogenization and (ii) an increase in both disposal space and processing costs in conjunction with additional vitrified rods obtained by flushing the glass melter. To solve these issues, we have developed a process for efficient removal of PGMs and Mo from HLLW prior to introducing into the glass melter, using metal hexacyanoferrates (HCF) as a sorbent [1-3]. It is important to reveal the sorption mechanisms of PGMs and Mo for design of high performance HCF sorbents. In this talk, we will present the results on the sorption characteristics of Prussian blue (PB) nanoparticles, one of the HCFs, to the above metal ions in nitric acid solution, obtained using various spectroscopies and first-principles calculations.

[1] S. Watanabe et al., J. Appl. Phys. 119, 235102 (2016).



- [2] S. Watanabe et al., AIP Adv. 8, 045221 (2018).
[3] S. Watanabe et al., Chem. Phys. Lett. 723, 76–81 (2019).

Keywords: ferrocyanide nanoparticles, sorption characteristics, nuclear wastes, vitrification processes

ABS-4 Crystal Size Effect in Polaritonic Luminescence from Atomic Cryocrystals

Alexander Ogurtsov, Olga Bliznjuk, Nataliia Masalitina

National Technical University "KhPI", Kyrpychova Str. 2,
Kharkov 61002, Ukraine

The exciton-photon interaction leads to the formation of polaritonic states energetically positioned at both sides of the initial exciton. In a large ideal crystal of cubic symmetry, where the interval of the longitudinal-transverse splitting does not contain excitonic levels, the polaritonic dispersion branches lie beyond this interval at both sides of its boundaries. On the contrary, in a crystalline grain comparable or less in size than the wavelength in the substance, the interval of the longitudinal-transverse splitting is filled in continuously by excitonic states intercepting a significant part of the oscillator strength of the excitonic transition. The photoluminescence spectra of atomic cryocrystals (solid Xe and Kr) were measured at the Superlumi experimental station at HASYLAB, DESY, Hamburg. Unlike previous works, where the red polaritonic shift was small commensurably with a weak inelastic polariton-photon scattering, a large polaritonic shift of luminescence is not due to energy dissipation, the energy conservation law being met due to equal probabilities for opposite-sign energy shifts. Such effect is possible if the crystalline grains are comparable in size with light wavelength, which provides the filling in the interval of the longitudinal-transverse splitting by excitons with sufficient oscillator strength. And the sample structure must be perfect enough to lowering the exciton scattering rate with respect to the rate of the polariton formation through exciton-photon coupling. For the first time the excitation spectra of free-exciton luminescence band were recorded simultaneously below the bottom of excitonic band E and within the interval of the longitudinal-transverse splitting. The luminescence of non-equilibrium polaritons was observed

both within the longitudinal-transverse splitting interval and at photoexcitation below E . The excitation spectrum below the bottom of excitonic band is determined by competition of two processes. The first one is the creation of excitons by photons with energy E at the Lorenz tail of excitonic absorption. The second process is a competing absorption related to the direct formation of two-site excitonic polarons (self-trapped excitons). Both excitation spectra of polaritonic luminescence below E and within the longitudinal-transverse splitting interval show high sensitivity to crystal quality of the samples

Keywords: exciton-photon interaction, longitudinal-transverse splitting, inelastic polariton-photon scattering

ABS-5

Supramolecular Rare Sugar*Tomohiko Ishii*Faculty of Engineering and Design,
Kagawa University**Abstract**

We introduce rare sugars including their derivatives and supramolecular rare sugars, that have been actively researched at Kagawa University. Although rare sugars are special sugars which hardly exist in nature, we have succeeded in mass synthesis of rare sugars by utilizing biological enzyme isomerization reaction. In addition, rare sugars have various functions such as blood sugar level suppressing function, cancer cell growth suppressing function, and antibacterial action. In addition, since rare sugars have a large number of hydroxyl groups, they have a possibility for using as devices for expressing various functions by employing them as ligands of transition metal complexes. In recent years, it is successful that we have synthesized the supramolecular rare sugars (SRSs) from the different types of rare sugars, that is not as simple mixtures but as single crystals having superlattice structure, that can be freely controlled optical rotation. It has been found that the crystal structures of these SRSs mostly follow the Wallach rule, but do not satisfy the Wallach rule especially in the case of D,L-psicose. Therefore, we have investigated the single crystal X-ray structural analyses of SRSs, obtaining the detailed crystal structure data, and analyzed the intermolecular interaction between their sugar molecules in the



crystal by means of the DV-X α molecular orbital calculation. According to our detailed analysis of the research, calculating the intermolecular interaction revealed that the stability of the intermolecular interaction in the crystal can not be explained only by following the simple Wallach rule. Specifically, for example in the case of D,L-psicose, the total energy in a crystal can be stabilized by aligning the polarization vectors of the molecules, and as a result, we have clarified that the SRSs crystal structure can be stabilized, even if it does not follow the simple Wallach rule.

Keywords: supramolecule, rare sugar, crystal structure

Topic: DV-Xa Method

ABS-6 X-ray Absorption Spectroscopy and Its Simulation for Some Metal Oxides

Deok-Yong Cho

Department of Physics, Chonbuk National University

X-ray absorption spectroscopy (XAS) is an easy and nice method to examine the chemical and local structural properties of materials which measures the X-ray absorption coefficients with various incident X-ray energies. XAS can reveal the atomic coordination and the intersite or intrasite electron-electron interactions for each atomic species, so that it can be commonly employed for identifying the local electronic structures of condensed matter.

In this talk, I would like to overview the general principles of XAS for the local structural identification, and showcase some examples of theoretical considerations for metal oxide systems including manganites, double perovskite cobaltites and hafnia-zirconia. For the simulation of soft XAS spectra, the concept of configuration interactions with atomic multiplets is employed. For the simulation of hard XAS spectra (so-called XANES), the ab-initio real space multiplet scattering calculation (FEFF) is employed.

Keywords: X-ray absorption spectroscopy, XANES, FEFF, Configuration interactions, Atomic multiplets

ABS-7 **Study on The Higher-Order Structure and Hydrogen Bonding of Biodegradable Polymer by Low-Frequency Vibrational Spectroscopy**

Dian Marlina (a, b)*, Harumi Sato (a)

a) Graduate School of Human Development and Environment, Kobe University
3-11, Tsurukabuto Nada-ku, Kobe 657-8501, Japan
b) Fakultas farmasi, Universitas Setia Budi
Jalan Letjend Sutoyo Mojosongo, Solo 57127, Indoneisa
*157d841d@stu.kobe-u.ac.jp

Synthetic polymers, including plastic products, are known as one of the most important materials in modern life. They have been widely used in various fields such as energy, information technology, medical, living, and environment. However, most of them cannot be decomposed in the environment causing environmental problems and harmful for living organisms. More than that, the excessive use of synthetic polymer will result in the depletion of irreplaceable natural resources since most of them are made from petrochemical materials. Therefore, development of new biopolymers is expected as a material to replace fuel-based polymers. Among them, chitosan is abundant in nature. Development of polymers with chitosan contributed to solve the environmental and resource depletion problems. It is getting more attention as a new material that mitigate energy waste. Chitosan is a polysaccharide produced by deacetylation of chitin and is extensively used in drug delivery applications. Chitosan is difficult to use industrially because it has high glass transition temperature (T_g) and high crystallinity, and exhibits high elastic modulus and low strain fracture. Therefore, it is necessary to improve their physical properties by blending



or copolymerizing with other polymers. Intermolecular hydrogen bonding between chitosan and other polymer molecules is expected to play an important role in stabilizing the higher-order structure. The information related to this interaction can be observed in the low-frequency region (3.3-330 cm⁻¹) using terahertz (THz) and low-frequency Raman spectroscopy. Since low-frequency vibrational spectra are derived from intermolecular and intramolecular interactions, THz and low-frequency Raman spectroscopies can be powerful tools to reveal the formation of higher order structures of not only homopolymer but also polymer blends. In this study, by analysing this low-frequency vibrational spectra, we attempted to elucidate the change in higher-order structure and intermolecular hydrogen bonding formed by polymer blends of chitosan and other polymers.

Keywords: Chitosan, higher-order structure, hydrogen bonding, low-frequency vibrational spectroscopy

ABS-8 **Systematic First-Principles Calculations of Charge Transfer Transitions of Transition Metal Ions (Sc³⁺, Ti³⁺, V³⁺, Cr³⁺, Mn³⁺, Fe³⁺) in α -Al₂O₃ with Structural Optimization**

Shota Takemura¹, Mega Novita², Kazuyoshi Ogasawara¹

1- Department of Chemistry, Kwansei Gakuin University

2- Faculty of Engineering and Informatics, Universitas PGRI Semarang

The position of the impurity level within the band gap is very important in optical materials. It can be estimated by calculating the charge transfer transition energy from the valence band to the impurity levels (Ligand to Metal Charge Transfer: LMCT). Recently, we successfully reproduced the LMCT energies and their experimental trend of transition metal (TM) ions in α -Al₂O₃ using the DVME method considering the configuration-dependent correction and the lattice relaxation based on the Shannons crystal radii. We also clarified that the LMCT energy is very sensitive to the bond length. In this work, in order to calculate the LMCT energies for various trivalent TM ions in α -Al₂O₃ using more accurate model clusters, we performed structural optimization using the CASTEP code and constructed the optimized TMO₆⁹⁻ and TMA₁₃O₆³⁰⁺ clusters (TM = Sc³⁺, Ti³⁺, V³⁺, Cr³⁺, Mn³⁺, Fe³⁺). As a result, the trend from Sc to V was reproduced better by calculations using the optimized clusters. The differences between the LMCT energies of the optimized clusters and those of the relaxed clusters based on the crystal radii originate from the multiplet splitting of the LMCT states.

Keywords: Charge transfer transition, Multiplet, Structural optimization



ABS-9 **Prediction of 4f²→4f15d1 transition energy of Pr³⁺ in fluorides based on first-principles calculations and machine learning**

Hayato Obata

Department of Chemistry, Kwansei Gakuin University

The 4f²-4f15d1 transition energies of Pr³⁺ in fluorides are utilized for various optical materials such as solid-state lasers, phosphors, and scintillators. Therefore, it is important to predict such energies of unknown materials for theoretical design of novel optical materials. In this study, we tried to predict the 4f²-4f15d1 transition energies of Pr³⁺ in fluorides based on first-principles calculations and machine learning. The first-principles calculations were performed based on the relativistic discrete variational multi-electron (DVME) method using the model clusters composed of the central Pr³⁺ and the anions closer than the nearest cation. Although the calculated 4f²-4f15d1 transition energies of Pr³⁺ in fluorides showed a relatively good correlation with the experimental ones, the theoretical values tend to be overestimated by ca. 2 eV. In order to improve the accuracy of the prediction, we used the calculated transition energies as an attribute for machine learning. As a result, the regression formula to predict the 4f²-4f15d1 transition energy of Pr³⁺ in fluorides has been derived by machine learning using the theoretical 4f²-4f15d1 transition energy as well as some other electronic and structural parameters as the attributes. The accuracy of the prediction was significantly improved compared to the simple first-principles calculations

Keywords: The 4f²-4f15d1 transition energies of Pr³⁺ in fluorides, machine learning

ABS-10 Optimization of first-principles calculation conditions of multiplet energies in Fe³⁺ and Co³⁺ in α -Al₂O₃

Judo Fusamoto, Kazuyoshi Ogasawara

Department of Chemistry, Kwansai Gakuin University

The prediction of multiplet energies of dn ions such as transition metal ions in crystals is important for the development of novel optical materials. The first-principles calculations of the multiplet energies can be performed by the discrete variational multi-electron (DVME) method which is based on the configuration-interaction (CI) approach. Although the multiplet energies calculated by CI calculations are generally overestimated, the overestimation can be corrected by considering the configuration-dependent correction (CDC) and the correlation correction (CC). For example, the experimental multiplet energies of V³⁺ and Cr³⁺ in α -Al₂O₃ were well reproduced by first-principles calculations considering CDC-CC. However, for d⁵ and d⁶ ions, the conditions of CDC-CC such as the considered transition processes have not been established. In this study, in order to optimize the conditions of CDC-CC, we performed first-principles calculations of the multiplet energies of Fe³⁺ and Co³⁺ in α -Al₂O₃ considering CDC-CC based on various transition processes. As a result, we could optimize the conditions of CDC-CC for Fe³⁺ and Co³⁺ in α -Al₂O₃ and successfully reproduced the experimental multiplet energies. By consideration of CDC-CC, the repulsion between electrons was suppressed and the overestimation of the multiplet splittings was reduced.

Keywords: First-principles calculation, Multiplet, Transition metal

ABS-11 Cultivation of *Oreochromis niloticus* with an Environmentally friendly Bioflog System

Endah Rita Sulistya Dewi, Ary Susatyo Nugroho, Maria Ulfah

Departement of Mathematics, Sciences and Information Technology, PGRI Semarang University

Increasing population growth demands the availability of continuous food. Aquaculture is one of the food production sectors that has the highest growth rate in the world. Facing this opportunity aquaculture is faced with several challenges related to the limited natural resources such as water and land, and aquaculture waste disposal. Furthermore, the application of best aquaculture practices in aquaculture product certification requires an environmentally friendly aquaculture practice. *Oreochromis niloticus* is a kind of freshwater consumption fish and is now a popular pet fish in freshwater pools in Indonesia. This research aims to increase aquaculture production in *Oreochromis niloticus* maintenance through an environmentally friendly Bioflog system. This research method was designed in form of Research and Development. The results showed that the operation of ponds using minimal water exchange, the development of dense microbial populations and management of microbial populations through C/N ratio adjustments could control the concentration of inorganic nitrogen in water. Bacteria in ponds form bioflog, producing microbial proteins that make it possible to recycle unused feed proteins, there by minimizing the occurrence of pollution.

Keywords: Bioflog, *Oreochromis niloticus*, Environmentally Friendly

ABS-12 **Prediction of 4f⁷→4f⁶5d¹ transition energies of Eu²⁺ in oxides based on first-principles calculations and machine learning**

Hiroyuki Hori, Shota Takemura, Kazuyoshi Ogasawara

Department of Chemistry, Kwansei Gakuin University

Eu²⁺ ions are utilized as luminescent ions in solid-state lasers and phosphors. Therefore, the prediction of the 4f⁷→4f⁶5d¹ transition energy of Eu²⁺ in crystals is important to develop novel luminescent materials. In this work, we tried to predict the 4f⁷→4f⁶5d¹ transition energy of Eu²⁺ in oxides using first-principles calculations and machine learning. The first-principles calculations were performed based on the discrete variational multi-electron (DVME) method using small clusters composed of Eu²⁺ and all anions closer than the closest cation. Although the calculated 4f⁷→4f⁶5d¹ transition energies and the experimental ones showed some correlation, the theoretical values tend to be larger than the experimental ones by ca. 1.5 eV. Since machine learning enables one to create a predictive model of an output based on attributes, we tried to create a predictive model of the 4f⁷→4f⁶5d¹ transition energy of Eu²⁺ in oxides by machine learning using the calculated 4f⁷→4f⁶5d¹ transition energies and other electronic and structural parameters as the attributes. The obtained predictive model significantly improved the correlation between the predicted 4f⁷→4f⁶5d¹ transition energies and the experimental ones.

Keywords: Machine learning, Multiplet, 4f-5d transition



ABS-13 Lattice Relaxation Effects on the Multiplet Energies of Ruby Under Pressure using One-Electron Calculations

Mega Novita (a*), Nur Cholifah (b), Kazuyoshi Ogasawara (c)

a) Faculty of Engineering and Informatics, Universitas PGRI Semarang, Jl. Sidodadi-Timur No.24 Semarang, Central Java 50232, Indonesia

*novita@upgris.ac.id

b) Lembaga Penelitian dan Pengabdian Kepada Masyarakat, Universitas PGRI Semarang, Jl. Lontar No.1 Semarang, Central Java 50232, Indonesia

c) School of Science and Technology, Kwansei Gakuin University, 2-1 Gakuen Sanda, Hyogo 669-1337 Japan

Up to recently, it has been difficult to calculate the multiplet energies of compounds using one-electron approach. Since it only considers one electron and one nuclei, the interaction among the electrons are neglected. Previously we have successfully estimated the 2E and 4T₂ levels of ruby at 0 pressure using one-electron approach based on Ohnisi's method. We also compare the result with those calculated by many-electron approach. The one-electron approach carried out in this work is a first-principles Discrete Variational-X α (DV-X α) calculation. In this work, we want to perform similar study not only at 0 pressure but up to 110 GPa. We estimated the lattice relaxation effect due to the Cr³⁺ substitution and due to the applied pressure by using two different methods i.e., Shannon's crystal radii and geometry optimizations. Two different types of model cluster consisting of 7 and 63 atoms will be used. The 4T₂ level is simply estimated by the crystal field splitting (10Dq). Whereas the 2E level is estimated by the barycenter of t_{2g}³ configuration.

Keywords: pressure, first-principles, CASTEP, a-Al₂O₃



ABS-14 Optical properties of Co³⁺ doped in α -Al₂O₃ with Considering Lattice Relaxation Effect

Mega Novita (a*), Duwi Nuvitalia (b), Nur Cholifah (c),
Kazuyoshi Ogasawara (d)

a) Faculty of Engineering and Informatics, Universitas PGRI Semarang, Jl. Sidodadi-Timur No.24 Semarang, Central Java 50232, Indonesia

*novita@upgris.ac.id

b) Faculty of Mathematics, Natural Science and Information Technology Education, Universitas PGRI Semarang, Jl. Sidodadi-Timur No.24 Semarang, Central Java 50232, Indonesia

c) Lembaga Penelitian dan Pengabdian Kepada Masyarakat, Universitas PGRI Semarang, Jl. Lontar No.1 Semarang, Central Java 50232, Indonesia

d) School of Science and Technology, Kwansai Gakuin University, 2-1 Gakuen Senda, Hyogo 669-1337 Japan

In this work we estimated the optical properties of α -Al₂O₃:Co³⁺. We constructed model clusters consisting of 7 and 63 atoms. The lattice relaxation effects due to the Co³⁺ substitution were calculated using Shannon's crystal radii method and geometry optimizations in the Cambridge Serial Total Energy Package (CASTEP) method. The one-electron Discrete Variational-X α (DV-X α) method was used to estimate the molecular orbital energies, while the many-electron Discrete Variational Multielectron (DVME) method was used to estimate the d-d absorption spectra.

Keywords: Co³⁺, α -Al₂O₃, first-principles, DV-X α , DVME



ABS-15 Effects of Transition Metal Doping to Cerium-Oxides in Thermal Reduction Reaction

Takaki Nishimura¹, Yuki Tsutsumi², Tomohiko Ishii²,
Tatsuya Kodama³, and Genta Sakane⁴

¹ Master course student, ^{1,2} Department of Advanced Materials Science, Graduate School of Engineering, Kagawa University, 2217-20 Hayashi-cho, Takamatsu, Kagawa 761-0301, Japan,
E-mail : s18g576@stu.kagawa-u.ac.jp, Tell : +81-87-864-2414

³ Faculty of Engineering, Niigata University, 8050 Ikarashi 2 nocho, Nishi-ku, Niigata, 950-2181, Japan

⁴ Department of Chemistry, Faculty of Science, Okayama University of Science, 1-1 Ridaicho, Kita-ku, Okayama 700-0005, Japan

Considering our lives, novel energy production is so required, and many scientists have been investigated this problem from a lot of views. One of the ways for the new energy productions is the solar power thermochemical production by an oxidation-reduction reaction of the ceramics CeO₂ (ceria). In this research, the ceria is used for obtaining an H₂ gas in vapor reaction in high-temperature thermal equipment which has a heat production system consisting by focused sun light. Nowadays technology has been also developed to keep a human living, not to harm the Earth environment. From these points, it is realized that how the H₂ gas is important for our lives, so we are trying to improve the thermochemical H₂ gas production by means of a ceria as a catalyst. Our purpose in this study is, to point out what effects occur in doped ceria's thermal reduction (TR) reaction by using DV-X α method. Comparing the



models between the pure ceria and the Mn doped ceria, we have realized the difference by the reaction. In this appearance, one of the considerations is a view from p-DOSs (partial-Density of States), in which there are different O-2p DOSs by different crystal structures. From this view, we considered that the ceria structure would become more stable and more flexible to cut the Ce-O bonding to keep the stable cyclic thermal reaction in Mn doped ceria.

Keywords: Ceria, H₂ gas, Thermal reduction (TR) , Mn doped ceria

ABS-16 Study of LCAO-MO calculation by using completely numerical basis functions

Katsumi Nakagawa

MO BASICS Research

Introduction)

DV-X α method uses numerical basis functions adaptable for the circumstance in a molecule. But they are calculated based on spherically symmetric potential for each atom. Newly proposed method inherits DV-X α 's merit but isn't restricted by the premise of atomic spherical symmetry. Aim of the study). This method is being developed as a powerful tool to calculate MOs under complicated outer potential, especially vector potential.

Method)

Basis functions are calculated by operations of matrices representing physical quantities and vectors representing wave functions. Analytical assists, like spherical harmonic functions, are not necessary at all. Hartree-Fock-Slater equations for molecules can be expressed numerically even for arbitrary outer potential and solved easily as eigenvalue problems of matrix. But basis functions for an atom are defined only on sample points proper to that atom. On the other hand, basis functions need to be defined on all sample points to calculate Fock matrices for LCAO-MO calculation. The author will explain some techniques to pass over this gap at this presentation.

Results)

LCAO-MO calculations were tested within real wave functions. Total energy vs interatomic-distance curves for



N₂, F₂ and CO were calculated and showed minima near experimental equilibrium distances. Dipole moments of CO, O₃ and NH₃ were calculated and matched semi-quantitatively experimental moments. Calculations of basis functions were heavy but showed O(n) dependence.

Conclusion)

It was demonstrated that basis functions calculated by matrix operation can be used well for LACO-MO calculation. When wave functions are extended to complex, this method will serve as a useful tool to treat MOs under complicated magnetic fields.

Keywords: DV-X α method, numerical basis functions, vector potential, dipole moment

ABS-18 Controlling of the Spin State by Jahn-Teller Distortion in Octahedral Hexa- Coordinate Metal Complex

T. Oguma ,a Y. Huzikawa ,a D. Tsuneda, a Y. Tsutsumi, a M. Nakano, a T. Ishii, a G. Sakane, b

a. Department of Advanced Materials Science, Faculty of Engineering, Kagawa University

2217-20 Hayashi-cho, Takamatsu, Kagawa 761-0369 JAPAN

b. Department of Chemistry, Faculty of Science, Okayama University Science

c. Department of Chemistry, School of Science and Technology, Kwansai Gakuin University

Metal complexes exhibit significant physical properties such as electrical conductivity, magnetism and optical properties. They can be controlled by changing the ligand field splitting of the d-orbital of the central metal ion. We have investigated the reversible spin crossover phenomenon by changing the ligand field splitting in six-coordinated metal complexes ML₆. In this study, we focused on the change of the spin states by JahnTeller distortion in the case of an octahedral hexa-coordinated metal complexes ML₆, involving 2 types of different ligand molecules. In addition, we have used DV-X α and DVME methods in order to calculate the electronic state. In order to reproduce the effect of the JahnTeller distortion in the calculation, a uniaxial anisotropic distortion has been applied into the octahedral hexa-coordinate metal complex. The purpose of this research is to study the mechanism and phenomenon of the spin state changing by studying the spin crossover phenomenon.

Keywords: Complex



ABS-19 Single-Crystal X-Ray Structural Analysis and Electronic Structures Calculation of Rare Sugar

Bui Quoc Huy, Tomohiko Ishii, Norito Taniguchi, Natsumi Nagayama, Genta Sakane

Department of Advanced Materials Science, Faculty of Engineering, Kagawa University Hayashi-cho, Takamatsu, Kagawa 761-0396, JAPAN.

Department of Chemistry, Faculty of Science, Okayama University of Science 1-1 Ridaicho, Kita-ku, Okayama 700-0005, JAPAN

There are two types of mono-saccharide sugars in nature, which are natural monosaccharide and rare sugar. The sugar that exists in extreme few amount in nature is called as rare sugar e.g. D-psicose, D-allose. The reasons we choose the rare sugar as a research target are because the rare sugar has almost no calories and is very useful for diabetic and dieter patients. Specifically, D-psicose is a no-calorie rare sugar with the sweetness of about 70 % of a sucrose. It has been confirmed that taking the D-psicose with meals suppresses the rise of sugar into the blood. Further, it is very effective in improving and preventing the diabetes. In order to investigate a research on the rare sugars, the first step must be to obtain the correct molecular structure. However, few crystal structures of rare sugars have been reported up to now. Therefore, the purpose of this research is to analyze the single-crystal X-ray structure of unknown rare sugar 'L-glucose'. In addition, the DV-X α molecular orbital method is used to compare the differences in a hydrogen bonding between D-glucose and L-glucose, and to make a novel supramolecular rare sugar.

Keywords: Rare Sugar, D-psicose, L-glucose, Supramolecule, Hydrogen bonding.



ABS-20 Data Mining Application with Fuzzy Logic Method for Monitoring Vehicle Position Based on Android

Munaf Ismail, Sri Arttini Dwi Prasetyowati, Ida Widiastuti

Universitas Islam Sultan Agung

Cases of vehicles theft in the Central Java region occurred 869 cases, the percentage was 57.86% against other crimes. To prevent vehicles theft, the police need public participation by increasing awareness of their vehicles. Vehicle theft can occur due to owner negligence and lack of safety locks on the vehicle. This study aims to create data mining application as monitoring vehicle position. The transmitter WiFi 2.4 GHz placed on a vehicle, then measured the RSSI signal by an android application based on fuzzy logic. Indication of theft is that the vehicle is positioned away from the owner, so the signal RSSI is less than -90 dBm. Fuzzy logic monitors this change in distance. When the membership function is lost, fuzzy logic instruct android to activated the alarm . Fuzzy logic was able to make decision give a alarm to user when RSSI signal less than -90 dBm. RSSI signal is less than -90 dBm when the distance is 20 meters if there is a wall barrier, without obstacle the distance is up to 80 meters. RSSI signal is influenced by the measurement environment.

Keywords: Data mining application, Fuzzy Logic, Monitoring vehicle position

**ABS-21 An elaborate response in molecular random network -
Approach for information processing functionality-**

Megumi Akai-Kasaya

An animate brain consists of interconnected complex biological networks of locally active components. The informational processing exploits massive parallelism with low energy consumption, utilizing the multiplicity and stochasticity of signals. Recently, utilization of disordered random networks of inanimate matter for computation or informational processing has started to be proposed. Here we introduce two nanomaterial networks, which show complex response showing potency for advanced future signal processing.

We succeeded in showing experimental evidence to prove Coulomb blockade taking place on Two-dimensional organic conducting polymer films. Electrical conductivity of hexylthiophene monolayer showed temperature-dependent threshold voltage in temperature range of 150K—4K and increased in power law of the current-voltage beyond the threshold. The onset of two-dimensional Coulomb blockade in the organic thin film was theoretically verified through the calculation of the delocalization of electric charge in the molecular film. This system meets the criteria to be a suitable building block for the physical realization of a cellular neural network. We further found a neuromorphic device that consist of extremely dense and random single-walled carbon nanotube (SWNT) complexed with polyoxometalate molecule generating spontaneous spikes similar to nerve impulses of neurons. A simulation calculation of the random molecular network model, which are able to store electric charges, replicate spikes generated from the network. This molecular model would very likely become a component of reservoir computing that is

anticipated as next-generation artificial intelligence. These results indicate the possibility that complex functional networks can be constructed using molecular devices, and contribute to the development of neuromorphic devices.

Keywords: Nanomaterial Molecular Device Neuromorphic Neural Network

ABS-22 Prediction of emission energy of Cr³⁺ in oxides based on first-principles calculations and machine learning

Tsubasa Hori, Kazuyoshi Ogasawara

Kwansei Gakuin University

Transition-metal ions with d³ electronic configuration such as Cr³⁺ and Mn⁴⁺ are utilized as emission centers in various luminescent materials such as ruby (Cr³⁺-doped α -Al₂O₃) or alexandrite (Cr³⁺-doped chrysoberyl). Since it is difficult to investigate the multiplet energy levels of Cr³⁺ in a wide variety of oxide crystals experimentally, the theoretical prediction of the energy of the emission level (2E_g) is indispensable for the efficient development of novel Cr³⁺-doped luminescent materials. In this study, we predicted the multiplet energy of the emission level (2E_g) of Cr³⁺ in oxides by first-principles calculations based on the discrete variational multi-electron (DVME) method using relatively small clusters consisting of seven atoms. However, the correlation coefficient between the calculated 2E_g energy and the experimental ones was relatively low (0.52). In order to improve the accuracy of the prediction, we also performed a machine learning modeling. By using the calculated 2E_g energy and the other electronic and structural parameters as the attributes, we created the predictive model of the 2E_g energy of Cr³⁺ in oxides. The predicted 2E_g energies are in good agreement with the experimental ones, showing a high correlation coefficient of 0.92.

Keywords: Cr, DV-X α

ABS-23 Consideration of Super Atomic Hibrid Orbital in Super Atom M@Al₁₂ (M = Be, Mg, Ca, Sr)

Teruyuki Toba¹, Ishii Tomohiko¹, Syoya Yamasaki¹, Sakane Genta²

1 Department of Advanced Materials Science, Graduate School of Engineering, Kagawa University, 2217-20 Hayashi-cho, Takamatsu, Kagawa 761-0301, Japan

2 Department of Chemistry, Faculty of Science, Okayama University of Science, , 1-1 Ridaicho, Kita-ku, Okayama 700-0005, Japan

We introduce SuperAtom, which is called a modern alchemy and has been studied around the world. The SuperAtom is a nanocluster composed of multiple atoms, for example Al₁₃. But very interestingly, the "SuperAtom" can mimic the chemical properties of "atom", for example, the Al₁₃ exhibits the halogen atom Cl. If this SuperAtom is constructed to an appropriate size and can be synthesized freely, there is the possibility to be placed as an "artificial atom" in a three-dimensional periodic table such as the conventional two-dimensional periodic table. In recent years, the exhaustion of rare metals has become a big problem, so that in the next generation of materials and devices, the SuperAtoms play as an important role, and it is expected to be used for various materials or as a substitute for expensive metal catalysts such as Pt. From these points of view, the SuperAtoms have been studied all over the world up to now, but there is not much studies on their electronic structures. Therefore, in this study, we have analyzed the electronic state of the chalcogen-inclusive type SuperAtom M@Al₁₂ (M = Be, Mg, Ca, Sr) by means of DV-X α molecular orbital calculation. Our analysis revealed that the electronic configuration of the SuperAtom, which



changed only the central atom, depends largely on the kind of doped atom. And we can also confirm the “SuperAtomic hybrid orbital” composed of SuperAtomic orbitals, and chalcogen characteristics.

Keywords: SuperAtom, SuperAtomic Hibrid Orbital

ABS-24 **Mechanism of Optical Rotation of Amino Acids Using Electronic State Calculation**

Shuuichi Katumoto,¹ Manami Mimura,¹ Tomohiko Ishii,¹
and Genta Sakane,²

¹Department of Advanced Materials Science, Faculty of Engineering, Kagawa University,

²Department of Chemistry, Faculty of Science, Okayama University of Science

E-mail : s19g558@stu.kagawa-u.ac.jp

Because of the chiral nature of the building blocks of living matter, an optical phenomena associated with the chirality constitute an important topic in physical chemistry. The specific optical rotation, which is a parameter for the characterization of the natural optical activity, depends strongly on the conformation of the molecule. In this study, we investigated the dependence of the optical rotation on a molecular conformation in the gas phase by calculating the electronic states of seven kind of chiral amino acids using the DV-X α method. As a result, we can confirm the existence of an antibonding orbital on the side chain and the optical rotations are strongly related.

Keywords: Chiral Optical Rotation



ABS-25 Generation of Two-Dimensional Spectrochemical Series in Tetrahedral Complexes

Takuhiro Yamamoto, Takeyoshi Oguma, Momoe Nakano,
Akira Tsuneda, Yuki Tsutsumi, Tomohiko Ishii

Department of Advanced Materials Science,
Graduate School of Engineering
Kagawa University

The physical properties of metal complexes that are present in many places are due to the ligand field splitting of the d-orbital of the central metal ion, and the series representing the relationship of the ligand field splitting is called a spectrochemical series. This spectrochemical series represents the relationship of the ligand field splitting in the octahedral hexa-coordination structure, but no equivalent index has been shown in the tetrahedral four-coordination complex. In this study, we have investigated the creation of a novel spectrochemical series in tetrahedral four-coordinated metal complexes. As a result, it is possible to accurately understand the ligand field splitting of the tetrahedral four-coordinated metal complex, and to more effectively apply the energy for obtaining a spin crossover phenomenon in which the magnetic spin state reversibly changes. As a research flow, first, a tetrahedral structural model is created, and then the ligand field splitting is calculated using the DV-X α method to create a spectrochemical series. As a result, we succeeded in creating a spectrochemical series of tetrahedral metal complexes. It has been found that the spectrochemical series of tetrahedral four-coordination complexes tend to hold as they are in octahedral hexa-coordination complexes to some extent, but there are cases where the series does

not hold according to the peculiarity in the case of the four-coordinated metal complex.

Keywords: Spectrochemical Series, Four Coordination Complex



ABS-26 Effect of Molecular Distortion of Ligand Field Splitting in Five-Coordinated Metal Complex

Yohiki Fujikawa, Takeyoshi Oguma, Takuhiro Yamamoto,
Yuki Tsutsumi, Dan Tsuneda, Momoe Nakano, Tomohiko
Ishii, Genta Sakane

Department of Advanced Materials Science,
Graduate School of Engineering,
Kagawa University

Metal complex indicates the significant physical properties such as magnetism, electrical conductivity and an optical property due to the electronic interaction between the ligand molecule and the central metal ion. We have evaluated the magnetism of the metal complex by controlling the ligand field splitting, which is the energy splitting of the d-orbitals. The magnetic phenomenon of such compounds are realized as a result of the electronic state based on the transition metal coordination complexes, and which can be switched between high spin (HS) and low spin (LS), that is called a spin-crossover (SCO) phenomenon. Our group have been investigated the way how to control the spin state according to the concept of a distortion. The five-coordinated metal complex ML₅ has been experimentally confirmed to be a molecular distortion called the Berry pseudorotation (BPR). In the BPR process, both the axial and equatorial ligands can move at the same rate of increasing the angle between the other axial or equatorial ligands in five-coordinated metal complex. The classical BPR mechanism changes its molecular symmetry between two independent trigonal bipyramidals (TBPs) of D_{3h} symmetry via a square pyramidal (SP) of C_{4v} symmetry. The distortion parameters 5 are proposed in order to represent the distortion of the BPR process quantitatively. In this work, we predicted the possibility of the SCO

phenomenon in intramolecular exchange mechanism through the BPR process in a five-coordinated metal complexes.

Keywords: Spin-crossover, Berry pseudorotation, Five-coordinated Metal complexes

ABS-27 Single crystal X-ray structural analysis of sugar molecules and electronic theory

Norito TANIGUCHI⁽¹⁾, Tomohiko ISHII⁽¹⁾, Bui Quoc Huy⁽¹⁾,
Natsumi NAGAYAMA⁽¹⁾, Genta SAKANE⁽²⁾

(1)Department of Advanced Materials Science, Faculty of Engineering, Kagawa University

(2)Department of Chemistry, Faculty of Science, Okayama University of Science

Rare sugar is a monosaccharide that there are trace amounts in nature, natural sugars present in large amounts in nature is referred to as a natural monosaccharide. Recently, the isomerase called D-tagatose-3-epimerase (DTE) was discovered, and it has a possibility to synthesize quantity of the rare sugar from a natural sugar through the enzymatic reaction. And all of hexoses can be synthesized by four kinds of enzymatic reactions (oxidoreductase reaction, aldose isomerase reaction, aldose reductase reaction and DTE reaction). In this study, single-crystal X-ray structure analysis of a sugar alcohol and its derivative have been performed in order to determine the absolute coordinates of these sugar molecules. We give an electronic state calculation by means of the DV-X α method using the obtained data of the absolute coordinates. We also consider that mechanism of the enzymatic reaction and intermolecular energy properly such as a hydrogen bond in order to synthesize the supramolecular rare sugars (SRSs) which can be controlled their structures and the hydrogen bonds.

Keywords: supramolecule rare sugar, hydrogen bond, D-tagatose-3-epimerase



ABS-28 Structural changes in graphene oxide induced by silver nanoparticles – TERS study

Prompong Pienpinijtham (a,b,c,*), Sanpon Vantasin (a), Oraporn Wong-u-ra (b), Yasutaka Kitahama (a), Sanong Ekgasit (b,c) and Yukihiro Ozaki (a,*)

a) Department of Chemistry, School of Science and Technology, Kwansai Gakuin University, Sanda, Hyogo 669-1337, Japan.

b) Sensor Research Unit (SRU), Department of Chemistry, Faculty of Science, Chulalongkorn University, Pathumwan, Bangkok 10330, Thailand.

c) Research Network NANOTEC-CU on Advanced Structural and Functional Nanomaterials, Faculty of Science, Chulalongkorn University, Bangkok 10330, Thailand.

This work has investigated structural changes of graphene oxide (GO) in silver/graphene oxide (AGO) nanocomposites using tip-enhanced Raman scattering (TERS), which is a combination technique of surface-enhanced Raman scattering (SERS) and scanning probe microscopy (SPM). Based on SERS, a Raman spectrum of trace molecules adsorbed on rough metal surfaces/nanoparticles can be collected. On the other hand, nano-scaled topology of samples can be bestowed by SPM. As a result, TERS provides benefits of those two techniques, enabling one to determine molecular information of diminutive samples at specific nano-scaled areas. Because of this markedly high spatial resolution of the TERS technique, the structural changes of GO in AGO can be achieved by constructing line-profile TERS spectra straight from the center of silver nanoparticles (AgNPs) on GO layers. The results show evidences that AgNPs cause shortening of C–C bonds beneath AgNPs, flattening of GO layers, and critical bending on GO layers. Additionally, a connection of carbon atoms via

C–C network subsequently expands structural changes with the distance of 200–250 nm from the center of AgNPs, even though this distance is larger than the size of AgNPs. The proposed model of GO structural changes opens new understanding about changes in properties from GO to AGO nanocomposites, which will contribute to a development of advanced nanostructures /nanocomposites in the near future.

Keywords: Tip-enhanced Raman scattering (TERS), Graphene oxide (GO), Silver nanoparticles (AgNPs), Silver/graphene oxide (AGO) nanocomposites, Structural changes.

ABS-29 Week Hydrogen Bonding of Polymers Studied by Terahertz and Low-frequency Raman Spectroscopy

Harumi Sato

Kobe University

The absorption peaks observed in the terahertz (THz) region reflect a higher-order structure, crystalline structure, and intermolecular interactions such as hydrogen bonding. Therefore, THz spectroscopy and low frequency Raman spectroscopy are unique techniques for analysing higher-order conformations and intermolecular interactions in semicrystalline polymers. We have investigated changes of the higher-order structure and hydrogen bonding of several kinds of polymers such as poly(glycolic acid) (PGA), poly-(R)-3-hydroxybutyrate (PHB), polylactic acid (PLA), poly(ϵ -caprolactone) (PCL), poly (butylene succinate) (PBS), poly (ethylene terephthalate) (PET) and poly (butylene terephthalate) (PBT), and so on by THz spectroscopy and THz Raman spectroscopy with quantum chemical calculations (QCCs).

Keywords: terahertz spectroscopy, low frequency Raman spectroscopy, polymer

ABS-30 **Metal complexes with sulfur-containing ligands**

Masahiro Mikuriya and Makoto Handa

Kwansei Gakuin University and Shimane University

Organic chelate ligands are useful to develop metal-organic systems. Especially N,N,S-tridentate thiolic ligands have a great affinity for many kinds of metal ions to form various metal complexes from mononuclear to polynuclear species. On the other hand, the corresponding N,N,O-tridentate alcoholic ligands form dinuclear CuII species exclusively. We focused on these systems in the hope of obtaining new feature of metal thiolates, elucidating interesting crystal structures and spectral properties, although most thiolato complexes are diamagnetic. We obtained mononuclear and trinuclear CoIII species with 2-[(3 aminopropyl) amino]ethanethiol. The CoIII species reacted with oxygen to form the corresponding sulfinato CoIII species. Their conversion was monitored by UV-vis spectra and DFT calculations were performed for the thiolate and sulfinato complexes. We also obtained molybdenum complexes. We will discuss these features based on the crystal structures.

Keywords: metal complexes, thiolic ligand, cobalt complexes, molybdenum complexes

ABS-31 Contribution of edge-carbon atoms and non-benzenoid rings in graphitic carbons to π^* peak profiles in CK-XANES

Yasuji Muramatsu and Yuma Hirai

University of Hyogo

To analyze the local structure of non-benzenoid rings in graphitic carbons by soft X-ray absorption spectroscopy, CK-XANES of aromatic compounds having pentagonal or heptagonal rings were theoretically analyzed by the first principle calculations. Edge-carbon atoms split the π^* peak into two peaks in CK-XANES. Pentagonal rings in polycyclic aromatic hydrocarbon structure disturb delocalization of π electrons in hexagonal rings. Hence, pentagonal rings split the π^* peak into two peaks in CK-XANES. It is also confirmed that heptagonal rings play a role of a hole in hexagonal carbon layer, and the heptagonal carbon atoms can be regarded as edge carbon atoms. Hence heptagonal rings also split the π^* peak into two peaks. Consequently, edge-carbon atoms and non-benzenoid rings make the π^* peak wider than hexagonal rings.

Keywords: Soft X-ray, XANES

ABS-33 Smart Aquaponics Design Using Internet of Things Technology

R Hafid Hardyanto (1), Prahenusa Wahyu Ciptadi (2)

(1)hafid@upy.ac.id, Informatics Department, Universitas PGRI Yogyakarta

(2)nusa@upy.ac.id, Informatics Department, Universitas PGRI Yogyakarta

Smart Aquaponic is a system of planting plants and maintaining fish in one container with integrated Internet of Things technology. Aquaponic is the process by which plants utilize nutrients derived from fish feces which, if left in the pond, will become toxic to the fish. Plants function as a filter of vegetation which will break down these toxic substances into substances that are not harmful to fish. Plants also supply oxygen to the water used to maintain fish. With this cycle there will be a mutually beneficial cycle between plants and fish. This system is very profitable, because the land used will not be too large. The purpose of this study was to develop a smart aquaponic system using Internet of Things technology. In this system aquaponics are equipped with sensors that function as monitors and actuators. Actuators can be controlled by users using smart phones. Sensor monitoring can be observed by users using smart phones. The method used in this study uses the Microsoft Solution Framework (MSF) with the waterfall system development method and Object Oriented Development (OOD) method. Stages in this study includes problem identification, preliminary planning and design, piloting and implementation. Conclusion: This smart aquaponics design is ready to be implemented in small scale trials.

Keywords: Aquaponic, Internet of Things, Smart Aquaponic System

ABS-34 **Stability of carbon-vacancy complexes in α -Fe**

Masataka Mizuno, Waka Yamada, Kazuki Sugita, Hideki Araki

Division of Materials and Manufacturing Science, Graduate School of Engineering, Osaka University, Osaka University 2-1 Yamadaoka, Suita, Osaka 565-0871, Japan

[Introduction & aim of the study] Recent experimental and theoretical studies suggest that the carbon-vacancy complexes are formed in Fe-C alloys. In this work, we have performed first-principles calculations to evaluate the stability carbon-vacancy complexes in α -Fe. For comparison, nitrogen-vacancy complexes have been investigated. [Method] We employed VASP code to obtain formation energies and stable structures of interstitial solutes X_n (X = C and N, n = 1 to 6) and vacancy (V) complexes. The chemical bonding was analyzed using DV-X α molecular orbital calculations.

[Results & conclusion] The formation energies of carbon-vacancy complexes have been calculated with reference to the total energy of an isolated interstitial atom and an isolated mono vacancy. In the case of carbon-vacancy complexes, the C₂-V complex is most stable due to the energy gain by forming the C-C bonding. On the one hand, the N₂-V complex is less stable than the N₁-V complex. This is due to the repulsive interaction between N atoms. The overlap population analysis reveals that the C 2s – C 2s anti-binding interaction is reduced by the C 2s – C 2p bonding interaction. The interaction between the N 2s and N 2p orbitals is not strong enough to cancel the N 2s – N 2s anti-binding interaction.

Keywords: Steel; defect; solute; interstitial; first-principles



ABS-35 Recent applications of discrete variational multi-electron method

Kazuyoshi Ogasawara

Department of Chemistry, Kwansai Gakuin University, 2-1
Gakuen, Sanda 669-1337, JAPAN

The discrete variational multi-electron (DVME) program is a first-principles many-electron calculation program especially designed to calculate multiplet states of transition-metal ions and rare-earth ions in crystals. It is a configuration interaction calculation program based on the discrete-variational X α cluster method. It has been applied to analyze various optical materials such as ruby, alexandrite, Mn⁴⁺-doped fluorides and RE-doped YLF. The variety of multiplet spectra such as d-d, f-f, f-d spectra, and X-ray absorption near-edge structures have been reproduced without any empirical parameters and the origins of the peaks in the spectra have been clarified based on the configuration analysis of the many-electron wave functions. The DVME method have been also utilized to create various energy-structure maps for Mn⁴⁺ and Ce³⁺ in oxides which can be used for theoretical design of novel red phosphor materials for white LEDs. Since systematic first-principles calculations are quite time consuming, we have also begun to create simple predictive models based on machine learning by using the results of the first-principles calculations as the training data. This could accelerate the creation of detailed energy-structure maps. In this presentation, I will show some of the recent applications and future perspective of the DVME method.

Keywords: First-principles calculation, multiplet, phosphor, machine learning

ABS-36 Alternative Solutions For Determining Village Funds Using Weight Product Method

Tri Hastono, Firdiyan Syah

Universitas PGRI Yogyakarta

A country is said to be progressing, if all regions of the country prosper. In Indonesia, the smallest area that runs the government function is the village. The legal basis regarding the village government itself is stated in Law No. 6 of 2014 Republic of Indonesia. In carrying out its functions as a government, the village is supported by village funds. Village funds sourced from the state budget are intended for the development and improvement of the welfare of rural communities. Village funds can go down based on the proposal submitted. The proposal for submitting village funds is processed by a work team called Tim 11. So far, the determination of village funds is still manual and less transparent. Therefore we need a system that can overcome the problem of determining village funds. This research provides alternative solutions regarding the determination of recipients of village funds using the weight product method. There are four criteria in this research, namely the number of residents in an RT area, the standard of living of an RT area, the extent of the area, the level of geographical damage in an RT area. And the object of the research chosen was Kasihan sub-district, Tamantirto village. the results obtained are a priority submission in writing the proposal submitted.

Keywords: Decision Support System;Village Funds;Weight Product

**ABS-37 DESIGN OF ELECTRICITY ENERGY SOURCES IN SEMARANG
PGRI UNIVERSITY USING OFF-GRID SOLAR PANEL
SYSTEMS**

Adhi Kusmantoro(a*), Mega Novita(b), Th.Indriati
Wardani(c)

(a)Faculty of Engineering and Information, University of
PGRI Semarang, Jl. Sidodadi Timur No.24, Central Java
50232,Indonesia

*adhiteknic@gmail.com

(b)Faculty of Engineering and Information, University of
PGRI Semarang, Jl. Sidodadi Timur No.24, Central Java
50232,Indonesia

(c)Faculty of Mathematics and Natural Sciences
Information Technology, University of PGRI Semarang, Jl.
Sidodadi Timur No.24, Central Java 50232,Indonesia

The source of electricity in each university building in PGRI Semarang is obtained from each distribution transformer or power transformer. The power transformer used to distribute electricity in the central building (GP) has a capacity of 400 KVA, while the power transformer used in the main building (GU) has a capacity of 160 KVA. In the previous study, identification of the quality of electrical power in the PGRI University Semarang building, obtained a low power data factor and a large electrical network harmonics. The problem was solved by designing and installing bank capacitors and harmonic filters. To avoid dependence on PLN electricity and not using a generator when PLN electricity goes out, the source of electrical energy is designed by using solar energy sources through Photovoltaic. Solar Panel as an important component of solar power plants, converts sunlight into electricity. Generally we calculate the maximum amount of sunlight that is converted into electricity throughout the day is 5

hours. Electricity in the morning is stored in the battery, so electricity can be used at night, where without sunlight. Based on the identification of the load in the GP is 26.12 KW, then in the design using solar energy as many as 27 solar panels are needed. As for the storage system uses 44 batteries 100Ah battery. Off-grid solar panel systems are only used when PLN electricity goes out or can be used at night. The research method used is observing the amount of electrical load, calculating the capacity of solar panels and batteries, and simulating the model using a Power Simulator.

Keywords: Solar Panel, Battery, PLTS off-grid, Charger Controller

ABS-38 Novel shape analysis method for single bioparticles in aqueous solutions

Sou Ryuzaki(a,b*), Makusu Tsutsui(c), Masateru Taniguchi(c)

a, IMCE Kyushu University

b, JST PRESTO,

*ryuzaki@ms.ifoc.kyushu-u.ac.jp

c, ISIR Osaka University

Rapid structural analysis methods for biomolecules and biomaterials consisting of single or several molecules in solution represent innovative technologies to reveal their functions because the functions strongly depend on their own structures. However, there presently exist no rapid structural analysis methods for single nanomaterials suspended in liquid environment. Nanopore sensors have been widely used to investigate the volume of particles and molecules passing through the pore by probing temporal changes in the ionic current pulses. These pulse sensors have been developed for not only size but also shape of analyte during recent years. Smaller aspect ratio defined as the ratio of the depth to the diameter and a high-speed current detection system provide greater spatial resolution, i.e. tomograms of a material passing through a nanopore. Here we will report the development of low-aspect-ratio nanopores with a spatial resolution of ca.35.5 nm and the 10 MHz-current-amplifier, resulting in realization of ultrafast time resolutions of 1.0 μ s for the tomography analysis of a material passing through a nanopore. Combining state-of-the-art technologies with multiphysics simulation methods to translate ionic current data into tomograms of nanomaterials passing through a nanopore, we have achieved rapid structural analysis of single and dabble polystyrene (Pst) beads, and bionanomaterials such

as E-colis in aqueous solutions [1]. In addition, we will also report plasmonic nanopore devices, which enable us to detect Surface-enhanced Raman Spectrum of a material inside a nanopore. The nanopore devices will be innovative technologies for the fields of nanobiodevices and structural biology. [1] S. Ryuzaki, et al., Nanotechnology 28 (2017) 15550

Keywords: nanopore

ABS-39 Design of Product Monitoring System using Internet of Things Technology for Smart Manufacturing

Marti Widya Sari ^(a*); Herianto ^(b); IGB Budi Dharma ^(c); Alva Edy Tontowi ^(d)

(a,b,c,d) Department of Mechanical and Industrial Engineering
Universitas Gadjah Mada

Yogyakarta, Indonesia

(a) Department of Informatics

Universitas PGRI Yogyakarta

Yogyakarta, Indonesia

*marti.widya.sari@mail.ugm.ac.id

The application system in the industrial era 4.0 is currently growing rapidly. One of them is the implementation of smart manufacturing in the industrial sector. In smart manufacturing, production processes can be monitored automatically and real time, starting from material selection, then the production process to the planned product. In this study discussed the design of a monitoring system on smart manufacturing based on internet of things technology. Smart technology is implemented on material scans automatically based on color sensors, then from the material it will also be known that with these materials produce certain products and require any material. In addition, the system will also automatically detect through the initial material to find out which processes will be passed. The methods used include literature studies, system requirements analysis, system design and system testing. The results of this study are in the form of a hardware system design and display of production process monitoring system applications. In the system that is made also a notification is displayed if a bottle neck occurs in the production process

Keywords: product, desgin; monitoring; smart manufacturing; internet of thing

ABS-40 **A Novel and Fast Memory Perturbation Method to Increase Exploration in Particle Swarm Optimization Algorithm**

Muchamad Taufiq Anwar, Edy Winarno, Iswatun Chasanah, Saptono Nugrohadi

Universitas Stikubank, Satya Wacana Christian University, Universitas PGRI Semarang

Particle Swarm Optimization (PSO), one of the versatile nature-inspired optimization algorithm, continue to suffer from premature convergence regardless of the rigorous amount of research. Many research had tried to address this issue but often use a complex algorithm which tax on computational time and complexity. This research introduced a novel perturbation method to mitigate premature convergence / to increase exploration while keeping the computational cost at a minimum. The particles memories (i.e the position of personal and global best) are modified by a random multiplier which in turn will 'perturb' the particles' velocity. The implementation of this novel perturbation method in early iterations had resulted in 100% success rate in finding global optima in our benchmark tests (Sphere, Schwefel 2.22, Rosenbrock, Rastrigin, and Ackley problem) – whereas the original PSO failed in all benchmark tests – without adding a significant amount of computational complexity and time.

Keywords: PSO, premature convergence, perturbation, exploration

ABS-41 BIOCONVERSION ON WASTEWATER OF SOYBEAN USING MICROBIAL FUEL CELL

Yohanes A Cahyono, Tilana Madurani, Widya F Azzahra and Retno A. S. Lestari

Chemical Engineering Department, 17 Agustus 1945 University, Pawiyatan Luhur Bendan Dhuwur, Semarang 50233, Indonesia

Microbial fuel cell (MFC) is a technology developed to obtain new sources of renewable energy to produce electricity. It can be an alternative for wastewater treatment and bioenergy producers of renewable electricity. This method requires bacteria to convert substrate in wastewater into electrical energy. The mechanism of MFC were oxidation of substrate by bacteria to produce electrons and protons at the anode. The proton in anode chamber passes through a membrane exchange to the cathode chamber, however the electrons couldn't through. It caused accumulation of electron in anode chamber and then both of electrode had a potential difference, so electron in anode chamber passed through membrane exchange to cathode chamber. In this study used dual-chambers reactors with each compartment having 8 cm \times 10 cm \times 10 cm of dimensions and 5 mm of thickness. This study was subjected to evaluate the performance of MFC in soybean washing wastewatertreatment with bacteria of EM4 to analyze the potentials production of electricity energy. The focus of this study was to evaluate the effect of time to electricity. MFC system was observed for 40 hours, measurement of voltages and electric currents performed every 2 hours. The results showed that there was potential of electricity production from soybean wastewater treatment by MFC. The maximum electricity reached in soybean wastewater

media were voltage 441 mV (at 24 h), the electric currents 170 μ A and the power density 51, 35 mW/m²(at 24 h after acclimatization). Increasing of time effect to decreasing of electricity produced.

Keywords: bioenergy, electricity, microbial fuel cell, membrane, wastewater soybean

ABS-42 Microscopic crystal field effects in impurity centers formed by the transition metal ions

Mikhail G. Brik

1. College of Sciences, Chongqing University of Posts and Telecommunications, Chongqing 400065, People's Republic of China
2. Institute of Physics, University of Tartu, W. Ostwald Str. 1, Tartu 50411, Estonia
3. Institute of Physics, Jan Długosz University, Armii Krajowej 13/15, PL-42200 Częstochowa, Poland

First-principles methods of calculations of electronic and optical properties of impurity ions in solids allow for a quantitative treatment of microscopic crystal field effects. In other words, they give a possibility of calculating the impurity ions energy levels for different geometric configurations of impurity centers, including changing chemical bond lengths and angles between those chemical bonds. From these calculations, it is possible to extract the distance dependence of the crystal field strength $10Dq$, estimate the electron-vibrational interaction constants, Jahn-Teller stabilization energy, Huang-Rhys factors, Stokes shift between the absorption and emission spectra. The obtained results can be used not only for explanation of the already existing experimental results, but for the prediction of optical properties of new materials. In this presentation several examples of such calculations will be given in detail [1-6] along with discussion of practical importance of the obtained results and their potential predictive power.

References:

- [1] M.G. Brik, K. Ogasawara, Phys. Rev. B 74 (2006) 045105.
- [2] M.G. Brik, J. Phys. Chem. Solids 68 (2007) 1341.
- [3] M.G. Brik, N.M. Avram, J. Phys.: Condens. Matter 21



- (2009) 155502.
[4] M.G. Brik, N.M. Avram, C.-G. Ma, Comput. Mater. Sci. 50
(2011) 2482.
[5] M.G. Brik, C.N. Avram, J. Lumin. 131 (2011) 2642.
[6] M.G. Brik, Physica B 532 (2018) 178.

Keywords: first-principles calculations; crystal field splitting; transition metal ions; Stokes shift

ABS-43 **Internet of Things (IOT) with Cloud Computing and Machine to Machine (M2M) communication in Aquaponics Farming Systems**

Prahenusa Wahyu Cipadi, R.Hafid Hardyanto

Universitas PGRI Yogyakarta

The IoT is gaining increasing attention. The overall aim is to interconnect the physical with the digital world. Therefore, the physical world is measured by sensors and translated into processible data, and data has to be translated into commands to be executed by actuators. Three important factors propelling the IoT forward are sensing nodes, embedded processing, and communication. This transformation is accompanied with the emergence of cloud computing capabilities supported by an increase in storage capacity, high end processing, and Machine to Machine (M2M) communication for data transport with security. With cloud computing technology, data will be stored and accessed anywhere needed with a support of powerful computing performance, efficient storage infrastructure for heterogeneous systems and software which configures and controls these different devices. The Machine to Machine (M2M) communication technology is needed, because the massive explosion of online services, further inspired by the smartphone and handheld revolution, which made these services highly accessible, has created a demand to leverage technology. In this paper, a brief introduction aquaponics farming system is given and the IoT with cloud storage and M2M communication is explained. This will also benefit the IoT in its ease of use in day-to-day life and believe to be future in this area which is getting ready for its revolution.

Keywords: Aquaponics, Internet of Things, Cloud Computing, Machine to Machine communication.

ABS-44 Potential of Ketapang Seed Oil (*Terminalia catappa* Linn) as Basic Material Mono-diglyceride Biodegradable Surfactant

Ery Fatarina Purwaningtyas (1*) ; Khoir Fadilah(2)

1*)Lecture staff in Chemical Engineering Department,
University of 17 Agustus 1945 Semarang

2) Students of the Chemical Engineering Department,
University of 17 Agustus 1945, Semarang

Surfactants are widely used in pharmaceuticals, perfumes, cosmetics, food and beverages. One type of surfactant produced from the synthesis of palm oil is mono-diglyceride which can function as an emulsifier. Ketapang (*Terminalia catappa* Linn) is a beach tree with a wide spread area, whose seeds have not been utilized optimally. The content of Ketapang seed oil has the potential to be converted into mono-diglyceride surfactant. This study uses experimental laboratory methods, while parameter optimization is done by the two level factorial design method. The raw materials used were Ketapang seeds from around the campus of Semarang 17 Agustus 1945 University, n-butanol solvents, glycerol, MgO catalysts. During the glycerolysis process, 8 run experiments were carried out with 3 variables which changed in temperature (60 & 90OC); MgO catalyst (2 & 4%) and solvent / 10 g oil volume 20 and 40 ml, while the variable is the weight of 25 gram ketapang seed oil; stirring speed of 400 rpm; reaction time of 4 hours; ratio of glycerol 2.5 ml / 10 g oil; and 24-hour deposition time. From the results of the study, the influential variable is temperature. The optimum results were obtained at conditions of 90OC, 4% MgO catalyst, solvent volume of 20 ml / 10g ketapang seed oil and yield 18.27 %. The resulting surfactant has the characteristics of acid number 57.2 mg KOH / gr, saponification number 218 mg KOH / gr.

Surfactant has HLB value 14.75, meaning that the surfactant functions as an O / W type emulsifier or as detergent agent.

Keywords: ketapang seed oil, glycerolysis, surfactant

ABS-45 BIOPLASTIC FROM JACKFRUIT SEEDS AND RICE WASTE

Retno A.S. Lestari, Mega Kasmiyatun, Kevin Dermawan,
Alfia N. Aini, Nur Riyati, Finka R. Putri

Chemical Engineering Department, University of 17
Agustus 1945 Semarang, Indonesia

This study tried to explore characterization of biodegradable plastic from jackfruit seeds and rice waste. Jackfruit seeds and rice waste contain starch, so it can be used for plastic material with addition of several additives such as polyvinyl alcohol (PVA), glycerol, chitosan and sorbitol. Starch is one of polysaccharides that be used as raw material for biofilms. Polyvinyl alcohol is the most important plastic in making water-soluble films. It was ability to form films, emulsifiers, and its adhesive properties. Polyvinyl alcohol has high tensile strength, good flexibility, and good oxygen barrier. Sorbitol and glycerol are also an additive in the manufacture of biodegradable plastics from starch, which is a plasticizer that increase elasticity. Chitosan have effects to bioplastic, such as biodegradable, hydrophilicity, and anti bacterial. This research aims to determine the effect of addition of PVA, glycerol, chitosan and sorbitol to optimum characteristics of bioplastic. The parameters to be examined bioplastic include thickness, moisture content, tensile strength, and % elongation. Increasing of glycerol added to the plastic material effect to decrease of tensile strength of bioplastic and elongation, lower water resistance and more easily degraded. Increasing of chitosan added to the plastic material effect to higher tensile strength, lower elongation, higher resistance to water and lower degradation. Increasing of PVA added to the plastic material effect to increasing of the tensile strength of the bioplastic and decreasing of elongation of bioplastic, but the increasing of

sorbitol effect to decreasing of tensile strength and increasing of elongation of bioplastic.

Keywords: biodegradable plastic, elongation, jackfruit, starch, tensile strength, waste of rice.

ABS-46 Carotenoid Dynamics Free and Bound to Pigment Protein Complexes: The Role of the 11Bu- State

Ferdy S. Rondonuwu

Department of Physics, Faculty of Science and
Mathematics, Universitas Kristen Satya Wacana, Jl.
Diponegoro 52-60, Salatiga, Indonesia

Introduction

The study focuses on the light-harvesting function of carotenoids. How carotenoids capture the photon energy and efficiently transfer its singlet energy to the neighboring bacteriochlorophyll in the pigment-protein antenna complexes (LH2) has been studied. The 11Bu- state in the light-harvesting function are critically investigated, tracing the excited-state dynamics of carotenoids free in solution and bound to the LH2.

Aim of the study

The goal of this study is to reveal the detailed mechanisms of light-harvesting function. The research focuses on the roles of the 11Bu- state in the light-harvesting function of Cars. A series of investigations were carried out to critically examine the roles of the 11Bu- state in the light-harvesting function.

Method

The internal conversion processes in Cars were investigated by the use of subpicosecond time-resolved absorption spectroscopy for a series of Cars in solution and bound to the pigment-protein antenna complexes. including neurosporene (n = 9), spheroidene (n = 10), lycopene (n = 11), anhydrorhodovibrin (n = 12) and spirilloxanthin (n = 13).



The SVD and global fitting were applied to of spectral-data matrices.

Results

The results lead to a branched relaxation scheme of energy deactivation including (1) the singlet internal conversion in the sequence of $11\text{Bu}^- \rightarrow 11\text{Bu}^- \rightarrow 21\text{Ag}^- \rightarrow 11\text{Ag}^-$ (ground), and (2) the singlet-to-triplet conversion of $11\text{Bu}^- \rightarrow 13\text{Ag}$ followed by triplet internal conversion of $13\text{Ag} \rightarrow 13\text{Bu}$. The relaxation scheme took place in subpicosecond to decapicosecond time scales.

Conclusion

The 11Bu^- state plays essential roles in light-harvesting mechanisms as a mediator state in the process of internal conversion, an acceptor state in the Car-to-BChl singlet-energy transfer and a precursor state for the rapid formation of the Car triplet state.

Keywords: Carotenoid singlet energy, relaxation scheme, SVD, Light-harvesting

ABS-47 Detecting Hoaxes in Indonesian News Using TF-IDF and K Nearest Neighbor

Eri Zuliarso, Muchamad Taufiq Anwar, Kristophorus Hadiono, Iswatun Chasanah

Universitas Stikubank, Satya Wacana Christian University

The presence of the internet and the rapid growth of social media had given rise to the blossoming of hoax creation and distribution through it. A hoax can cause anxiety and reactivity to its readers and could harm a certain party. Thereby, it is important to detect and report hoaxes to stop its spreading as soon as possible. This research aims to utilize the K Nearest Neighbor (KNN) classification algorithm to detect whether a piece of news is a hoax or not. Experiments were done by using 74 hoaxes compiled from Indonesian hoax-debunking community websites and were being compared against 74 real news from various reputable news websites in Indonesia. The result showed that the model could give detection/classification accuracy up to 83.6% and that the model is prone to false positives detections. The characteristics of the resulted model and further research directions are then discussed.

Keywords: hoax, fake news, TF-IDF, K Nearest Neighbor, KNN

ABS-48 Designing Enterprise Architecture in Koperasi Karyawan using TOGAF Architecture Development Method

Noora Qotrun Nada*, Setyoningsih Wibowo, Mega Novita

Faculty of Engineering and Informatics
Universitas PGRI Semarang
Jalan Sidodadi Timur 24 Semarang Indonesia
*noora@upgris.ac.id

The Open Group suggests that Enterprise Architecture (EA) can produce an IT blueprint that can be used to determine business, technology and information to support the achievement of the vision and mission of an organization. EA can also be seen as a model, a set of principles and methods used to produce an actual enterprise organizational structure with business processes used covering business operations, business planning, information technology and information systems. EA is also in the form of models, graphs and narratives that will explain the scope of business processes and designs of an enterprise. The purpose of this study is to design EA, especially in Koperasi Karyawan TEKAD Semarang using TOGAF framework. In other words, this architecture is created by emphasizing the relationship between IT and business. EA model is intended to maximize business alignment with IT in all situations, but depends on the nature of business and capacity in business and IT management in cooperatives.

Keywords: Enterprise Architecture, TOGAF, Koperasi

**ABS-49 DECISION SUPPORT SYSTEM MUSEUM AMBASSADOR
USING TOPSIS METHOD**

Setyoningsih Wibowo*, Maichifa Ayuning Tyas, Noora
Qotrun Nada, Mega Novita

Faculty of Engineering and Informatics Universitas PGRI
Semarang

Generally, the museum ambassador is based on various criteria from several assessment stages. The evaluation criteria for the audition stage are: interview, communication, appearance. Criteria for evaluating the quarantine stage are: creativity, ability, activity, loyalty, responsibility, and written test. Final stage assessment criteria: ideas, communication, and appearance. In the selection process, the committee uses the manual method so it is prone to errors. From these problems to the process of selecting the museum for ambassadors. The decision support system is using Topsis method. SDLC (System Development Life Cycle) is used as a design method. Research shows a score of 100 which means that the system is accepted by the user. While in testing the user test obtained from four respondents got a percentage of 89.24% which means that the system is worth of use. The system is developed in the web. So that is need more research to be developed in mobile.

Keywords: dss, museum, topsis

ABS-50 Design of Augmented Reality Game app "IntroME" using ADDIE Model

Rahmat Robi Waliyansyah (a*), Irfan Nur Ridwan (a),
Khoirul Huda (a), Febrian Murti Dewanto (a)

a) Faculty of Engineering and Informatics Universitas PGRI
Semarang

Dr. Cipto, Jl. Sidodadi Timur No.24, Karangtempel, Kec.
Semarang Tim., Kota Semarang, Jawa Tengah 50232

*rahmat.robi.waliyansyah@upgris.ac.id

An understanding of animal classification based on the type of food, is one of the sub themes in the subject of Natural Sciences. In school learning activities an understanding of the classification is conveyed through cooperative approaches such as presentations, group activities and tests. By doing the delivery method it is less attractive for students to pay attention and study in school, as well as to understand it themselves outside of teaching activities by reading books, can reduce students interest in learning it. With the existence of a different alternative in the form of an educational game, it can help in supporting the process of understanding animal classification based on the type of food carried out during teaching activities and outside of school hours. Coupled with the application of augmented reality technology on learning features that can increase user interest in learning, and for the application of learning in the form of interactive games. This educational game uses the ADDIE model and uses the design of the Unified Modeling Language (UML) which consists of Use Case Diagrams, Activity Diagrams, Squares Diagrams and as a programming language in making the application is expected to help solve existing problems. Furthermore, this application will be tested using the Black Box and User Acceptance Test. Based on the results of testing using the

User Acceptance Test obtained a percentage of 86%. Then the test results using Black Box Testing obtained a 100% valid percentage.

Keywords: Educational Game; Augmented Reality; Android; ADDIE

ABS-51 Phosphorus recovery through crystallization of struvite (MgNH₄PO₄·6H₂O): From basic research to applications

S.Muryanto

Department of Chemical Engineering,
Universitas 17 Agustus 1945 Semarang

Phosphorus (P) is the eleventh most abundant element on Earth and is essential for life, but its presence in surplus quantities is problematic. Excessive P is responsible for eutrophication of water bodies which causes deterioration of aquatic environments, naturally and economically. Also, excess of P instigates operational problems for industrial facilities due to scale formation and accumulation, which may considerably hamper heat and mass transfer and translates into substantial financial loss. In many cases, the major component of the scale is a phosphate mineral (MgNH₄PO₄·6H₂O), widely known as struvite. In contrast, due to its low solubility and its P content, struvite is an excellent fertilizer. The primary source of P is the phosphate rock, a non-renewable resource, of which about 80% is used as raw material for fertilizers. In line with increasing global population, demand for fertilizers increases rapidly, resulting in the dwindling availability of P world-wide. Thus serious attention is paid for P recovery and reuse. This paper focuses on P recovery from wastewater through crystallization of struvite. Firstly, a brief description of the theory of crystallization is presented. Then, basic research on crystallization of struvite is discussed. Finally, the different processes for P recovery as struvite are described.

Keywords: crystallization, Phosphorus, struvite

ABS-52 The Multi-agents System for Job Recommendation

Meilany Nonsi Tentua(a,b)*, Azhari Azhari (b), Aina Musdholifah(b)

(a) Teknik Informatika, Fakultas Teknik, Universitas PGRI Yogyakarta, Indonesia

(b) Department of Computer Science and Electronics, FMIPA UGM, Yogyakarta, Indonesia

Abstract. The number of available job portals causes abundant information. Therefore, a system of recommendations is needed by job seekers to find jobs that fit their profile. Offering job vacancies on job portals are changing every time because there are always additional job search data and job opening data. Multi-agents system is a technology that can be used to handle information changes.

This article proposed a recommendation system is expected to help job seekers to get jobs in accordance with the field of science they have. The system will monitor what work is offered by online job portals. From the results of observations, the job content offered will be used as a reference so that if new content is entered, the agent will automatically provide input to the job seeker. Based on the results of the implementation of the recommendation system using a multi-agent system can provide search results that are in accordance with what is inputted by the user based on the profile they have. These search results can recommend jobs that match the job seeker profile.

Keywords: multi-agents system, recommender system
Categorized in Computer Engineering

ABS-53 DESIGNING ANDROID BASED EDUCATION GAME AKSARA JAWA USING SHUFFLE RANDOM ALGORITHM

Nugroho Dwi Saputro^(a*), Tri Ana Romadhani^(b), Febrian Murti Dewanto^(b)

a) Engineering and Informatics Faculty, Universitas PGRI Semarang, Jalan Lontar No. 1, Semarang, Indonesia

*nugputra@upgris.ac.id

b) Engineering and Informatics Faculty, Universitas PGRI Semarang, Jalan Lontar No. 1 Semarang, Indonesia

The lack of appropriate supporting media in learning at the age of children makes the young generation now less familiar and still having difficulty learning Javanese script (Aksara Jawa called in Indonesia). For this reason, we need appropriate media in the introduction of Javanese script, one of which is through interesting educational games and can provide information about Javanese script. Games are forms of application that can be used as learning media where the process can be done with the concept of learning while playing for children. In this study the algorithm used was the Shuffle Random Algorithm. With this randomization algorithm, it is expected to avoid repetition of the problem so that the learning process can run effectively. Furthermore, this application was tested using the Black Box, White Box and User Acceptance Test. With the Black Box and White Box testing get a percentage of 100% while with the User Acceptance Test this application has an average of 86%, so it can be concluded that this educational game is feasible to be used as a learning medium for Javanese script.

Keywords: Javanese script, Educational Game, Randomization, Android, Shuffle Random

ABS-54 **Electronic state of sulfide-based alkali-ion conducting solid-state electrolytes applied to all-solid-state secondary batteries**

Yoshiyuki Kowada(a*), Yasukazu Akaki(a), Akiotoshi Hayashi(b), and Masahiro Tatsumisago(b)

a) Hyogo University of Teacher Education
942-1 Shimokume, Kato-shi, Hyogo 673-14, JAPAN
* ykowada@hyogo-u.ac.jp

b) Department of Applied Chemistry, Osaka Prefecture University
Gakuen-cho 1-1, Nakaku, Sakai 599-8531, JAPAN

All-solid-state batteries are expected as the next generation secondary batteries for their higher energy density, inflammable properties, and so on. In order to develop these batteries, there are several problems to improve. One of them is the lower ionic conductivities of the solid-state electrolyte. In order to improve the ionic conductivity, electronic states of the sulfide-based alkali-ion conducting solid-state electrolyte were calculated by the DV-X α cluster method. The cluster models were constructed by the coordination number reported by experimental methods and the bond length estimated from the ionic radii of each ion. The movement of the Li ion was simulated by several model clusters with different positions of the moving ion. The relationship between ionic conductivity and the differential total bond overlap population (DBOP) of the moving ion was discussed for the sulfide-based solid-state electrolytes in the several different systems. In any cases, the smaller change of DBOP of the moving cations played an important role for the fast ion movement in the solid-state electrolytes. This bonding state of the moving cations is one

of the characteristics of the electronic state in the sulfide-based solid-state electrolytes.

Keywords: all-solid-state battery, solid-state electrolyte, bonding state analysis, DV-X α method



ABS-55 Geometry and Stability of Small Gold Cluster Ions by Graph Theory and Hückel Model

Rika Sekine, Takashi Kasai, Eri Furuta, and Masaya Okakura

Department of Chemistry, Faculty of Science, Shizuoka University
836 Ohya, Shizuoka, 422-8529, Japan

Structure and stability of the gold cluster ions of which skeleton are synthesized as a complex was analyzed using the Hückel method based on graph theory. Hückel Energy (HE) and Topological Resonance Energy (TRE) were determined for neutral, positive ion, and negative ion clusters, where all the isomers of the gold cluster up to octamer were considered. Since some graphically designed isomers include bonds that can not be realized in three-dimensional space, the screening was carried out by a molecular force field calculation with LAMMPS (lammmps.sandia.gov/). Among the isomers thus obtained, both HE and TRE were most stable when the tetramer was Au₄²⁺ with a tetrahedral structure, and with the hexamer, Au₆²⁺ with two tetrahedrons sharing one side. The complexes with these structures have actually been synthesized. On the other hand, there is no reported synthetic example of the most stable cluster of octamer Au₈ with TRE and HE.

Keywords: Graph Theory, Hückel method, Gold Cluster

ABS-57 **THE EFFECT OF CHLORIDE ACID (HCL) CONCENTRATION AND FERMENTATION ON BIOETHANOL LEVELS FROM BREADFRUIT (*Artocarpus Artilis*) SUBSTRATE**

Nasrul Rofiah Hidayati
Chemical Engineering Study Program, Faculty of
Engineering, Universitas PGRI MADIUN

ABSTRACT

At present the public demand for fuel oil (BBM) is increasing inversely proportional to its availability. One of the efforts to reduce peoples dependence on BBM is to utilize alternative vegetable-based energy such as bioethanol. Bioethanol is a fermented product that can be made from a substrate containing carbohydrates. Breadfruit (*Artocarpus artilis*) is one of the agricultural products that has a fairly high starch content of 89% and is not included as a staple food source in Indonesia. Utilization of breadfruit is still not optimal because breadfruit is only used as a snack or made into flour. This study aims to make bioethanol with breadfruit raw material (*Artocarpus artilis*). The method of this study consisted of two stages, the first step was hydrolysis of breadfruit flour with a hydrochloric acid (HCL) catalyst. The concentrations of HCL used in this study were 1%, 1.5%, and 2%. The second stage of fermentation using *Saccharomyces cerevisiae* with 72 hours fermentation time, 120 hours and 168 hours. From the results of the research it was found that the highest levels of bioethanol to the treatment of HCL 2% concentration and fermentation duration of 168 hours was 17.6%.

Keywords: bioethanol, breadfruit, HCL concentration

ABS-58 Interaction Analysis Application of Arduino Industrial Automation Trainer Based on Project to Improve Cognitive Ability and Bodily Kinesthetic Ability

Ir. Sulistyaning Kartikawati, MM, M.Pd., Sekreningsih Nita, S.Kom, MT.

Universitas PGRI Madiun

S. Kartikawati^{1, 2}, S. Nita,³.

¹Program Studi Pendidikan Teknik Elektro, FKIP, Universitas PGRI Madiun

Jl. Setiabudi No. 85 Madiun

²kartikawati@unipma.ac.id

³Program Studi Teknik Informatika, FT, Universitas PGRI Madiun

Jl. Setiabudi No. 85 Madiun

Abstract: This research aims to determine the interaction/influence of the application of Arduino Industrial Automation trainer based on projects to enhance cognitive abilities and bodily kinesthetic abilities. college student This research is in the form of quantitative descriptive research. The sample used in this study is one class taken using Saturated Sampling techniques. The research class was treated by using a project-based Arduino Industrial Automation trainer. The independent variables in this study are project-based Arduino Industrial Automation trainer, while the dependent variable is cognitive abilities and bodily kinesthetic abilities.. There are 3 kinds of analysis techniques, namely instrument analysis techniques, prerequisite analysis techniques, and hypothesis analysis techniques. The measured research data are cognitive ability data and bodily kinesthetic ability data. The analysis of the hypothesis test of this study used the T-Test with a significance level of 5%. T-Test Criteria are as follows, Ho is

accepted / H1 is rejected if T count is smaller than T table and Ho is rejected / H1 is accepted if T count is greater than T table. From the results of the instrument test, namely the validation test, there are 75% questions stated valid. For the prerequisite test for the normality of cognitive abilities and bodily kinesthetic abilities on the pretest and posttest obtained L count < L table means the sample is normally distributed. For the prerequisite test for homogeneity of cognitive abilities and bodily kinesthetic abilities obtained F count < F table means the sample is homogeneous. The results of the T test for the first dependent variable cognitive ability obtained T count = 24.41 and T table = 1.697, because T count is greater than T table then Ho1 is rejected, meaning that there is an interaction/influence of the implementation of the Arduino Industrial Automation trainer based on projects in improving cognitive abilities. While the results of the T test for the dependent variable bodily kinesthetic ability obtained T count = 18.33 with T table = 1.697, because T count is greater than T table then Ho2 is rejected, it means that there is an interaction/influence of the application of the Arduino Industrial Automation trainer based on the project in increasing bodily kinesthetic abilities. So that it can be concluded that there is an interaction/influence of the application of the Arduino Industrial Automation trainer based on projects in improving cognitive abilities and bodily kinesthetic abilities.

Keywords: Industrial Automation Trainer, Arduino, Project,

ABS-59 **ANALYSIS OF THE IMPACT NARRATIVE ALGORITHM METHOD, PSEUDOCODE AND FLOWCHART TOWARDS STUDENTS UNDERSTANDING OF THE PROGRAMMING ALGORITHM COURSES**

Sekreningsih Nita, Sulistyaning Kartikawati

University PGRI Madiun

The definition of an algorithm is to describe a problem into a sequence of systematic steps in solving problems. Arranging the algorithm in sequence from beginning to end so that it can solve the problem. The results of the implementation of the algorithm are programs. So the purpose of making an algorithm is to facilitate the making of the program. There are 2 types of algorithms, 1) Text (narration and pseudocode), 2) Visual / image (flowchart). Comparison between the two, students understand it which is higher between the text or visual model. This study aims to see how much influence the value of students understands the algorithm in text or in an image. In conclusion, students are easier to understand the types of algorithms in an image. The test uses the sample T test, with a 5% significance level obtained value = 0.5. This means that the t count is greater than t table of 2.06390

Keywords: algorithm, pseudocode, flowchart, t test

ABS-60 Prediction of the Number Job Seeker Using Backpropagation Neural Network Method

Khoiriya Latifah, Agung Handayanto, Anita Rahayu

Universitas PGRI Semarang

Problem solving techniques for predicting include Artificial Neural Network. This research aims to apply the Backpropagation Neural Network method in predicting the number of job seekers in the city of Semarang. The data used in this study are secondary data, namely job seekers data of Semarang city residents from Semarang City Manpower Office from 2015-2018. In this study the data will be divided into 2 namely 2015 training data and testing data in 2018. Design and testing in this study using matlab software using 2-3-1 network architecture, which is a multilayer network with 2 input layers, 3 hidden layers and 1 output layer.

Keywords: Job Seeker, Neural Network, Backpropagation

ABS-61 **Extraction Sentiment Analysis Using naive Bayes Algorithm and Reducing Noise Word applied in Indonesian Language**

Aris Tri Jaka Harjanta, Bambang Agus Herlambang

Universitas PGRI Semarang

Sentiment Analysis is now very important and very useful in machine learning technology where a contextual mining of text to identify and extract subjective information in the source, and in helping to understand social sentiment from comments. In general, sentiment analysis can be classified into three broad categories namely sentiment positive and negative. One method of machine learning is the Deep Belief Network (DBN). DBN which is included in the Deep Learning method, is by stacking several algorithms with several extraction features that utilize all resources optimally. This research has two points. First, it aims to classify positive, negative, and neutral sentiments for the test data. The following experiments provide a system of sentiment analysis through the naive Bayes algorithm to calculate sentiment and to improve accuracy by reducing noise in words applied in Indonesian language.

Keywords: Sentiment Analysis, Machine Learning, Naive Bayes

ABS-64 **ABSTRACT****Color Based Segmentation Using Fuzzy C Means for Bamboo as Environmentally Friendly Material**

Khoiriya Latifah, Joko Siswanto, Bambang Supriyadi,
Abdul Rochim

Universitas PGRI Semarang
Universitas Islam Sultan Agung Semarang
*email: khoiriyalatifah@upgris.ac.id*¹

Bamboo is an environmentally friendly material. Many benefits of bamboo that we can get. Indonesia as a bamboo-producing country needs easy techniques to make good classification of bamboo. Bamboo is composed of fibers and fiber adhesives. There are various kinds of bamboo in Indonesia. This study uses digital image processing with fuzzy c means based segmentation to identify bamboo. Segmentation is an important thing in image processing. By using fuzzy c means in segmentation in this study obtained good segmentation results. This study uses 4 types of bamboo, namely Javanese bamboo, Ori bamboo and Petung bamboo and Wulung bamboo. There are 40 images as training images and 12 test images. The results of segmentation show that fuzzy c means produces good segmentation with the number of iterations between 20-23 and time ranging from 0.11 to 0.15. The accuracy of this test reaches 80%.

Keywords:

GALA DINNER, 2 September 2019
The 7th International Conference on DV-X α Method
(ICDM 2019)
Grand Candi Hotel, Semarang, INDONESIA

MC: Dr. Dyah Nugrahani (Universitas PGRI Semarang, Indonesia)

Time	Activities
19:00-19:10	Opening
19:10-19:20	“Indonesia Raya” Indonesia National Anthem
19:20-19:25	Report on ICDM 2019 Mega Novita, S.Si., M.Si., M.Nat.Sc., Ph.D (Chairperson of ICDM 2019, Universitas PGRI Semarang, Indonesia)
19:25-19:30	Speech Prof. Yoshiyuki Kowada (Chair of DV-X α Society, Hyogo University of Teacher Education, Japan)
19:30-19:40	Welcoming speech Dr. Muhdi, S.H., M.Hum. (Rector of Universitas PGRI Semarang)
19:40-19:50	Token appreciation for: - Keynote speakers - Invited speakers - Co-host
19:50-20:45	Dinner
20:45-21:00	Closing

GUIDELINES FOR PRESENTER(S)
2-4 September 2019
The 7th International Conference on DV-X α Method
(ICDM 2019)
Grand Candi Hotel, Semarang, INDONESIA

ORAL PRESENTER

1. ICDM 2019 distinguished the types of the speaker *i.e.*, keynote speaker, invited speaker, and general speaker
2. Keynote speaker(s) has in total 40 minutes' timeslot to talk which is 30 minutes' presentation and 10 minutes' discussion
3. Invited speaker(s) has in total 20 minutes' timeslot to talk which is 15 minutes' presentation and 5 minutes' discussion
4. General speaker(s) has in total 15 minutes' timeslot to talk which is 10 minutes' presentation and 5 minutes' discussion
5. When registering your attendance at ICDM 2019 front desk, all oral presenter(s) are required to submit their power point in PDF., PPT., or PPTX., only
6. The file name should indicate your schedule such as day (Day1), room (Room1), type of presenter (General), and abstract number, for example:

Keynote speaker	Day1_Room1_Keynote_ABS-1.pptx,
Invited speaker	Day1_Room1_Invited_ABS-1.pptx,
General speaker	Day1_Room1_General_ABS-1.pptx



7. Oral presenter(s) must submit their short paper for about 4-6 pages and follow the IOP template before **15 September 2019 (WILL NOT BE EXTENDED)**

POSTER PRESENTER

1. Poster should be prepared on A0 size (841 x 1189 mm) in a portrait format.
2. Poster presenter must prepare the 3 minutes' presentation which consists of (1) Title page, (2) Introduction, (3) Method, and (4) Result
3. When registering your attendance at ICDM 2019 front desk, all poster presenter(s) are required to submit their power point in PDF., PPT., or PPTX., only.
4. The file name should be your schedule such as day (Day1), room (Room1), type of presenter (Poster), and abstract number, for example

Poster presenter	Day1_Room1_Poster_ABS-1.pptx,
------------------	-------------------------------

5. The A0 poster will be hung on the back of Room1
6. The poster can only be put off on the Conference day 3, September 4, 2019
7. Poster presenter(s) must submit their short paper for about 4-6 pages and follow the IOP template before **15 September 2019 (WILL NOT BE EXTENDED)**


EXCURSION

CANDI BOROBUDUR TOUR

3 September 2019

The 7th International Conference on DV-X α Method (ICDM 2019)
Grand Candi Hotel, Semarang, INDONESIA

1. CANDI BOROBUDUR tour is a free tour sponsored by PT. UPGRI TOUR
2. All ICDM 2019 participants are recommended to join this activity to enjoy the wonderful Indonesia.
3. In order to get the best services from PT. UPGRI TOUR, it is very important to make reservation through this link <https://forms.gle/NhWZmoF2rmcGtD81A> before **24 August 2019**.
4. Since CANDI BOROBUDUR is a holly temple, visitor shall adhere to the dress code. Visitor may wear cool loose clothing that cover up your shoulders and knees. Wear pants as the steps leading to the top of the temple are high and not suited to skirts. Comfortable footwear is also a must.



- Postgraduate Programme
- Faculty of Education
- Faculty of Social Sciences, Health and Recreation Education
- Faculty of Mathematics, Natural Sciences, and Information Technology
- Faculty of Language and Arts Education
- Faculty of Engineering and Informatics
- Faculty of Law
- Faculty of Business and Economics



Universitas PGRI Semarang

Jl. Sidodadi Timur No. 24 - Semarang
Website: www.upgris.ac.id | Email: upgris@upgris.ac.id
Telp. (024) 8316377 | Faks. 8448217

4. Publish artikel (23 May 2020)



Meilany Nonsi Tentua <meilany@upy.ac.id>

Notification of ICDM 2019 Manuscript Publication

1 message

icdm icdm <icdm@upgris.ac.id>

Sat, May 23, 2020 at 12:59 PM

To: novita@upgris.ac.id

Bcc: meilany@upy.ac.id

Dear authors,

It is with great pleasure to inform you that the manuscripts of ICDM 2019 have been published in IOP Conference Series: Materials Science and Engineering.

For more detailed information, please click the following link:

<https://iopscience.iop.org/issue/1757-899X/835/1>

On behalf of the committee, we would like to thank you for your participation and apologize for any inconvenience, particularly for the rejected manuscripts.

Warm Regards,

ICDM 2019 Committee

Table of contents

Volume 835

2020

← Previous issue Next issue →

The 7th International Conference on DV-X α Method 2-4 September 2019, Semarang, Indonesia

Accepted papers received: 09 April 2020

Published online: 22 May 2020

[Open all abstracts](#)

Papers

OPEN ACCESS	011001
Preface	
Open abstract View article PDF	
OPEN ACCESS	011002
Peer review statement	
Open abstract View article PDF	
OPEN ACCESS	012001
Chemical forms of rhodium ion in pure water and nitric acid solution studied using ultraviolet-visible spectroscopy and first-principles calculations	
Shinta Watanabe, Toshikazu Sato, Miki Harigai, Yusuke Inaba, Kenji Takeshita and Jun Onoe	
Open abstract View article PDF	
OPEN ACCESS	012002
Supramolecular Rare Sugar	
Tomohiko Ishii, Tatsuya Senoo, Hiroaki Taguchi, Kouta Sogo, Bui Quoc Huy, Norito Taniguchi, Syuichi Katsumoto, Natsumi Nagayama and Genta Sakane	
Open abstract View article PDF	
OPEN ACCESS	012003
X-ray Absorption Spectroscopy and Its Simulation for Some Disordered Oxide Systems	
Deok-Yong Cho	
Open abstract View article PDF	
OPEN ACCESS	012004
Study on the higher-order structure and hydrogen bonding of biodegradable polymer by low-frequency vibrational spectroscopy	
Dian Marlina and Harumi Sato	
Open abstract View article PDF	
OPEN ACCESS	012005
Systematic First-Principles Calculations of Charge Transfer Transitions of Transition Metal Ions (Sc³⁺, Ti³⁺, V³⁺, Cr³⁺, Mn³⁺, Fe³⁺) in α-Al₂O₃ with Structural Optimization	
Shota Takemura, Mega Novita and Kazuyoshi Ogasawara	
Open abstract View article PDF	
OPEN ACCESS	012007
Optimization of first-principles calculation conditions for multiplet energies of Fe³⁺ and Co³⁺ in α-Al₂O₃	
Judo Fusamoto and Kazuyoshi Ogasawara	
Open abstract View article PDF	
OPEN ACCESS	012008
Water Quality in the Maintenance of <i>Oreochromis niloticus</i> through Environmentally Friendly Biofloc Technology	

JOURNAL LINKS

- [Journal home](#)
- [Journal scope](#)
- [Information for organizers](#)
- [Information for authors](#)
- [Contact us](#)
- [Reprint services from Curran Associates](#)

15:10



IOP Publishing

Progress in Physics
2024

Join us in-person or online

Register today



[Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012009

Prediction of $4f^2-4f^15d^1$ transition energy of Pr^{3+} in fluorides based on first-principles calculations and machine learning

Hayato Obata, Shota Takemura and Kazuyoshi Ogasawara

[Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012010

Lattice Relaxation Effects on the Multiplet Energies of Ruby Under Pressure using One-Electron Calculations

Mega Novita, Nur Cholifah and Kazuyoshi Ogasawara

[Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012011

Optical properties of Co^{3+} doped in $\alpha\text{-Al}_2\text{O}_3$ with Considering Lattice Relaxation Effect

Mega Novita, Duwi Nuvitalia, Nur Cholifah and Kazuyoshi Ogasawara

[Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012012

Effects of Transition Metal Doping to Cerium-Oxides in Thermal Reduction Reaction

Takaki Nishimura, Yuki Tsutsumi, Tomohiko Ishii, Tatsuya Kodama and Genta Sakane

[Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012013

Study of LCAO-MO calculation by using completely numerical basis functions

Katsumi Nakagawa

[Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012014

The magnetism change by Jahn-Teller distortion in octahedral hexa-coordinate complex

Takeyoshi Oguma, Dan Tsuneda, Yuki Tsutsumi, Yoshiaki Fujikawa, Momoe Nakano, Tomohiko Ishii, Genta Sakane and Kazuyoshi Ogasawara

[Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012015

Single-Crystal X-Ray Structural Analysis and Electronic Structures Calculation of Rare Sugar

Bui Quoc Huy, Tomohiko Ishii, Norito Taniguchi, Natsumi Nagayama and Genta Sakane

[Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012016

Data Mining Application with Fuzzy Logic Method for Monitoring Vehicle Position Based on Android

Munaf Ismail, Sri Artini Dwi Prasetyowati and Ida Widiastuti

[Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012017

Coulomb blockade transport emerged in quasi one-dimensional PEDOT: PSS fiber

Megumi Akai-Kasaya, Naoya Ogawa and Soichiro Kakinoki

[Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012018

Prediction of emission energy of Cr^{3+} in oxides based on first-principles calculations and machine learning

Tsubasa Honi and Kazuyoshi Ogasawara

[Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012019

Consideration of Superatomic Hybrid Orbital in superatom $\text{M}@\text{Al}_{12}$ ($\text{M} = \text{Be}, \text{Mg}, \text{Ca}, \text{Sr}$)

Teruyuki Toba, Masaya Yamasaki, Tomohiko Ishii and Genta Sakane

[Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012020

Mechanism of Optical Rotation of Amino Acids Using Electronic State Calculation

Shuichi Katsumoto, Manami Mimura, Tomohiko Ishii, Genta Sakane and Kazuyoshi Ogasawara

[Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012021

Generation of two-dimensional spectrochemical series in tetrahedral complexes

Takuhiro Yamamoto, Takeyoshi Oguma, Yuki Tsutsumi, Shino Tsuboi and Tomohiko Ishii

[Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012022

Effect of Molecular Distortion of Ligand Field Splitting in Five-Coordinated Metal Complex

Yoshiki Fujikawa, Dan Tsuneda, Yuki Tsutsumi, Momoe Nakano, Takeyoshi Oguma, Tomohiko Ishii, Genta Sakane and Kazuyoshi Ogasawara

[Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012023

Single Crystal X-ray Structural Analysis and Electronic Theory of Sugar Molecules

Norito Taniguchi, Tomohiko Ishii, Bui Quoc Huy and Genta Sakane

[Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012024

Cobalt Thiolates with Organic *N,N,S*-Tridentate Ligand: Conversion of Thiolate Complex to Sulfinate Complex via Sulfenate

Takanori Kotera, Atsushi Fujita, Masahiro Mikuriya, Daisuke Yoshioka, Takashi Shibahara, Kichisuke Nishimoto and Makoto Handa

[Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012025

Contribution of edge-carbon atoms and non-benzenoid rings in graphitic carbons to π^* peak profiles in CK-XANES spectra

Yasuji Muramatsu and Yuma Hirai

[Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012026

Smart Aquaponics Design Using Internet of Things Technology

R. Hafid Hardyanto and Prahonusa Wahyu Ciptadi

[Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012027

Stability of carbon-vacancy complexes in α -Fe

Masataka Mizuno, Waka Yamada, Kazuki Sugita and Hideki Araki

[Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012028

Recent applications of discrete variational multi-electron method

Kazuyoshi Ogasawara

[Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012029

Alternative Solutions for Determining Village Funds Using Weight Product Method

Tri Hastono and Firdiyan Syah

[Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012030

Design Of Electricity Energy Sources In University Of PGRI Semarang Using Off-Grid Solar Panel Systems

Adhi Kusmantoro, Mega Novita and W Th.Indriati

[Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012031

A Novel and Fast Memory Perturbation Method to Increase Exploration in Particle Swarm Optimization Algorithm

Muchamad Taufiq Anwar, Edy Winarno, Iswatun Chasanah and Saptono Nugrohadi

[Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012032

3d Ions in Solids and Microscopic Crystal Field Effects: Theoretical Analysis and Relations with Experimental Spectroscopic Data

Mikhail G. Brik

[Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012033

EKA v1: Emergency Call Auto-register, an Emergency Warning System based on Internet of Things for Intensive Care Patient at Hospital

Prisma Megantoro and Hendra Ari Winarno

[Open abstract](#) [View article](#) [PDF](#)

- OPEN ACCESS** 012034
Potential of Ketapang Seed Oil (*Terminaliacatappa Linn*) as Basic Material Mono-diglyceride Biodegradable Surfactant
Ery Fatarina Purwaringtyas and Khoir Fadilah
▼ Open abstract View article PDF
- OPEN ACCESS** 012035
Bioplastic from Jackfruit Seeds and Rice
Retno A.S. Lestari, Mega Kasmiyatun, Kevin Dermawan, Alfia N. Aini, Nur Riyati and Finka R. Putri
▼ Open abstract View article PDF
- OPEN ACCESS** 012036
Detecting Hoaxes in Indonesian News Using TF/TDM and K Nearest Neighbor
Eri Zuliarso, Muchamad Taufiq Anwar, Kristophorus Hadiono and Iswatu Chasanah
▼ Open abstract View article PDF
- OPEN ACCESS** 012037
Decision Support System Museum Ambassadors Using Topsis Method
Setyoningsih Wibowo, Maichifa Ayuning Tyas, Noora Qotrun Nada and Mega Novita
▼ Open abstract View article PDF
- OPEN ACCESS** 012038
Kinetics of Oxidation Decomposition on *Jatropha Curcas* Biodiesel
Supriyono, Sumardiyono, Enny Purwanti and S.L. Retno Ambarwati
▼ Open abstract View article PDF
- OPEN ACCESS** 012039
The Multi Agent System for Job Recommendation
Meilany Nonsi Tentua, Azhari Azhari and Aina Musdholifah
▼ Open abstract View article PDF
- OPEN ACCESS** 012040
Designing Android Based Education Game Aksara Jawa Using Shuffle Random Algorithm
Nugroho Dwi Saputro, Triana Romadhani and Febrian Murti Dewanto
▼ Open abstract View article PDF
- OPEN ACCESS** 012041
Electronic state of sulfide-based alkali-ion conducting solid-state electrolytes applied to all- solid-state secondary batteries
Yoshiyuki Kowada, Yasukazu Akaki, Akitoshi Hayashi and Masahiro Tatsumisago
▼ Open abstract View article PDF
- OPEN ACCESS** 012042
The Effect Of Chloride Acid (Hcl) Concentration And Fermentation On Bioethanol Levels From Breadfruit (*Artocarpus Artilis*) Substrate
Nasrul Rofiah Hidayati
▼ Open abstract View article PDF
- OPEN ACCESS** 012043
Interaction Analysis Application of Arduino Industrial Automation Trainer Based on Project to Improve Cognitive Ability and The Bodily-Kinesthetic Ability
S. Kartikawati and S. Nita
▼ Open abstract View article PDF
- OPEN ACCESS** 012044
Analysis Of The Impact Narrative Algorithm Method, Pseudocode And Flowchart Towards Students Understanding Of The Programming Algorithm Courses
Sekreningsih Nita and Sulistyning Kartikawati
▼ Open abstract View article PDF
- OPEN ACCESS** 012045
Color Based Segmentation Using Fuzzy C Means For Bamboo As EnvironmentallyFriendly Material
Khoirya Latifah, Joko Siswanto, Bambang Supriyadi and Abdul Rochim
▼ Open abstract View article PDF
- OPEN ACCESS** 012046



OPEN ACCESS

The Effect Of Decomposer Microorganism Additions On The Natural Hydroponic Technology (Nht) Systems Of Pakchoy Growth

Ary Susatyo Nugroho, Endah Rita Susatya Dewi and Muhammad Anas Dzakiy

Open abstract View article PDF

OPEN ACCESS

Damarjati Application for Development of Javanese Language Learning in High Schools

Suyitno, Ngatmini and Setyoningsih Wibowo

Open abstract View article PDF

OPEN ACCESS

Design of Product Monitoring System Using Internet of Things Technology for Smart Manufacturing

Marti Widya Sari, Herianto, IGB Budi Dharma and A.E. Tontowi

Open abstract View article PDF

OPEN ACCESS

Designing Enterprise Architecture in Koperasi Karyawan using TOGAF Architecture Development

Noora Qotrun Nada, Setyoningsih Wibowo and Mega Novita

Open abstract View article PDF

OPEN ACCESS

Geometry and Stability of Small Gold Cluster Ions by Graph Theory and Hückel model

Rika Sekine, Takashi Kasai, Eri Furuta and Masaya Okakura

Open abstract View article PDF

OPEN ACCESS

Extraction Sentiment Analysis Using naive Bayes Algorithm and Reducing Noise Word applied in Indonesian Language

Aris Tri Jaka Harjanta and Bambang Agus Herlambang

Open abstract View article PDF

OPEN ACCESS

Temperature and Humidity Observation System in ATmega8 Microcontroller-Based Homes

Abdul Latif and Kuart Supriyadi

Open abstract View article PDF

OPEN ACCESS

Design and Development of Heart Rate Per Minutes Based on Atmega16 Microcontroller with Alarm Warning

Nur Hudha Wijaya, Achmad Kholilul Mujib, Agung Budi Santoso and Kuart Supriyadi

Open abstract View article PDF

OPEN ACCESS

Decision Support System in Recommending Climbing Tourism Destinations with Profile Matching Method

Ni Luh Wiwik Sri Rehayu Ginantra, Gita Widi Bhawika, Ahmad Zamsuri, Fadjar Budianto and GS Achmad Daengs

Open abstract View article PDF

OPEN ACCESS

Conjugate Gradient Polak Ribiere In Improving Performance in Predicting Population Backpropagation

Medi Herman Tinambunan, Erna Budhiarti Nababan and Benny Benyamin Nasution

Open abstract View article PDF

OPEN ACCESS

Effect of chitosan-zinc oxide coated in PVDF membrane; morphology and performance testing

A Febriasari, D Siswanta, N Riyanto, N Hidayat Aprilita and F Silvianti

Open abstract View article PDF

OPEN ACCESS

Text Mining Pre-Processing Using Gata Framework and RapidMiner for Indonesian Sentiment Analysis

S Kurniawan, W Gata, D A Puspitawati, I K S Parthama, H Setiawan and S Hartini

Open abstract View article PDF

OPEN ACCESS

Application of Data Mining Technique using K-Medoids in the case of Export of Crude Petroleum Materials to the Destination Country

Fathur Rahman, Ihda Innar Ridho, M. Muflih, Sefto Pratama, Mokhammad Ramdhani Raharjo and Agus Perdana Windarto



Application of MCDM using PROMETHEE II Technique in the Case of Social Media Selection for Online Businesses.

I Gede Iwan Sudipa, Cici Astria, Khairunnissa Fanny Imanda, Agus Perdana Windarto, Nelly Khairani Daulay, Wildan Suharso and Harna Oktafia Lingga Wijaya



IOPSCIENCE

Journals

Books

IOP Conference Series

About IOPscience

Contact Us

Developing countries access

IOP Publishing open access policy

Accessibility

IOP PUBLISHING

Copyright 2024 IOP Publishing

Terms and Conditions

Disclaimer

Privacy and Cookie Policy

PUBLISHING SUPPORT

Authors

Reviewers

Conference Organisers



This site uses cookies. By continuing to use this site you agree to our use of cookies.



IOP Publishing

PAPER • **OPEN ACCESS**

The Multi Agent System for Job Recommendation

To cite this article: Meilany Nonsi Tentua *et al* 2020 *IOP Conf. Ser.: Mater. Sci. Eng.* **835** 012039

View the [article online](#) for updates and enhancements.

You may also like

- [Machine Learning Techniques for True and Fake Job Posting](#)
Kamakshi Mehta, Navaneetha Krishnan Rajagopal, Sagar Balu Gaikwad *et al.*
- [Research on Position Recommendation System Based on Convolutional Neural Network](#)
Yuan Li, Guangjian Ning and Haoxi Zhang
- [Design and achievement of hardware-in-the-loop simulation system for strapdown semi-active laser seeker](#)
Wenbin Zhou, Chunxi Zhang, Siyuan Wang *et al.*



ECS The Electrochemical Society
Advancing solid state & electrochemical science & technology

ECS UNITED

247th ECS Meeting
Montréal, Canada
May 18-22, 2025
Palais des Congrès de Montréal

Showcase your science!

**Abstracts due
December
6th**

The Multi Agent System for Job Recommendation

Meilany Nonsi Tentua^{*1,2}, Azhari Azhari², Aina Musdholifah²

¹Teknik Informatika, Fakultas Teknik, Universitas PGRI Yogyakarta, Indonesia

²Department of Computer Science and Electronics, FMIPA UGM, Yogyakarta, Indonesia

meilany@upy.ac.id, arisn@ugm.ac.id, aina_m@ugm.ac.id

Abstract. The number of available job portals causes abundant information. Therefore, a system of recommendations is needed by job seekers to find jobs that fit their profile. Offering job vacancies on job portals are changing every time because there are always additional job search data and job opening data. Multi-agents system is a technology that can be used to handle information changes. This article proposed a recommendation system is expected to help job seekers to get jobs in accordance with the field of science they have. The system will monitor what work is offered by online job portals. From the results of observations, the job content offered will be used as a reference so that if new content is entered, the agent will automatically provide input to the job seeker. Based on the results of the implementation of the recommendation system using a multi-agent system can provide search results that are in accordance with what is inputted by the user based on the profile they have. These search results can recommend jobs that match the job seeker profile.

1. Introduction

Today, there are many portals that offer online job openings (Job Portal) in Indonesia, such as Indeed.com, Jobindo.com, karir.com, loker.id and komaskarir.com. Job portal provides job search features based on position, company name, job location and salary amount. The online job portal application allows job seekers and recruiters to be directly connected [1].

The number of available job portals causes abundant information. This often causes job seekers to get confused and eventually take jobs that are not in accordance with their profile. Therefore, a system of recommendations is needed by job seekers to find jobs that fit their profile. The recommendation system also helps people to find interesting items by utilizing previous user interactions [2]. In addition, the recommendation system has a successful use in e-commerce applications to handle problems related to information overload efficiently [3].

Offering job vacancies on job portals are changing every time because there are always additional job search data and job opening data. So that for job recommendation systems must contain technology that can adapt to changes in the environment in the system. In addition, the recommendation system must also be able to gather information from several existing job portals. Multi-agents system is a technology that can be used to handle information changes. An agent is a computer system, which is in a certain environment, which is capable of performing autonomous actions that are flexible, in order to meet the design goals [4]. Multi-agent system has been widely applied in several applications such as job search agencies[5], health care [6], biomedical [7], and so on. In this paper, a system of



recommendations for finding a job using a multi-agent system will be proposed. This job recommendation system platform uses agent technology, its JADE.

2. Literature Review

2.1. Recommendation System

The recommendation system is an information filtering system that handles the problem of information overload [8] [8] by filtering fragments of information from a large number of data dynamically information generated according to user preferences, interests, or observed behavior about items [9] [9].

Several methods have been applied by previous researchers about recommendation systems such as profile matching [10] [11], collaborative filtering [12] [13] [14], clustering [15] , decision tree [16], [16], classification [17], machine learning [18], content base [19], hybrid [20]

Basically, the methods in the recommendation system can be divided into three categories: content-based, collaborative filtering and hybrid models. The recommendation system based on content uses items that interact with users in the past, then calculates similarity scores between pairs of items and ranks new recommendations accordingly [21]. The collaborative filtering approach suggests items to certain users, who interact with other users who have positive interactions [12]. Hybrid method is a method which combines both techniques [22].

2.2. Multi-Agent System

Multi-agents system is a technology that can be used to handle information changes. An agent is a computer system, which is in a certain environment, which is capable of performing autonomous actions that are flexible, in order to meet the design goals [4].

Several characteristics of multi agent systems can be used for the purpose of this study

- Autonomy, that is, agents can carry out independent and indirect tasks by users, other agents or by the environment.
- Intelligence, Reasoning, and Learning: Every agent must have a minimum standard to be called an agent, namely intelligence (intelligence). In the concept of intelligence, there are three components that must be possessed: an internal knowledge base, the ability to reason based on the knowledge base that is owned, and the ability to learn to complete in a changing environment.
- Mobility and Stationery: specifically for mobile agents, he must have the ability to represent the characteristics he has is mobility. In contrast to stationary agents. However, it still must have the ability to send messages and communicate with other agents.
- Delegation: In accordance with the appearance and we have discussed in the resolution section, the agent moves in a move task that is ordered by the user. This phenomenon of delegation (delegation) is the main characteristic of a program called an agent.
- Reactivity: Another characteristic agent is the ability to be able to quickly with the help of changing information in the environment (enviornment). Environments that can include: other agents, users, outside information, etc.
- Proactive and Target Oriented: Proactive nature is arguably a continuation of reactivity. Agents are not only required to change the environment, but also must take step by step what to take. For this reason, the agent must be designed to have a clear goal (goal), and to always be oriented towards the goals it is carrying out (goal oriented).
- Communication and Coordination Ability: Agents have the ability to communicate with users and other agents. To be able to coordinate with other agents in carrying out tasks, standard languages are needed to communicate, such as Knowledge Queries and Language Manipulation (KQML) and Knowledge Interchange Format (KIF).

The use of multi-agents on the recommendation system has been done a lot before. A multi-agent is used in the TV recommendation system based on the user's desire to select thousands of TV programs broadcasted by hundreds of TV stations [23] [24]. News recommendations that use a multi-agent framework by utilizing Social Media Preferences [25].

3. Methodology

Based on the explanation above to overcome the problem, the researcher proposes a multi-agent system for job recommendations that are in accordance with the field of job seekers. An agent is a computer system that is in an environment and has the ability to act autonomously in the environmental situation in accordance with the objectives designed [4].

The system is used to recommend work in accordance with the field of science that has been taken by job seekers. The system will monitor what work is offered by online job portals. From the results of observations, the job content offered will be used as a reference so that if new content is entered, the agent will automatically provide input to the job seeker. This proposed system is expected to help job seekers to get jobs in accordance with the field of science they have.




Figure 1 . Agent Work Flow Chart

Figure 1. Shows agent workflows in this job recommendation system. On the basis of this system the agent is grouped into two criteria based on its function, namely:

- information gathering agent
This agent serves to collect information from various job portals currently available, such as indeed.com, jobsdb.com, karir.com, etc.
- agent user
This agent serves to search for job openings that have been collected in the database based on user profiles

Any changes that occur in the environment will also result in changes to the agent's actions. The environment used is an online job portal on the internet. If there is a change in the portal, it will give an impact on the system agent. This environment will be modeled into a database that will be updated based on updates that occur in online portal jobs.

4. Implementation

The implementation of the job recommendation system based on the design consists of two interfaces, an interface for the user and an interface for displaying search results.

4.1. a. User interface

The user will enter the name, status, and profile. After that when the user clicks ok, the agent will search for the appropriate profile in the database.

Figure 2. User interface

4.2. b. Interface results

The results of the search request performed by the user will be displayed on the results interface. The results displayed are the name of the appropriate job and the name of the company offering the job.

```

Nama : PT Surya Akapindo Semarang +13 lokasi
Pekerjaan : Staff Administrasi Pembukuan

Nama : SG Yogyakarta Yogyakarta
Pekerjaan : Admin

Nama : PT. Central Prima Kelola - Central Park Mall Jakarta
Pekerjaan : Admin Engineering / Data Entry

Nama : PT Indoreksa Lokamandiri Jakarta
Pekerjaan : Service Engineer Admin

Nama : PT. Asuransi Sinar Mas Yogyakarta Yogyakarta
Pekerjaan : Admin Klerikal

Nama : PT Sumber Daya Bergengsi Indonesia +20 lokasi
Pekerjaan : Staff Administrasi

```

Figure 3. Interface Results

5. Discussion

The online job portal application allows job seekers and recruiters to be directly connected. The number of available job portals causes abundant information. This often causes job seekers to get confused and eventually take jobs that are not by their profile. Utilization of the multi-agent system in the job recommendation system created can gather all job information offered by all online job portals. The same work from several portals will be sorted and only displayed once. So the user will not be offered repeatedly. In the system that has been created, data is collected from several online job portals such as, indeed.com, jobsdb.com, karir.com. Job data collection on the online job portal is done by the agent and then collected in one table. The results of data collection job can be seen in Figure 4.

id	name_job	company
130	ADMIN OFFICER	Angkasa Pura Supports Semarang
131	administrasi pembayaran	PT PELANGI FORTUNA GLOBAL Salatiga
132	Admin	SG Yogyakarta Yogyakarta
133	Recruitment Officer (RO)	PT. Ponseljobs Indonesia Jakarta
134	Staf Analis	SG Yogyakarta Yogyakarta
135	MAGANG DATA ENTRY	PT PELAYARAN TEMPURAN EMAS 4 ulasan Surabaya
136	Staff Administrasi Pembukuan	PT Surya Akapindo Semarang +13 lokasi
137	Cleaning Service OB Office Boy Asisten Masak	PT. Djalala Trading International Jakarta
138	Data Entry Team	PT Surya Akapindo Jakarta +13 lokasi
139	Salesman/Salesgirl Otomotif (Surabaya dan Sekitarnya)	PT Mobilkamu Group Indonesia Sidoarjo
140	Sales Executive (Mojokerto)	PT Mobilkamu Group Indonesia Mojokerto
141	Marketing Otomotif	PT Mobilkamu Group Indonesia Gresik
142	Marketing Mobil Baru All Brand	PT Mobilkamu Group Indonesia Mojokerto

Figure 4. Collecting data

Each online job portal has a different presentation in providing information, even from the same company. So that in this system each online job portal has one agent to retrieve information. Retrieval of information tailored to the needs of the system, so that it can be put together in the table. By utilizing the characteristics of the autonomous agent, the agent is assigned to retrieve job information every time there is a change in the online job portal

On the user side, the system will provide information about what work is by user requests. An agent is assigned to search for information based on the suitability of the user in the collection of work information that has been collected in the table. Furthermore, the system will provide information to the user automatically or job recommendations based on changes that occur in the online job portal.

6. Conclusion

The online job portal application allows job seekers and recruiters to be directly connected. The number of available job portals causes abundant information. This often causes job seekers to get confused and eventually take jobs that are not by their profile. Utilization of the multi-agent system in the job recommendation system created can gather all job information offered by all online job portals. The same work from several portals will be sorted and only displayed once. So the user will not be offered repeatedly. Based on the results of the implementation of the recommendation system using a multi-agent system can provide search results that are in accordance with what is inputted by the user based on the profile they have. These search results can recommend jobs that match the job seeker profile.

References

- [1] S. Arunkumar, S. Gautam, and A. Behera, "Multifunctional Online Job Portal," *Journal of Network Communications and Emerging Technologies (JNCET)* vol. 8, no. 4, p. 4, 2018.
- [2] K. C. Kara, S. Esen, N. Kahyalar, A. A. Karakaş, and T. Aytekin, "Design and implemenatation of a job recommender system," in *2017 International Conference on Computer Science and Engineering (UBMK)*, 2017, pp. 729–733.
- [3] S. T. Al-Otaibi, "A survey of job recommender systems," *Int. J. Phys. Sci.*, vol. 7, no. 29, Jul. 2012.
- [4] Wooldridge, "An Introduction to Multi Agent Systems." [Online]. Available: <https://www.pdfdrive.com/wooldridge-an-introduction-to-multi-agent-systems-d5107234.html>. [Accessed: 21-May-2019].
- [5] X. Qiu, N. Zhou, and X. Wang, "Design of Labour Agency Platform Based on Agent Technology of JADE," in *Computer and Computing Technologies in Agriculture V*, vol. 368, D. Li and Y. Chen, Eds. Berlin, Heidelberg: Springer Berlin Heidelberg, 2012, pp. 41–45.
- [6] R. Morejon Novales, "A Multi-Agent Approach To Data Mining Processes: Applications To Health Care," *Mestre Em Informática*, Pontifícia Universidade Católica Do Rio De Janeiro, Rio de Janeiro, Brazil, 2018.
- [7] M. Camara, O. Bonham-Carter, and J. Jumadinova, "A multi-agent system with reinforcement learning agents for biomedical text mining," in *Proceedings of the 6th ACM Conference on Bioinformatics, Computational Biology and Health Informatics - BCB '15*, Atlanta, Georgia, 2015, pp. 634–643.
- [8] J. A. Konstan and J. Riedl, "Recommender systems: from algorithms to user experience," *User Model. User-Adapt. Interact.*, vol. 22, no. 1, pp. 101–123, Apr. 2012.
- [9] F. O. Isinkaye, Y. O. Folajimi, and B. A. Ojokoh, "Recommendation systems: Principles, methods and evaluation," *Egypt. Inform. J.*, vol. 16, no. 3, pp. 261–273, Nov. 2015.
- [10] A. Gupta and D. Garg, "Applying data mining techniques in job recommender system for considering candidate job preferences," in *2014 International Conference on Advances in Computing, Communications and Informatics (ICACCI)*, 2014, pp. 1458–1465.

- [11] V. Desai, D. Bahl, S. Vibhandik, and I. Fatma, "Implementation of an Automated Job Recommendation System Based on Candidate Profiles," *Int. Res. J. Eng. Technol. IRJET*, vol. 04, no. 05, p. 4, May 2017.
- [12] C. Fry, "A Comparison of Collaborative Filtering Algorithms for Job Recommendation Using Apache Mahout," *Proj. Present. Fac. Calif. State Polytech. Univ. Pomona*, p. 101, 2016.
- [13] W. Shalaby *et al.*, "Help Me Find a Job: A Graph-based Approach for Job Recommendation at Scale," *ArXiv180100377 Cs*, Dec. 2017.
- [14] R. G. Belsare and D. V. M. Deshmukh, "Employment Recommendation System using Matching, Collaborative Filtering and Content Based Recommendation," *Int. J. Comput. Appl. Technol. Res.*, vol. 7, no. 6, pp. 215–220, Jun. 2018.
- [15] I. Editor, "Survey on Recommendation System using Data Mining and Clustering Techniques."
- [16] A. Utku, H. Karacan, O. Yildiz, and M. A. Akcayol, "Implementation of a New Recommendation System Based on Decision Tree Using Implicit Relevance Feedback," *JSW*, vol. 10, pp. 1367–1374, 2015.
- [17] G. Özcan and Ş. G. Ögüdücü, "Applying Classifications Techniques in Job Recommendation System for Matching of Candidates and Advertisements," *Int. J. Intell. Comput. Res.*, vol. 8, no. 1, p. 9, 2017.
- [18] M. Nilashi, "An Overview of Data Mining Techniques in Recommender Systems," *J. Soft Comput. Decis. Support Syst.*, vol. 3, no. 6, pp. 16–44, Oct. 2016.
- [19] A. Elsafty, M. Riedl, and C. Biemann, "Document-based Recommender System for Job Postings using Dense Representations," in *Proceedings of the 2018 Conference of the North American Chapter of the Association for Computational Linguistics: Human Language Technologies, Volume 3 (Industry Papers)*, New Orleans - Louisiana, 2018, pp. 216–224.
- [20] J. Séguéla and G. Saporta, "A hybrid recommender system to predict online job offer performance," p. 15.
- [21] N. D. Almalis, G. A. Tsihrintzis, and N. Karagiannis, "A content based approach for recommending personnel for job positions," in *IISA 2014, The 5th International Conference on Information, Intelligence, Systems and Applications*, 2014, pp. 45–49.
- [22] R. Burke, "Hybrid Web Recommender Systems," in *The Adaptive Web: Methods and Strategies of Web Personalization*, P. Brusilovsky, A. Kobsa, and W. Nejdl, Eds. Berlin, Heidelberg: Springer Berlin Heidelberg, 2007, pp. 377–408.
- [23] K. Kurapati, S. Gutta, D. Schaffer, J. Martino, and J. Zimmerman, "A Multi-Agent TV Recommender," Adaptive Systems Department, Philips Research Briarcliff 345 Scarborough Rd. Briarcliff Manor, NY 10510, USA p. 8.
- [24] Y. Blanco-Fernandez, J. J. Pazos-Arias, A. Gil-Solla, M. Ramos-Cabrer, B. Barragans-Martinez, and M. Lopez-Nores, "A Multi-Agent Open Architecture for a TV Recommender System: A Case Study Using a Bayesian Strategy," in *IEEE Sixth International Symposium on Multimedia Software Engineering*, Miami, FL, USA, 2004, pp. 178–185.
- [25] M. A. Ghalib Ahmed Tahir, "Personalized News Recommendation based on Multi-agent framework using Social Media Preferences - IEEE Conference Publication." [Online]. Available: <https://ieeexplore.ieee.org/document/8538403>. [Accessed: 24-May-2019].