

KORESPONDENSI
WITH: APPLIED ECOLOGY AND ENVIRONMENTAL RESEARCH (Q3)

Title:
Response of Rice Growth and Yield under Different Timing of NPK-Based Fertilizer Application in the Salibu System

Web SJR Journal:

<https://www.scopus.com/sourceid/100147015#tabs=2>

The screenshot shows the Scopus Preview page for the journal "Applied Ecology and Environmental Research". The page includes the following information:

- Source details:**
 - Scopus coverage years: 2003, from 2005 to Present
 - Publisher: Corvinus University of Budapest
 - ISSN: 1589-1623 E-ISSN: 1785-0037
 - Subject area: (Agricultural and Biological Sciences: Agronomy and Crop Science) (Agricultural and Biological Sciences: Ecology, Evolution, Behavior and Systematics)
 - Source type: Journal
- Metrics:**
 - CiteScore 2022: 1.9
 - SJR 2022: 0.233
 - SNIP 2022: 0.353
- CiteScore rank & trend:**
 - CiteScore 2022: 1.9 (4,178 Citations 2019 - 2022, 2,198 Documents 2019 - 2022)
 - CiteScoreTracker 2023: 1.4 (2,171 Citations to date, 1,513 Documents to date)

Web H-Index Journal:


<https://www.scimagojr.com/journalsearch.php?q=100147015&tip=sid&clean=0>

The screenshot shows the Scimago Journal & Country Rank page for the journal "Applied Ecology and Environmental Research". The page includes the following information:

- COUNTRY:** Hungary
- SUBJECT AREA AND CATEGORY:** Agricultural and Biological Sciences, Agronomy and Crop Science, Ecology, Evolution, Behavior and Systematics
- PUBLISHER:** Corvinus University of Budapest
- H-INDEX:** 38

Web Journal:

<http://www.aloki.hu/>



Manuscript Submission: **1 November 2024**

a. Cover Letter

COVER LETTER

To Applied Ecology and Environmental Research

Dear Editor,

I would like to send an original article entitled: **Response of Rice Growth and Yield under Different Timing of NPK-Based Fertilizer Application in the Salibu System** for Applied Ecology and Environmental Research to consider. I confirm that this work is genuine and has not been published elsewhere, nor is it considered for publication elsewhere. We believe and hope that this manuscript is worthy of publication by Applied Ecology and Environmental Research. We are interested in publishing articles in this journal because it has an excellent reputation, so it is a matter of pride if published in Applied Ecology and Environmental Research. Here I attach the manuscript.

Thank you
Best regards,

Paiman
Universitas PGRI Yogyakarta, Indonesia

b. Manuscript for Submission

RESPONSE OF RICE GROWTH AND YIELD UNDER DIFFERENT TIMING OF NPK-BASED FERTILIZER APPLICATION IN THE SALIBU SYSTEM

PAIMAN^{1*} – RAKASIWI, N¹, DELZA¹ – MUADIM, N¹ – SRIWATI, E¹, WIBOWO, W.T¹ –LAILI, L. N¹ – MUSTAKIM¹ – KUSUMASTUTI, C. T¹ – ARDIYANTA¹

¹Agrotechnology Department, Agriculture Faculty, Universitas PGRI Yogyakarta, Yogyakarta 55182, Indonesia

(e-mail/phone: paiman@upy.ac.id/ +62-821-3439-1616; riskiraka46@gmail.com/ +62-821-8035-8795; delzawess@gmail.com/ +62-813-9970-6506; nurmuadim@gmail.com/ +62-822-9880-0503; endangsriwati949@gmail.com/ +62-812-3751-0996; wahyutribowow281201@gmail.com/ +62-858-7715-2262; LatifahLaili8@gmail.com/ +62-882-1645-4912; mustakimagro@gmail.com/ +62-812-4685-7797; kusumastuti@upy.ac.id/ +62-821-3439-1664; ir.ardiyanta@upy.ac.id/ +62-812-275-7813)

**Corresponding author*

e-mail: paiman@upy.ac.id ; phone: +62-821-3439-1616; fax: +62-274-376-808

Abstract: In the Salibu system (SS), rice plants have a shorter growth cycle than their parent plants. Applying NPK as a based fertilizer can effectively stimulate early shoot growth, giving the plants optimal vegetative growth and maximum yield. This study aimed to determine the optimal timing of NPK-based fertilizer application to enhance rice growth and yield in the SS. The experiment was arranged in a randomized complete block design (RCBD) with three replications, where each replication acted as a block. The timing of NPK-based fertilizer application included four treatments: no application (control), at the time of harvesting, at the time of stem cuttings (seven days after harvest, or 7 DAH), and at seven days after stem cuttings (7 DAC). The results indicated that the application of NPK-based fertilizer can enhance the dry weight of shoots clump⁻¹ (g), grains clump⁻¹ (g), and the weight of 1000-filled grains (g) in the SS. The application of NPK-based fertilizer at the time of stem cuttings in the parent plants (7 DAH) resulted in a higher dry weight of milled grains clump⁻¹ (15.3 g) compared to the control (13.2 g). The research findings indicate that applying NPK-based fertilizer at the time of stem cuttings of the parent plants is the best time to enhance the rice growth and yield in the SS. Future research should recommend determining the optimal dosage of NPK-based fertilizer to maximize rice growth and yield.

Keywords: *based fertilizer, NPK, rice, stem cuttings, Salibu system*

INTRODUCTION

As Indonesia's population grows, the demand for food, particularly rice, is increasing. Various measures have been implemented to achieve national rice self-sufficiency, including adapting the SS. The abbreviation of Salibu is derived from the Indonesian phrase “Salin ibu”, which translates to “change of the parent plants” in English. Salibu is modified from ratoon rice. However, farmers have not widely accepted this method due to its relatively low production. Farmers prefer methods that provide higher and more stable yields, even with smaller profit margins. The SS features a shorter life cycle than parent plants, enabling increased productivity by applying NPK-based fertilizer. Furthermore, using NPK-based fertilizer can accelerate the growth of rice shoots during the early stages, thereby enhancing the potential for higher yields.

The SS is a rice cultivation method based on local wisdom, originally implemented in West Sumatra, Indonesia. The technology represents a modification of the ratoon rice cultivation technique and has been tested in various regions across Indonesia. Although farmers have not widely adopted it due to perceptions of lower yields, this technology can enhance rice harvests and improve land-use efficiency with proper management. According to Abdurachman et al. (2015),

the SS offers advantages such as faster harvest times, more efficient water usage, and reduced production costs by up to 45% compared to the transplanting method. This method reduces land preparation, seedling production, and planting expenses. Therefore, the SS has the potential to improve the welfare of farmers.

The success of the SS technology is influenced by various factors, one of which is the timing of NPK-based fertilizer application. The NPK fertilizer plays a crucial role in supporting the growth of the new shoots after the parent plants are cut. Applying the fertilizer can provide the essential nutrients the new shoots need, promoting their development. Since the SS has a shorter harvest cycle than its parent plants, it is important to determine the optimal fertilizer application timing to accelerate the growth of the new shoots. However, research on the timing of NPK-based fertilizer in the SS remains limited. Therefore, further studies are necessary to identify the optimal timing for fertilizer application in the SS.

In Indonesia, rice cultivation in irrigated land typically occurs one to three times annually, depending on the irrigation system employed. In rainfed fields, rice can be planted only once a year, while fully irrigated fields can support up to three planting cycles each year. Each planting season involves various activities, including land preparation, seedling production, planting, plant maintenance, and harvesting. Conventional rice cultivation often incurs relatively high costs, lowering profit margins. In contrast, the SS offers greater advantages due to its land and resource utilization efficiency.

The SS can enhance rice production and land productivity annually. This method utilizes the shoots that emerge from the basal stems after cuttings. To maximize the number of tillers, apply the optimal dosage of NPK-based fertilizer. New shoots emerge from the nodes of the remaining stem, followed by the emergence of new roots. Eventually, these shoots grow into independent plants. The formed shoots subsequently develop into new tillers, similar to the rice planting system used in transplanting techniques (Juanda, 2016). The SS can enhance food availability in Indonesia and contribute to food security and resilience. It is more effectively implemented on land that is not always flooded (Jahari & Sinaga, 2019). Rice productivity in tropical regions primarily relies on fertilizers, as soils often lack nitrogen and phosphorus (Hindersah et al., 2022).

Nitrogen (N), phosphorus (P), and potassium (K) are applied in large quantities as fertilizers in rice fields, and a deficiency of any of these nutrients can reduce crop yields (Shrestha et al., 2020). Nitrogen plays a crucial role in the synthesis of chlorophyll and protein. Phosphorus is involved in energy storage and transfer, and it is a key element in nucleic acids, coenzymes, nucleotides, phosphoproteins, phospholipids, and phosphate sugars. Meanwhile, potassium supports starch formation, activates enzymes, and acts as a catalyst in storing photosynthetic products.

Nitrogen is an essential element that forms plants' amino acids, proteins (enzymes), nucleic acids, and chlorophyll. A nitrogen deficiency in rice cultivation leads to reduced plant height, lower tiller formation, decreased leaf area index, and diminished photosynthesis rates (Chauhan et al., 2017). Nitrogen plays a crucial role in promoting growth, enhancing yields, and improving grain quality by improving the number of tillers, expanding leaf area, and facilitating protein synthesis. Plants absorb nitrogen through nitrate ions (NO_3^-) and ammonium ions (NH_4^+) (Kaya, 2013). A lack of nitrogen can limit rice productivity, while excessive nitrogen can result in agronomic and economic losses (Hussain et al., 2023). The yield increased when nitrogen was applied at $202.15 \text{ kg ha}^{-1}$. The fertilization frequency was four times (He et al., 2024). Optimal nitrogen fertilization can effectively compensate for the negative impacts of N fertilizer,

enhancing grain quality and nitrogen use efficiency (NUE) while sustaining high yields (Tian-Yang et al., 2022).

Phosphorus (P) is essential in photosynthesis, respiration, energy transfer, storage, cell division, and enlargement. Plants absorb phosphorus in the form of primary orthophosphate ions, H_2PO_4^- (Irwanto, 2014). Additionally, phosphorus is a key component of high-energy compounds such as adenosine triphosphate (ATP) and synthesizes phospholipids, nucleotides, and glycoproteins. Phosphorus deficiency can hinder growth and yield in rice, which is evident through reduced tiller numbers and dry matter accumulation, along with symptoms such as stunted growth and upright leaves that exhibit a dark green colour (Chauhan et al., 2017).

Potassium (K) activates various enzymes in plant metabolic processes and is absorbed as potassium ions (K^+) (Irwanto, 2014). This nutrient is essential for enhancing root growth, promoting plant vigour, and increasing resistance to pests and diseases. Symptoms of potassium deficiency in rice include stunted growth with fewer tillers, dark green upper leaves, and chlorosis in the areas between leaf veins (Chauhan et al., 2017). Although chemical fertilizers can improve rice yield, the response highly depends on optimal fertilization management in paddy fields (Yang et al., 2021). Using a combination of $135\text{--}210\text{ kg ha}^{-1}\text{ N}$ and $115\text{--}137\text{ kg ha}^{-1}\text{ K}$ application to achieve relatively higher yields and improve quality in rice production (Chen et al., 2024).

The NPK Phonska (15-15-15) fertilizer is complex inorganic NPK with composition of 15% N, 15% P_2O_5 , 15% K_2O , and 10% S. This fertilizer is widely utilized by farmers due to its effectiveness in enhancing rice yields and grain quality. According to Shrestha et al. (2020), nitrogen (N), phosphorus (P), and potassium (K) are applied as fertilizers in large quantities to rice fields. Research conducted by Kurniadie (2002), using a dosage of $300\text{ kg ha}^{-1}\text{ NPK Phonska}$ along with $333\text{ kg ha}^{-1}\text{ ZA}$ resulted in the highest performance for the IR 64 rice variety, including parameters such as the number of panicle clump $^{-1}$, the number of grain panicle $^{-1}$, the harvest index, the weight of 1000 grains, as well as dry grain yields (7.50 tons ha^{-1}) and milled rice yields (6.17 tons ha^{-1}).

Optimized fertilization practices represent a promising management approach for achieving sustainable rice production (Zhuang et al., 2022). The application of NPK Phonska fertilizer at a rate of 300 kg ha^{-1} , combined with $200\text{ kg ha}^{-1}\text{ urea}$, significantly increased plant height, the number of tillers, and yield up to $7.448\text{ tons ha}^{-1}$, with relative agronomic effectiveness (RAE) value of 106.46 (Fidiyawati et al., 2022). Combining NPK fertilizer with urea is essential to meet nitrogen nutrient requirements during various rice growth. Without urea addition, higher dosages of NPK were needed; specifically, $300\text{ kg ha}^{-1}\text{ NPK Phonska (15-15-15)}$ required an additional $150\text{ kg ha}^{-1}\text{ urea}$ (Hartatik & Widowati, 2015). The greatest yield of 5.81 tons ha^{-1} was achieved with the combined treatment of $250\text{ kg ha}^{-1}\text{ NPK} + 150\text{ kg ha}^{-1}\text{ urea} + 1,000\text{ kg ha}^{-1}\text{ dolomite}$ with a yield of 5.81 tons ha^{-1} (Ratmini et al., 2023).

Applying NPK fertilizer can enhance plant height and yield by 3%, although it may reduce the weight of 1000 grains. On the other hand, nitrogen fertilizer significantly contributed to an increase in productive tillers and boosted yield by 6%. The most effective combination for rice fields with high phosphorus levels was $300\text{ kg ha}^{-1}\text{ NPK}$ and $100\text{ kg ha}^{-1}\text{ urea}$ (Siska & Ismon, 2019). The NPK application also enhances the absorption of nutrients N, P, and K, and improves grain yield (Shrestha et al., 2020). Additionally, Srinivas (2017) noted a significant increase in the number of tillers and panicles m^{-2} with the application of $120\text{--}60\text{--}60\text{ kg ha}^{-1}\text{ NPK}$.

The application of NPK fertilizer significantly influenced plant height at 35, 45, and 90 days after planting (DAP), the number of tillers at 35 and 45 DAP, and various production parameters

such as the number of panicles clump⁻¹ and the weight of 1000-filled grains of rice (Waty et al., 2014). However, a dosage of 50 kg ha⁻¹ NPK Phonska did not show significant differences in rice production across various agroecological zones (highlands, midlands, and lowlands) using the SS (Fitri et al., 2019). In the SS, fertilization adhered to the parent plants: the first application of 40% recommended dosage at 15-20 DAP and the second consisting of 60% at 30-35 DAP (Abdulrachman et al., 2015).

Based on the background and literature review, NPK fertilizer is more commonly used under the transplanting system in rice cultivation. In contrast, its application as base fertilizer in the SS has not been thoroughly investigated. Therefore, the study aimed to determine the optimal timing of NPK-based fertilizer application to enhance rice growth and yield in the SS.

MATERIALS AND METHODS

Study Site

The research was conducted from January to June 2024 in Tirtonirmolo Village, Kasihan District, Bantul Regency, Special Region of Yogyakarta, Indonesia. The site was located at an elevation of 70 m above sea level, with an average air temperature of 26.6 °C and humidity of 83%. Kasihan District is 110°20'40" East Longitude and 7°48'42" South Latitude.

Experiment Design

The experiment was arranged in an RCBD with three replications, where each replication acted as a block. The timing of NPK-based fertilizer application included four treatments: no application (as a control), application at the time of harvesting the parent plants, application at the time of cutting the parent plant stems or seven days after harvest (DAH), and application at seven days after cutting the stems (DAC). This study required 12 treatment plots, each measuring 2 × 2 m² (or 4 m²) and comprising 64 plant stems. The research flow can be seen in *Figure 1*.




Figure 1. The flow diagram in research activity

Research Procedures

Parent plants

In the SS, the parent plants cultivated by farmers were utilized. The variety employed was Inpari 42, which has a harvesting period of 87 DAP. The spacing was set at 23×23 cm. The land used for cultivation was technically irrigated, ensuring a consistent water supply, and consisted of alluvial soil.

Salibu system

After harvesting the parent plants, the stems at seven days after harvest (7 DAH) were cut to a height of 3 cm from the soil surface. The new shoots emerged from the remaining cut of parent stems and were carefully maintained. Plant care included irrigation, fertilization, and the management of pests, diseases, and weeds to ensure optimal growth.

The application of NPK-based fertilizer occurred according to the treatment schedule, which took place at the time of harvesting, at the time of stem cuttings (7 DAH), and seven days after stem cuttings (7 DAC). The dosage of NPK-based fertilizer used was 60 g plot^{-1} , equivalent to 150 kg ha^{-1} NPK Phonska, except in the control plots. After applying NPK-base fertilizer, a subsequent fertilizer was recommended with 150 kg ha^{-1} NPK Phonska and 200 kg ha^{-1} urea. It was given in two stages: the first stage (40% of the recommended dosage) was applied at 14 DAC, followed by the second stage (60% of the recommended dosage) at 28 DAC.

Weeding was performed at 12 and 26 DAC to reduce weed competition. Pest and disease management was implemented according to integrated pest management (IPM) principles at 56 and 63 DAC, using Decis as the control agent. Rice harvesting occurred at 75 DAC when the rice reached physiological maturity (95% yellowing). With the appropriate maintenance and fertilization practices, the rice yield in the SS is expected to be maximized.

Parameters

The research observations included the growth and yield components from each plot treatment. The plant growth components comprised the percentage of tillering stems (%), plant height (cm), the number of tillers clump⁻¹ (stems), and dry weight of shoots (leaves and stems) clump⁻¹ (g). In addition, the observed yield components included the dry weight of grains clump⁻¹ (g), the weight of 1000-filled grains (g), and panicle length (cm).

Statistical Analysis

The observation data were analyzed using analysis of variance (ANOVA) at 5% significant level (Gomez & Gomez, 1984) with IBM SPSS Version of 23 software. If differences were found between treatment means, further testing was conducted using Duncan's new multiple range test (DMRT) at 5% significant level.

RESULTS

Component of the Rice Growth

Tillering Stems

Observations of the tillering stems were conducted at 14 DAC. The results of the ANOVA (*Appendix 1*) indicates that the timing of NPK-based fertilizer application did not significantly affect the percentage of tillering stems that sprouted. The number of tillering stems is presented in *Table 1*.

Table 1. The number of tillering stems (%) at 14 DAC

The timing of NPK-based fertilizer application			
No application (control)	At the time of harvesting	At the time of stem cuttings	At seven days after stem cuttings
92.7 a	93.8 a	97.9 a	97.4 a

Remarks: The values in the same row with different letters are significantly different based on DMRT at 5% significant level.

Figure 2a shows shoot emergence from the basal stem of the parent at 7 DAC, while *Figure 2b* illustrates the tiller growth at 28 DAC.



a. The shoot emergence from the basal stem of the parents at 7 DAC



b. The tiller growth in the SS at 28 DAC

Figure 2. Photo of the shoot emergence from the basal stem of the parents at 7 DAC (a) and the tiller growth at 28 DAC (b)

Plant height

Plant height was observed at regular intervals of 14, 28, 42, and 56 DAC. The results of the ANOVA (*Appendix 2*) find that the timing of NPK-based fertilizer application significantly affected the plant height at 28 and 42 DAC, while there was no significance at 14 and 56 DAC. The average plant height is presented in *Table 2*.

Table 2. The plant height at 14, 28, 42, and 56 DAC

Time of observations (DAC)	The timing of NPK-based fertilizer application			
	No application (control)	At the time of harvesting	At the time of stem cuttings	At seven days after stem cuttings
14	33.9 a	32.3 a	42.1 a	36.0 a
28	62.4 b	70.3 ab	73.3 a	63.3 b
42	87.0 a	86.9 a	84.3 ab	80.8 b
56	90.1 a	91.3 a	89.5 a	86.8 a

Remarks: The values in the same row with different letters are significantly different based on DMRT at 5% significant level.

Table 2 shows that applying NPK-based fertilizer increased plant height at the 28 DAC. The application of NPK-based fertilizer at the time of cutting the stem of the parent plants resulted in taller plants compared to applying it at 7 DAC. Additionally, the treatment of NPK-based fertilizer application at the time of cutting showed a significant difference compared to applying it at the time of harvesting. However, at the 42 DAC observation, it appeared that the application of NPK-based fertilizer led to a decrease in plant height. At the same time at the 56 DAC, there were no significant differences among the treatments.

Tiller numbers

The number of tillers was also observed at regular intervals of 14, 28, 42, and 56 DAC. The results of the ANOVA (*Appendix 3*) shows that the timing of NPK-based fertilizer application had no significant effect on the number of tillers clump⁻¹ at 14, 28, 42, and 56 DAC. The average number of tillers is presented in Table 3.

Table 3. The number of tiller clump⁻¹ at 14, 28, 42, and 56 DAC

Time of observations (DAC)	The timing of NPK-based fertilizer application			
	No application (control)	At the time of harvesting	At the time of stem cuttings	At seven days after stem cuttings
14	7.3 a	6.4 a	7.9 a	6.0 a
28	14.6 a	15.9 a	16.9 a	13.6 a
42	17.5 a	18.2 a	19.0 a	15.2 a
56	16.2 a	15.9 a	16.2 a	13.8 a

Remarks: The values in the same row with different letters are significantly different based on DMRT at 5% significant level.

Component of the Rice yield

The dry weight of shoots clump⁻¹ (g), grains clump⁻¹ (g), weight of 1000-filled grain (g), and panicle length (cm) were observed at 75 DAC. The results of the ANOVA (*Appendix 4*) indicates that the timing of NPK-based fertilizer application significantly affected the dry weight of shoots, grains clump⁻¹, and weight 1000 grains, while panicle length showed no significant effect. The dry weight of shoots, grains, weight of 1000-filled grains, and panicle length are presented in Table 4.

Table 4. The means of shoot dry weight, grain dry weight, the weight of 1000-filled grain, and panicle length at 75 DAC.

Parameters observed	The timing of NPK-based fertilizer application			
	No application (control)	At the time of harvesting	At the time of stem cuttings	At seven days after stem cuttings
Shoots dry weight clump ⁻¹ (g)	23.1 b	31.3 a	32.2 a	30.1 a
Grains dry weight clump ⁻¹ (g)	13.2 b	13.9 ab	15.3 a	14.6 ab

Weight of 1000-filled grains (g)	13.3 b	19.0 a	19.3 a	17.3 a
Panicle length (cm)	18.9 a	18.6 a	20.5 a	19.8 a

Remarks: The values in the same row with different letters are significantly different based on DMRT at 5% significant level.

Table 4 shows that the application of NPK-based fertilizer at harvest, at the time of stem cuttings of the parent plants, and at 7 DAC can significantly increase the shoot dry weight clump⁻¹ (g), grain dry weight clump⁻¹ (g), and weight of 1000-filled grains (g).

DISCUSSION

Components of Rice Growth

The application of NPK-based fertilizer in the SS did not influence the percentage of new shoots emerging from cut stems and tillers number. This may be because the new shoots primarily rely on the food reserve stored in the parent stems. After cutting, the emerging shoots utilize the resources available in the remaining stem for growth. Therefore, although NPK provides essential nutrients, internal factors such as the energy stored in the parent stem may be more dominant in determining the success of new shoot emergence. While NPK contains essential nutrients, the growth of tillers is influenced by various factors, including the genetic traits of the rice variety, soil conditions, and management techniques. In the SS, which emphasizes effective resource use, the number of tillers may be more influenced by care practices and other cultivation techniques rather than by fertilizer alone. Thus, applying NPK as a fertilizer was insufficient to increase the number of tillers significantly.

In contrast, the application of NPK-based fertilizer significantly impacted plant height in the SS. NPK provides nitrogen, phosphorus, and potassium, which are crucial for vegetative growth. Nitrogen plays a role in protein synthesis and leaf growth, directly affecting plant height. Additionally, phosphorus supports better root development, allowing the plant to absorb more nutrients from the soil. With optimal nutrient availability, the plant can grow taller, demonstrating that while NPK did not affect new shoots and tillers numbers, it was key in supporting plant height growth. According to Samira et al. (2012), the application of NPK can increase the plant height of rice.

Components of Rice Yield

In the early stages of growth, new shoots utilize the food reserves from the remains of the parent stems while forming new roots that replace the function of the parent plant's roots. New shoots were ready to utilize nutrients available in their environment 10 days after the stem cuttings. The application of NPK-based fertilizer at the time of cutting the parent stems enhanced the availability of essential nutrients, including nitrogen (N), phosphorus (P), and potassium (K), which were crucial for rice growth in the SS. The new shoot's roots quickly absorbed available nutrients to support their development. This base fertilizer helps provide the essential nutrients needed by the new shoots that emerge from the stem cuttings. New shoots will appear the day after cutting, although small and without leaves.

The application of NPK-based fertilizer significantly increased shoot dry weight compared to the control. This fertilizer contains essential nutrients such as nitrogen, phosphorus, and potassium, crucial for plant growth. Nitrogen plays a vital role in chlorophyll formation and leaf development. With the application of NPK, plants can develop denser leaves and wider shoots, enhancing their photosynthetic capacity. As a result, the shoots' dry weight was higher than those that did not receive the fertilizer. Yang et al. (2004) emphasized that the rice root system was vital in absorbing water and nutrients from the soil. According to Wang et al. (2006), root system formation was closely coordinated with shoot growth, where the shoot regulated the root development through the circulation of materials between the two. Root distribution determines the structure of the root system within the soil, affecting the acquisition of soil resources such as nutrients and water.

In addition to increasing shoots dry weight, it positively influenced grain dry weight clump⁻¹ higher than the control. The nutrients provided by NPK support the grain-filling process, enabling the plants to produce more and heavier grains. Applying nitrogen and phosphorus is crucial for optimal grain development during the grain-filling phase. Plants receiving NPK-based fertilizer show a significant increase in the dry weight of grains clump⁻¹, indicating that a good nutrient supply contributes to better harvest yields. The product of photosynthesis in the leaves can be translocated in the seed-filling process if the plants are supplied with this nutrient. According to Rawat et al. (2022), phosphorus is the most abundant element after nitrogen and is involved in many physiological processes such as photosynthesis and translocation. Sarkar et al. (2021) stated that the capacity for source-sink dynamics and assimilate translocation can play an important role during grain formation. As explained by Wei et al. (2018), rice grain yield depends on the supply assimilates (source) and the grain capacity to utilize them (sink).

The application of NPK resulted in the resulting grains being heavier and of higher quality than the control. Adequate nutrients aid in grain-filling, allowing the grains to fill out properly and achieve greater dry weight. The application of NPK-based fertilizer not only increased the quantity of the harvest but also improved the quality of the rice grains, which is essential for determining the market value of the produce. According to Metwally et al. (2020), the grain quality characteristics can be enhanced by the NPK fertilizer application. Explain by Pasaribu et al. (2018), the parent plant's ability to generate new shoots highly depends on the carbohydrate content and phytohormones retained in the intercalary meristem tissues of the stubble after harvest. Additionally, increasing the rice yield with the SS can be achieved by enhancing fertilization, especially nitrogen.

Therefore, applying NPK-based fertilizer at the time of stem cutting was more effective in increasing shoot dry weight, grain dry weight, and the weight of 1000-filled grains than using it at harvest or 7 DAC. The application of NPK-based fertilizer at the time of stem cutting provided essential nutrients for the new shoots that emerged from the remaining parts of the parent plants. The fertilizer applied at this time can be directly utilized by the shoots after ten days, as nitrogen, phosphorus, and potassium are crucial for accelerating the formation of new tissues and photosynthesis. In contrast, applying fertilizer at harvest or 7 DAC was less effective because the growth of new shoots had not yet started during that period. The availability of nutrients was not optimally utilized, leading to lower dry weight and harvest yields. Thus, the timing of fertilizer-based application significantly influenced nutrient absorption efficiency and the final yield of the plants.

Conclusions

The results indicated that the application of NPK-based fertilizer can enhance the plant height (cm), dry weight of shoots clump⁻¹ (g), grains clump⁻¹ (g), and weight of 1000-filled grains (g) in the SS. The application of NPK-based fertilizer at the time of stem cuttings in the parent plants (7 DAH) resulted in a higher dry weight of milled grains clump⁻¹ (15.3 g) compared to the control (13.2 g). The research findings indicate that applying NPK-based fertilizer at the time of stem cuttings of the parent plants is the best time to enhance rice growth and yield in the SS. Future research should recommend determining the optimal dosage of NPK-based fertilizer to maximize rice yields.

Acknowledgements. We sincerely thank the Institute for Research and Community Service at Universitas PGRI Yogyakarta for funding and supporting this research. We also wish to thank the Faculty of Agriculture, Universitas PGRI Yogyakarta, for providing the necessary financial support for greenhouse facilities and equipment used in this study.

REFERENCES

- [1] Abdulrachman, S., Suhartatik, E., Erdiman, Susilawati, Zaini, Z., Jamil, A., Mejaya, M. J., Sasmita, P., Abdulah, B., Suwarna, Baliadi, Y., Dhalimi, A., Sujinah, Suharna, Ningrum, E. S. (2015): Panduan teknologi budidaya padi salibu. Badan Penelitian dan Pengembangan Pertanian, Kementerian Pertanian. (in Indonesian)
- [2] Chauhan, B. S., Jabran, K., Mahajan, G. (2017): Fertilizer management in rice. – In: Rice Production Worldwide (Issue February 2017, pp. 1–563).
- [3] Chen, G., Duan, Q., Wu, C., He, X., Hu, M., Li, C., Ouyang, Y., Peng, L., Yang, H., Zhang, Q., Jiang, Q., Lan, Y., Li, T. (2024): Optimizing rice yield, quality, and nutrient use efficiency through combined application of nitrogen and potassium. – *Frontiers in Plant Science* 15: 1–16.
- [4] Fidiyawati, E., Latifah, E., Krismawati, A., Sihombing, D., Setyorini, D., Bakar, A., Sugiono. (2022): Effectivity of inorganic fertilizer NPK (15-15-6) to growth and yield of lowland rice (*Oryza sativa* L.) on alfisol soil. – In: IOP Conference Series: Earth and Environmental Science. The 4th International Conference on Food and Agriculture 980: 012014.
- [5] Fitri, R., Erdiman, Kusnadi, N., Yamaoka, K. (2019): Salibu technology in Indonesia: An alternative for efficient use of agricultural resources to achieve sustainable food security. – *Paddy and Water Environment* 17: 403–410.
- [6] Gomez, A. G., & Gomez, K. A. (1984): Statistical procedures for agricultural research (Second edition). John Wiley & Sons, Inc. New York.
- [7] Hartatik, W., Widowati, L. R. (2015): Effect of compound fertilizer NPKS and NPK on rice growth and yield on Inceptisol. – *Jurnal Penelitian Pertanian Tanaman Pangan* 34(3): 175–186. (in Indonesian)
- [8] He, X., Zhu, H., Shi, A., Wang, X. (2024): Optimizing nitrogen fertilizer management enhances rice yield, dry matter, and nitrogen use efficiency. – *Agronomy* 14: 919.
- [9] Hindersah, R., Kalay, A. M., Talahaturuson, A. (2022): Rice yield grown in different fertilizer combination and planting methods: Case study in Buru Island, Indonesia. – *Open Agriculture* 7: 871–881.
- [10] Hussain, T., Mulla, D. J., Hussain, N., Qin, R., Tahir, M., Liu, K., Harrison, M. T., Sinutok, S., Duangpan, S. (2023): Optimizing nitrogen fertilization to enhance productivity and profitability of upland rice using CSM–CERES–Rice. – *Plants* 12: 3685.
- [11] Irwanto. (2014): Pengaruh pemberian pupuk NPK terhadap pertumbuhan tanaman buah naga di Kecamatan Pemayung, Kabupaten Batanghari, Provinsi Jambi. Balai Pelatihan Pertanian Jambi. (in Indonesian)

Indonesian)

- [12] Jahari, M., Sinaga, P. H. (2019): Overcome limited growing season with cultivation ratoon and salibu. – Jurnal Dinamika Pertanian 3: 65–72.
- [13] Juanda, R. (2016): Peningkatan produksi padi melalui potensi dan pengembangan wilayah produksi benih unggul di Provinsi Aceh. – Agrosamudra 3(2): 72–80. (in Indonesian)
- [14] Kaya, E. (2013): Effect of rice straw compost and fertilizer NPK on N-available of soil, N-uptake, growth and yield of paddy (*Oryza sativa* L.). – In: Prosiding FMIPA Universitas Pattimura 2013 (pp. 41–47). (in Indonesian)
- [15] Kurniadie, D. (2002): Pengaruh kombinasi dosis pupuk majemuk NPK phonska dan pupuk N terhadap pertumbuhan dan hasil tanaman padi sawah (*Oryza sativa* L.) varietas IR 64. – Jurnal Bionatura 4(3): 137–147. (in Indonesian)
- [16] Metwally, T. F., Mohamed, A. A. E., Sorour, S. G. R., Elsayed, G. A. (2020): Stimulating compounds affect the grain quality characteristics and the nutritional value of rice (*Oryza sativa*). – Applied Ecology and Environmental Research 18(5): 6829–6840.
- [17] Pasaribu, P. O., Triadiati, Anas, I. (2018): Rice ratooning using the salibu system and the system of rice intensification method influenced by physiological traits. – Pertanika 41(2): 637–654.
- [18] Ratmini, N. P. S., Subiksa, I. G. M., Suprihatin, A., Suastika, I. W., Riyanto, D. (2023): Optimization of lowland productivity through fertilization as a food buffer against the impact of climate change. – E3S Web of Conferences, 9th ICCO 2023, 467: 01019.
- [19] Rawat, J., Pandey, N., Saxena, J. (2022): Role of potassium in plant photosynthesis, transport, growth, and yield. – In: Role of Potassium in Abiotic Stress. Springer Nature Singapore Ltd. (pp. 1–14).
- [20] Samira, D., Gani, A., Slavich, P., McLeod, M. (2012): Effect of NPK fertilizer and biochar residue on paddy growth and yield of second planting. – In: 2nd Annual International Conference Syiah Kuala University 2012 & The 8th IMT-GT Uninet Biosciences Conference, Banda Aceh, 22-24 November 2012, 2(1): 22–24.
- [21] Sarkar, S., Monshi, F. I., Uddin, M. R., Tabassum, R., Sarkar, M., Hasan, A. (2021): Source-sink manipulation influences the grain-filling characteristics associated with the grain weight of rice. – Journal of Innovative Agriculture 8: 20–29.
- [22] Shrestha, J., Shah, K. K., Timsina, K. P. (2020): Effects of different fertilizers on growth and productivity of rice (*Oryza sativa* L.): a review. – International Journal of Global Science Research 7(1): 1291–1301.
- [23] Siska, W., Ison, L. (2019): Pemupukan NPK dan nitrogen pada tanaman padi di lahan sawah berstatus P tinggi di Sumatera Barat. – Jurnal Pengkajian dan Pengembangan Teknologi Pertanian 22(2): 175–184. (in Indonesian)
- [24] Srinivas, M. (2017): Response of rice (*Oryza sativa* L.) to different nutrient levels under machine planted conditions. – In: 5th International Conference on Agriculture, Ecology and Biological Engineering, Thailand, May 2-4, 2017 (pp: 1–5).
- [25] Tian-Yang, Z., Zhi-Kang, L., En-Peng, L., Wei-Lu, W., Li-Min, Y., Hao, Z., Li-Jun, L., Zhi-Qin, W., Jun-Fei, G., Jian-Chang, Y. (2022): Optimization of nitrogen fertilization improves rice quality by affecting the structure and physicochemical properties of starch at high yield levels. – Journal of Integrative Agriculture 21(6): 1576–1592.
- [26] Wang, H., Inukai, Y., Yamauchi, A. (2006). Root development and nutrient uptake. – Critical Reviews in Plant Sciences 25(3): 279–301.
- [27] Waty, R., Musyassir, Syamaun, Chairunnas. (2014): Effects of NPK fertilization and biochar residual

to physical properties of soil, plant growth and result of rice field (*Oryza sativa* L.) second planting season. – Jurnal Manajemen Sumber Daya Lahan 3(1): 383–389.

- [28] Wei, H., Meng, L., Li, X., Dai, Q., Zhang, H., Yin, X. (2018): Sink-source relationship during rice grain filling is associated with grain nitrogen concentration. – Field Crops Research 2015: 23–38.
- [29] Yang, C., Yang, L., Yang, Y., Ouyang, Z. (2004): Rice root growth and nutrient uptake as influenced by organic manure in continuously and alternately flooded paddy soils. – Agricultural Water Management 70: 67–81.
- [30] Yang, G., Nabi, F., Sajid, S., Kaleri, A. R., Jakhar, A. M., Cheng, L., Raspor, M., Muhammad, N., Ma, J., Hu, Y. (2021): Response of root development and nutrient uptake of two Chinese cultivars of hybrid rice to nitrogen and phosphorus fertilization in Sichuan Province, China. – Molecular Biology Reports 48(12): 8009–8021
- [31] Zhuang, Y., Ruan, S., Zhang, L., Chen, J., Li, S., Wen, W., Liu, H. (2022): Effects and potential of optimized fertilization practices for rice production in China. – Agronomy 42(32): 1–16.

APPENDIX

Appendix 1. ANOVA of tillering stem

Source of variance	Degree of freedom	Sum of Squares	Mean square	F calc.	F table 5%
Block	2	37.028	18.514	2.703 ns	5.14
Treatment	3	60.832	20.277	2.960 ns	4.76
Error	6	41.097	6.850		
Total	11	138.957			

Remarks: ns = not significant at 5%

Appendix 2. ANOVA of plant height

Source of variance	Degree of freedom	Mean square				F table 5%
		14 DAC	28 DAC	42 DAC	56 DAC	
Block	2	9.027 ns	6.754 ns	5.127 ns	4.868 ns	5.14
Treatment	3	55.394 ns	84.675 *	25.636 *	10.640 ns	4.76
Error	6	20.947	18.678	4.934	13.549	
Total	11					

Remarks: ns = not significant at 5% and * = significance different at 5%

Appendix 3. ANOVA of tiller number

Source of variance	Degree of freedom	Mean square				F table 5%
		14 DAC	28 DAC	42 DAC	56 DAC	
Block	2	9.896 ns	17.453 ns	20.818 ns	12.766 ns	5.14
Treatment	3	2.200 ns	6.444 ns	8.132 ns	4.125 ns	4.76
Error	6	7.861	4.960	10.220	14.141	
Total	11					

Remarks: ns = not significant at 5%


Appendix 4. ANOVA of shoot dry weight, grain dry weight, weight of 1000-filled grain, and panicle length

Source of variance	Degree of freedom	Mean square				F table 5%
		Shoot dry weight	Grain dry weight	Weight of 1000-filled grain	Panicle length	
Block	2	13.898 ns	0.259 ns	0.250 ns	0.002 ns	5.14
Treatment	3	51.490 *	2.570 *	22.750 *	2.207 ns	4.76
Error	6	7.719	0.494	3.250	2.829	
Total	11					


Remarks: ns = not significant at 5% and * = significance different at 5%

Manuscript Submitted: 1 November 2024


URL: <https://mail.google.com/mail/u/0/#sent>




URL: <https://mail.google.com/mail/u/0/#sent/KtbxLwGzcRLSsswVGVDQHxPILsnRLRpwwg>




Pemberitahuan biaya publikasi dari jurnal: 6 November 2024






Approval to pay publication fees: 11 November 2024




Reviewing Process: 12 November 2024

<https://mail.google.com/mail/u/0/#inbox/FMfcgzQXKDbzTCznWBgmTKnZNBgdhVSC>



This screenshot shows a Gmail inbox with a list of 54 emails. The top email is from 'Applied Ecology and Environmental Research' with the subject 'Submitted manuscript: Applied Ecology and Environmental Research, ref. 16969 - Thank you for your informa...'. The inbox also shows other emails from 'eContent Pro', 'EcoSummit 2024', 'LinkedIn', 'Routledge Books', 'Google Scholar Aler.', 'Kudos: Science need.', 'ResearchGate', and 'Levi's® Red Tab™'.



This screenshot shows a Gmail email thread. The top email is from 'Applied Ecology' with the subject 'Submitted manuscript: Applied Ecology and Environmental Research, ref. 16969'. The email body contains the following text:

Dear Authors,

We would like to inform you that the full reviewing process of MS ref. 16969 (titled: *RESPONSE OF RICE GROWTH AND YIELD UNDER DIFFERENT TIMING OF NPK-BASED FERTILIZER APPLICATION IN THE SALIBU SYSTEM*; by authors: PAIMAN – RAKASIWI, N. – DELZA – MUADIM, N. – SRIWATI, E. – WIBOWO, W.T –LAILI, L. N – MUSTAKIM – KUSUMASTUTI, C. T – ARDIYANTA,) is in progress.

If you did not co-author this submission or disagree with the process for any reason, please contact us by e-mail as soon as possible!

Thank you in advance!

Best regards,

Technical Editors
Applied Ecology and Environmental Research
Open Access International Scientific Journal
<http://www.aloki.hu/>

The bottom email is from 'Dr. Ir. Paiman, M.P. - UPY' with the subject 'Submitted manuscript: Applied Ecology and Environmental Research, ref. 16969'. The email body contains the following text:

Thank you for your information.

Extract of referee opinions for MS 16969: 6 Januari 2025

This screenshot shows a Gmail inbox with a list of emails. The email titled "Extract of referee opinions for MS 16969 - Received, thank you. On Mon, Jan 6, 2025 at 2:00 PM Applied Ecol..." is selected. The left sidebar shows the "Inbox" with 429 emails. The top navigation bar includes "Compose", "Search mail", and "Active" status. The email list includes various newsletters and academic updates.

From	Subject	Date
Applied, me 2	Extract of referee opinions for MS 16969 - Received, thank you. On Mon, Jan 6, 2025 at 2:00 PM Applied Ecol...	Jan 6
BPP UMY	Newsletter UMY November 2024 Issue - Dear Paiman Paiman Hope you have an amazing day! With great ple...	Jan 6
SEA-Teacher.. me 3	The 13th SEAMEO-University of Tsukuba Symposium - Forwarded message ----- From: COLL...	Jan 6
Google Scholar Aler.	1 new citation to your articles - Pengaruh herbisida bahan aktif glifosfat dan parakuat terhadap gulma dan h...	Jan 6
Google Scholar Aler.	Recommended articles - [HTML] The Impact of Split Nitrogen Fertilizer Applications on the Productivity and ...	Jan 6
LinkedIn	Dr. Ir. Paiman., follow Vivek Ramaswamy - Founder at Vivek Ramaswamy - See your recommendations ...	Jan 6
LinkedIn	You have 1 new invitation - You have 1 new invitation	Jan 5
Kudos: Science need.	Claim your latest articles - Start growing your research impact	Jan 5
Google Scholar Aler.	Recommended articles - Exogenously applied proline with silicon and zinc nanoparticles to mitigate salt stre...	Jan 5
LinkedIn	You have 1 new invitation - You have 1 new invitation	Jan 4
ResearchGate	Paiman, a recent article cited your research - Paiman, a recent article cited your research Paiman, we found ...	Jan 4
Pubtext to Editor111	The effect of soil water content and biochar on rice cultivation in polybag - Journal of Soil Science and Plant ...	Jan 4
Google Scholar Aler.	Recommended articles - Residual effects of biochar and nano-modified biochar on growth and physiology u...	Jan 4
Howly	Download Your File Presentation3.pdf - Download Your File: Make the Most of Howly Docs! Hello! A quick rem...	Jan 4

This screenshot shows the content of the email titled "Extract of referee opinions for MS 16969". The email is from "Applied Ecology" to "me" on Monday, Jan 6, 2:00 PM (2 days ago). The content includes a greeting, a message about the manuscript review, and two attachments: "Extract from Re..." and "AEER_16969.docx". The email is also labeled as "External" and "Inbox".

Extract of referee opinions for MS 16969 External Inbox x

Applied Ecology
to me
Mon, Jan 6, 2:00 PM (2 days ago)

Dear Author,

We are sending you enclosed an extract of referee opinions for your manuscript ref. 16969.
Please consider the suggestions and resubmit the final ready-to-publish version within approximately 2 weeks if possible.

Thank you in advance!

Best regards,

Technical Editors
Applied Ecology and Environmental Research
Open Access International Scientific Journal
<http://www.aloki.hu/>

2 Attachments • Scanned by Gmail

Extract from Re... AEER_16969.docx

Dr. Ir. Paiman, M.P. - UPY <paiman@upy.ac.id>
to Applied
Mon, Jan 6, 10:31 PM (2 days ago)

Received, thank you.

Extract of referee opinions and editorial suggestions for manuscript ref. 16969

Authors aimed to test the effect of NPK fertilisation on a specific rice growth system to enhance productivity and profitability. The experimental design needs further clarification, the size of an experimental plot and the number of measured plants per plot has to be indicated. The conclusion has to be in harmony with the displayed results if the optimal timing is based on the yield performance.

Comments to the Authors:

1. Sentences between L25-27 and L32-35 contain the same information. Please revise.
 2. It is suggested to introduce the ratoon technique and afterwards, authors might highlight what is the modification compared to that in salibu rice cultivation.
 3. Please add a map of the study site and the environmental conditions might be given, too.
 4. It is not completely clear what is the differences between harvest and stem cutting, what is the usual difference between those (in days).
 5. Please add the schedule and amount of the irrigation.
 6. Please indicate the row length and row number of an experimental plot.
 7. The rice variety might be added.
 8. Please clarify how many plants were measured/sampled when measuring the variables.
 9. Please add $\pm$ SD values to the tables.
 10. Authors stated that at 42 DAC the NPK fertilizer led to a decrease in plant height, but this statement is only valid in the case of 7 days after stem cutting application.
 11. In the Discussion, the authors stated that the application of NPK-based fertilizer significantly impacted plant height in the SS but this statement is not valid for all measurements and each timing. Please be more specific.
 12. The authors concluded the following: The application of NPK-based fertilizer at the time of stem cuttings in the parent plants (7 DAH) resulted in a higher dry weight of milled grains clump-1 (15.3 g) compared to the control (13.2 g). The research findings indicate that applying NPK-based fertilizer at the time of stem cuttings of the parent plants is the best time to enhance rice growth and yield in the SS. On the other hand in L239-241 authors stated that the application of NPK-based fertilizer at harvest, at the time of stem cuttings of the parent plants, and 7 DAC can significantly increase the shoot dry weight clump-1 (g), grain dry weight clump-1 (g), and weight of 1000-filled grains (g). Therefore the conclusion and the results are slightly contradictory, no superior treatment timing could be identified based on the yield parameters.
 13. The references were cited in the Discussion only to confirm well-known facts.
- Keywords should not be repetitions of the title words, please find such words which are not in the title, this way search engines on the web will find your manuscript with a higher probability.

RESPONSE OF RICE GROWTH AND YIELD UNDER DIFFERENT TIMING OF NPK-BASED FERTILIZER APPLICATION IN THE SALIBU SYSTEM

The manuscript demonstrates a good overall command of English, with clear and professional communication appropriate for scientific discourse. The strengths lie in its logical organization, technical terminology, and readability. However, recurring issues include minor grammatical inaccuracies (e.g., inconsistent article usage), wordy phrasing, awkward expressions, and some typographical errors. Addressing these will improve clarity, conciseness, and precision.

Issues involved:

Line 2: Use plural “FERTILIZERS” when referring to general or multiple applications to align with scientific precision.

Line 8: Replace “based” with “base” for correct adjective use.

Lines 8-9: Streamline by removing redundant words. **Suggested revision:** “...promoting optimal vegetative growth and yield.”.

Lines 11-13: Simplify repetitive phrases (“at the time of”) for brevity. **Suggested revision:** “The timing of NPK-based fertilizer application included four treatments: no application (control), at harvest, at stem cuttings (seven days after harvest, 7 DAH), and seven days after stem cuttings (7 DAC).”.

Lines 16-18: Condense repetitive phrases and retain essential meaning. **Suggested revision:** “Applying NPK-based fertilizer at stem cuttings optimally enhances rice growth and yield in the SS.”.


Line 20: Ensure consistency in terminology across the manuscript. Change “based fertilizer” to “base fertilizer”.

Line 24: Use “term” instead of “abbreviation of” for smoother phrasing.

1. An Itemized Response Sheet

- Respond to all issue/referee opinions/editor suggestions
- All amendments made are highlighted in yellow on the revision paper.

Comments to the Authors from referee:	Addressed (Y/N)	Reply/Action taken
1. Sentences between L25-27 and L32-35 contain the same information. Please revise.	Y	Thank you the suggestion: The sentence in lines 25-27 has been removed. We have used lines 32-35 instead.
2. It is suggested to introduce the ratoon technique and afterwards, authors might highlight what is the modification compared to that in salibu rice cultivation.	Y	The following sentence has been added to the introduction: The ratoon rice technique is a method of rice cultivation that utilizes the rice stems remaining after harvest (ratoon) to regrow and produce the next rice crop. After the main harvest, the remaining rice stems are

		left to regenerate, and new roots and shoots will develop to produce rice in a shorter time frame. In the Salibu system, after the main rice harvest, the remaining stems are not only left to regenerate as in the conventional ratoon technique, but also undergo special management practices such as cutting the main stems, fertilization, weed control, and others to ensure higher and more stable yields.
3. Please add a map of the study site and the environmental conditions might be given, too.	Y	The location map and environmental conditions have been added in the Materials and Methods section (Figure 1):  The site was located at an elevation of 70 m above sea level, with an average air temperature of 26.6 °C and humidity of 83%. Kasihan District is 110°20'40" East Longitude and 7°48'42" South Latitude.
4. It is not completely clear what is the differences between harvest and stem cutting, what is the usual difference between those (in days).	Y	We have explained the difference between harvesting and stem cutting in the research procedure: Harvesting is done after the rice reaches physiological maturity. As parent plant, harvesting is done manually by cutting the stems (along with the panicle) at a height of 15-20 cm above the soil surface using a sickle. Then, 7 days after the harvest, the remaining stems above the soil are cut again at a height of 3 cm from the soil surface to stimulate new shoot growth in the SS technology.

5. Please add the schedule and amount of the irrigation.	Y	It has been added to the research procedures: Water irrigation was applied to the experimental plots until it reached a height of 3 cm above the soil surface. Once it was evenly distributed across all plots, the water channels were closed. This was done every 7 days until the plants reached 70 DAC.
6. Please indicate the row length and row number of an experimental plot.	Y	The suggestion has been added to the experiment design section: The size of each treatment plot can be seen in Figure 2. Figure 2. The size of treatment plot for experimental Each treatment plot size was 1.84 m (length) x 1.84 m (width). Plant spacing within the rows was 0.23 m, with 8 plants per row, making the row length 1.84 m. The spacing between rows was 0.23 m, with a total of 8 rows, resulting in a row width of 1.84 m. The number of rice clumps in each plot was 64.
7. The rice variety might be added.	Y	It has been included in the research procedure section: The rice variety used was Inpari 42
8. Please clarify how many plants were measured/sampled when measuring the variables.	Y	The number of samples measured for each variable has been added in result section: Observations of the tillering stems were conducted on 64 clump plot⁻¹. For plant height, tiller number, dry weight of shoots clump⁻¹ (g), panicle length (cm), weight of 1000-filled grain (g), and grains clump⁻¹ were observed on 10 clump plot⁻¹.

9. Please add $\pm$ SD values to the tables.	Y	Done In Table 1-4, we have added the SD for each observation.
10. Authors stated that at 42 DAC the NPK fertilizer led to a decrease in plant height, but this statement is only valid in the case of 7 days after stem cutting application.	Y	The statement regarding plant height has been revised to: <i>Table 2</i> shows that applying NPK-based fertilizer increased plant height only at the 28 DAC. The application of NPK-based fertilizer at the time of cutting the stem of the parent plants resulted in taller plants compared to applying it at control and 7 DAC. However, at the 42 DAC observation, it appeared that the application of NPK-based fertilizer led to a decrease in plant height.
11. In the Discussion, the authors stated that the application of NPK-based fertilizer significantly impacted plant height in the SS but this statement is not valid for all measurements and each timing. Please be more specific.	Y	In contrast, the application of NPK-based fertilizer significantly impacted plant height at 28 DAC in the SS, although this was not always the case for all plant height observations, particularly at 42 DAC. The delayed application of NPK base fertilizer at 7 DAC actually disrupted plant height growth during the maximum vegetative phase, with inhibition observed at 45 DAC. At 56 DAC, there was a tendency for plant height to be lower than control, although the difference was not significant.
12. The authors concluded the following: The application of NPK-based fertilizer at the time of stem cuttings in the parent plants (7 DAH) resulted in a higher dry weight of milled grains clump ⁻¹ (15.3 g) compared to the control (13.2 g). The research findings indicate that applying NPK-based fertilizer at the time of stem cuttings of the parent plants is the best time to enhance rice growth and yield in the SS. On the other hand in L239-241 authors stated that the application of NPK-based fertilizer at harvest, at the time of stem cuttings of the parent plants, and	Y	We have revised the explanation in Table 4, so the superior treatment is now clearly identified as the application of NPK-based fertilizer at the time of stem cuttings: However, the application of NPK-based fertilizer at the time of stem cuttings significantly increased the grain dry weight clump⁻¹ and was not significantly different with at the time of harvesting and 7 DAC. No significant differences were observed among the treatments at the time of harvesting, 7 DAC, and the control. And we have added it to the discussion in the end of discussion: The application of NPK-based fertilizer at the time of stem cuttings was superior

7 DAC can significantly increase the shoot dry weight clump ⁻¹ (g), grain dry weight clump ⁻¹ (g), and weight of 1000-filled grains (g). Therefore the conclusion and the results are slightly contradictory, no superior treatment timing could be identified based on the yield parameters.		treatment timing for achieving the highest grain dry weight clump ⁻¹ in the SS. The conclusion has been revised to: The results indicated that the application of NPK-based fertilizer can significantly increase the shoot dry weight clump ⁻¹ and weight of 1000-filled grains. Applying of NPK-base fertilizer at the time of stem cuttings enhanced plant height at 28 DAC, with no significant difference compared to the time of harvesting. Furthermore, this treatment resulted in a higher grain dry weight clump ⁻¹ (15.3 g) than the control (13.2 g), and was not significantly different from the treatments at the time of harvesting (13.9 g) or 7 DAC (14.6 g). No significant differences were observed among the treatments at the time of harvesting, 7 DAC, and the control. These findings suggest that applying NPK-base fertilizer at stem cuttings optimally enhances rice yield in the SS. Future research should focus on determining the optimal dosage of NPK-base fertilizer to further maximize rice yields.
13.The references were cited in the Discussion only to confirm well-known facts.	Y	Done
Keywords should not be repetitions of the title words, please find such words which are not in the title, this way search engines on the web will find your manuscript with a higher probability.	Y	The keywords have been changed to words not included in the title, namely: <i>harvesting, main stem, parent plant, ratoon, stem cutting</i>
Issues involved (Editor suggestions)	Addressed (Y/N)	Reply/Action taken
1. Line 2: Use plural “FERTILIZERS” when referring to general or multiple applications to align with scientific precision.	Y	Thank you. The word FERTILIZER is changed to FERTILIZERS
2. Line 8: Replace “based” with “base” for correct adjective use.	Y	Done. The word "NPK-based" has been replaced with " NPK-base " throughout this manuscript.
3. Lines 8-9: Streamline by removing redundant words.	Y	Thank. The sentence in Lines 8-9 has been revised to:

Suggested revision: “...promoting optimal vegetative growth and yield.”		Applying NPK-base fertilizer can effectively promote optimal vegetative growth and yield
4. Lines 11-13: Simplify repetitive phrases (“at the time of”) for brevity. Suggested revision: “The timing of NPK-based fertilizer application included four treatments: no application (control), at harvest, at stem cuttings (seven days after harvest, 7 DAH), and seven days after stem cuttings (7 DAC).”	Y	Done The phrases of “at the time of NPK-base fertilizer ...” in Lines 11-13 has been revised to: The timing of NPK-base fertilizer ...
5. Lines 16-18: Condense repetitive phrases and retain essential meaning. Suggested revision: “Applying NPK-based fertilizer at stem cuttings optimally enhances rice growth and yield in the SS.”	Y	Done
6. Line 20: Ensure consistency in terminology across the manuscript. Change “based fertilizer” to “base fertilizer”.	Y	Done
7. Line 24: Use “term” instead of “abbreviation of” for smoother phrasing.	Y	Done

2. A Paper Revised for Resubmit:

RESPONSE OF RICE GROWTH AND YIELD UNDER DIFFERENT TIMING OF **NPK-BASE** FERTILIZERS APPLICATION IN THE SALIBU SYSTEM

AEER_16969

Abstract: In the Salibu system (SS), rice plants have a shorter growth cycle than their parent plants. Applying **NPK-base** fertilizer can **promote** optimal vegetative growth and yield. This study aimed to determine the optimal timing of NPK-base fertilizer application to enhance rice yield in the SS. The experiment was arranged in a randomized complete block design (RCBD) with three replications, where each replication acted as a block. The timing of **NPK-base** fertilizers application included four treatments: no application (control), at **harvest**, stem cuttings (seven days after harvest, or 7 DAH), and seven days after stem cuttings (7 DAC). The results indicated that the application of NPK-base fertilizers can significantly increase the shoot dry weight clump⁻¹ and 1000-filled grains. Applying NPK-base fertilizer at stem cuttings enhanced plant height at 28 DAC, with no significant difference compared to harvest. Furthermore, this treatment resulted in a higher grains dry weight clump⁻¹ (15.3 g) than the control (13.2 g), and was not significantly different from the treatments at harvest (13.9 g) or 7 DAC (14.6 g). No significant differences were observed among the treatments at harvest, 7 DAC, and the control. The research findings indicate that applying NPK-

base fertilizer at stem cuttings of the parent plants optimally enhances the rice yield in the SS. Future research should recommend determining the optimal dosage of NPK-base fertilizer to maximize rice growth and yield.

Keywords: *harvest, main stem, parent plant, ratoon, stem cutting*

INTRODUCTION

As Indonesia's population grows, the demand for food, particularly rice, is increasing. Various measures have been implemented to achieve national rice self-sufficiency, including adapting the SS. The term of Salibu is derived from the Indonesian phrase “Salin ibu”, which translates to “change of the parent plants” in English.

The ratoon rice technique is a rice cultivation method that utilizes the rice stems remaining after harvest (ratoon) to regrow and produce the next rice crop. After the main harvest, the remaining rice stems are left to regenerate, and new roots and shoots will develop to make rice in a shorter time frame. In the SS, after the main rice harvest, the remaining stems are not regenerated as in the conventional ratoon technique but also undergo special management practices such as cutting the main stems, fertilization, weed control, and others to ensure higher and more stable yields.

The SS features a shorter life cycle than parent plants, enabling increased productivity by applying NPK-base fertilizer. Furthermore, using NPK-base fertilizer can accelerate the growth of rice shoots during the early stages, thereby enhancing the potential for higher yields.

The SS is a rice cultivation method based on local wisdom, originally implemented in West Sumatra, Indonesia. The technology has been tested in various regions across Indonesia. Although farmers have not widely adopted it due to perceptions of lower yields, this technology can enhance rice harvests and improve land-use efficiency with proper management. According to Abdulrachman et al. (2015), the SS offers advantages such as faster harvest times, more efficient water usage, and reduced production costs by up to 45% compared to the transplanting method. This method reduces land preparation, seedling production, and planting expenses. Therefore, the SS has the potential to improve the welfare of farmers.

The success of the SS technology is influenced by various factors, one of which is the timing of NPK-base fertilizer application. The NPK fertilizer plays a crucial role in supporting the growth of the new shoots after the parent plants are cut. Applying the fertilizer can provide the essential nutrients the new shoots need, promoting their development. Since the SS has a shorter harvest cycle than its parent plants, it is important to determine the optimal fertilizer application timing to accelerate the growth of the new shoots. However, research on the timing of NPK-base fertilizer in the SS remains limited. Therefore, further studies are necessary to identify the optimal timing for fertilizer application in the SS.

In Indonesia, rice cultivation in irrigated land typically occurs one to three times annually, depending on the irrigation system employed. In rainfed fields, rice can be planted only once a year, while fully irrigated fields can support up to three planting cycles each year. Each planting season involves various activities, including land preparation, seedling production, planting, plant maintenance, and harvesting. Conventional rice cultivation often incurs relatively high costs, lowering profit margins. In contrast, the SS offers greater advantages due to its land and resource utilization efficiency.

The SS can enhance rice production and land productivity annually. This method utilizes the shoots that emerge from the basal stems after cuttings. To maximize the number of tillers, apply the optimal dosage of NPK-base fertilizer. New shoots emerge from the nodes of the remaining stem, followed by the emergence of new roots. Eventually, these shoots grow into independent plants. The formed shoots subsequently develop into new tillers, similar to the rice planting system used in transplanting techniques (Juanda, 2016). The SS can enhance food availability in

Indonesia and contribute to food security and resilience. It is more effectively implemented on land that is not always flooded (Jahari & Sinaga, 2019). Rice productivity in tropical regions primarily relies on fertilizers, as soils often lack nitrogen and phosphorus (Hindersah et al., 2022).

Nitrogen (N), phosphorus (P), and potassium (K) are applied in large quantities as fertilizers in rice fields, and a deficiency of any of these nutrients can reduce crop yields (Shrestha et al., 2020). Nitrogen plays a crucial role in the synthesis of chlorophyll and protein. Phosphorus is involved in energy storage and transfer, and it is a key element in nucleic acids, coenzymes, nucleotides, phosphoproteins, phospholipids, and phosphate sugars. Meanwhile, potassium supports starch formation, activates enzymes, and acts as a catalyst in storing photosynthetic products.

Nitrogen is an essential element that forms plants' amino acids, proteins (enzymes), nucleic acids, and chlorophyll. A nitrogen deficiency in rice cultivation leads to reduced plant height, lower tiller formation, decreased leaf area index, and diminished photosynthesis rates (Chauhan et al., 2017). Nitrogen plays a crucial role in promoting growth, enhancing yields, and improving grain quality by improving the number of tillers, expanding leaf area, and facilitating protein synthesis. Plants absorb nitrogen through nitrate ions (NO_3^-) and ammonium ions (NH_4^+) (Kaya, 2013). A lack of nitrogen can limit rice productivity, while excessive nitrogen can result in agronomic and economic losses (Hussain et al., 2023). The yield increased when nitrogen was applied at $202.15 \text{ kg ha}^{-1}$. The fertilization frequency was four times (He et al., 2024). Optimal nitrogen fertilization can effectively compensate for the negative impacts of N fertilizer, enhancing grain quality and nitrogen use efficiency (NUE) while sustaining high yields (Tian-Yang et al., 2022).

Phosphorus (P) is essential in photosynthesis, respiration, energy transfer, storage, cell division, and enlargement. Plants absorb phosphorus in the form of primary orthophosphate ions, H_2PO_4^- (Irwanto, 2014). Additionally, phosphorus is a key component of high-energy compounds such as adenosine triphosphate (ATP) and synthesizes phospholipids, nucleotides, and glycoproteins. Phosphorus deficiency can hinder growth and yield in rice, which is evident through reduced tiller numbers and dry matter accumulation, along with symptoms such as stunted growth and upright leaves that exhibit a dark green colour (Chauhan et al., 2017).

Potassium (K) activates various enzymes in plant metabolic processes and is absorbed as potassium ions (K^+) (Irwanto, 2014). This nutrient is essential for enhancing root growth, promoting plant vigour, and increasing resistance to pests and diseases. Symptoms of potassium deficiency in rice include stunted growth with fewer tillers, dark green upper leaves, and chlorosis in the areas between leaf veins (Chauhan et al., 2017). Although chemical fertilizers can improve rice yield, the response highly depends on optimal fertilization management in paddy fields (Yang et al., 2021). Using a combination of $135\text{--}210 \text{ kg ha}^{-1}$ N and $115\text{--}137 \text{ kg ha}^{-1}$ K application to achieve relatively higher yield and improve quality in rice production (Chen et al., 2024).

The NPK Phonska (15-15-15) fertilizer is complex inorganic NPK with composition of 15% N, 15% P_2O_5 , 15% K_2O , and 10% S. This fertilizer is widely utilized by farmers due to its effectiveness in enhancing rice yields and grain quality. According to Shrestha et al. (2020), nitrogen (N), phosphorus (P), and potassium (K) are applied as fertilizers in large quantities to rice fields. Research conducted by Kurniadie (2002), using a dosage of 300 kg ha^{-1} NPK Phonska along with 333 kg ha^{-1} ZA resulted in the highest performance for the IR 64 rice variety, including parameters such as the number of panicle clump $^{-1}$, the number of grain panicle $^{-1}$, the harvest

index, the weight of 1000 grains, as well as dry grain yields (7.50 tons ha⁻¹) and milled rice yields (6.17 tons ha⁻¹).

Optimized fertilization practices represent a promising management approach for achieving sustainable rice production (Zhuang et al., 2022). The application of NPK Phonska fertilizer at a rate of 300 kg ha⁻¹, combined with 200 kg ha⁻¹ urea, significantly increased plant height, the number of tillers, and yield up to 7.448 tons ha⁻¹, with relative agronomic effectiveness (RAE) value of 106.46 (Fidiyawati et al., 2022). Combining NPK fertilizer with urea is essential to meet nitrogen nutrient requirements during various rice growth. Without urea addition, higher dosages of NPK were needed; specifically, 300 kg ha⁻¹ NPK Phonska (15-15-15) required an additional 150 kg ha⁻¹ urea (Hartatik & Widowati, 2015). The greatest yield of 5.81 tons ha⁻¹ was achieved with the combined treatment of 250 kg ha⁻¹ NPK + 150 kg ha⁻¹ urea + 1,000 kg ha⁻¹ dolomite with a yield of 5.81 tons ha⁻¹ (Ratmini et al., 2023).

Applying NPK fertilizer can enhance plant height and yield by 3%, although it may reduce the weight of 1000 grains. On the other hand, nitrogen fertilizer significantly contributed to an increase in productive tillers and boosted yield by 6%. The most effective combination for rice fields with high phosphorus levels was 300 kg ha⁻¹ NPK and 100 kg ha⁻¹ urea (Siska & Ismon, 2019). The NPK application also enhances the absorption of nutrients N, P, and K, and improves grain yield (Shrestha et al., 2020). Additionally, Srinivas (2017) noted a significant increase in the number of tillers and panicles m⁻² with the application of 120-60-60 kg ha⁻¹ NPK.

The application of NPK fertilizer significantly influenced plant height at 35, 45, and 90 days after planting (DAP), the number of tillers at 35 and 45 DAP, and various production parameters such as the number of panicles clump⁻¹ and 1000-filled grains of rice (Waty et al., 2014). However, a dosage of 50 kg ha⁻¹ NPK Phonska did not show significant differences in rice production across various agroecological zones (highlands, midlands, and lowlands) using the SS (Fitri et al., 2019). In the SS, fertilization adhered to the parent plants: the first application of 40% recommended dosage at 15-20 DAP and the second consisting of 60% at 30-35 DAP (Abdulrachman et al., 2015).

Based on the background and literature review, NPK fertilizer is more commonly used under the transplanting system in rice cultivation. In contrast, its application as base fertilizer in the SS has not been thoroughly investigated. Therefore, the study aimed to determine the optimal timing of NPK-base fertilizer application to enhance rice yield in the SS.

MATERIALS AND METHODS

Study Site

The research was conducted from January to June 2024 in Tirtonirmolo Village, Kasihan District, Bantul Regency, Special Region of Yogyakarta, Indonesia. A map of the study site can be seen in Figure 1.




Figure 1. A map showing the study site in Tirtonirmolo Village, Kasihan District, Bantul Regency, Special Region of Yogyakarta, Indonesia

The site was located at an elevation of 70 m above sea level, with an average air temperature of 26.6 °C and humidity of 83%. Kasihan District is 110°20'40" East Longitude and 7°48'42" South Latitude.

Experiment Design

The experiment was arranged in an RCBD with three replications, where each replication acted as a block. The timing of NPK-base fertilizer application included four treatments: no application (control), at harvest, at stem cuttings, or seven days after harvest (7 DAH), and at seven days after cutting the stems (7 DAC). This study required 12 treatment plots. The size of each treatment plot can be seen in Figure 2.




Figure 2. The size of the treatment plot for experimental

Each treatment plot size was 1.84 m (length) x 1.84 m (width). Plant spacing within the rows was 0.23 m, with 8 plants per row, making the row length 1.84 m. The spacing between rows was 0.23 m, with a total of 8 rows, resulting in a row width of 1.84 m. The number of rice clumps in each plot was 64. The research flow is shown in Figure 3.




Figure 3. The flow diagram in research activity

Research Procedures

Parent plants

In the SS, the parent plants cultivated by farmers were utilized. The rice variety used was Inpari 42, which has a harvesting period of 87 DAP. The spacing was set at 23×23 cm. The land used for cultivation was technically irrigated, ensuring a consistent water supply, and consisted of alluvial soil. As the parent plant, harvesting was done manually by cutting the stems (along with the panicle) at a height of 15-20 cm above the soil surface using a sickle.

Salibu system

Then, 7 days after the harvest, the remaining stems above the soil were cut again at a height of 3 cm from the soil surface to stimulate new shoot growth in the SS technology. The new shoots emerged from the remaining cut of parent stems and were carefully maintained. Plant care included irrigation, fertilization, and the management of pests, diseases, and weeds to ensure optimal growth.

Water irrigation was applied to the experimental plots until it reached a height of 3 cm above the soil surface. Once it was evenly distributed across all plots, the water channels were closed. This was done every 7 days until the plants reached 70 DAC.

The application of NPK-base fertilizer occurred according to the treatment schedule, which took place at harvest, at stem cuttings (7 DAH), and 7 DAC. The dosage of NPK-base fertilizer used was 60 g plot^{-1} , equivalent to 150 kg ha^{-1} NPK Phonska, except in the control plots. After applying NPK-base fertilizer, a subsequent fertilizer was recommended with 150 kg ha^{-1} NPK Phonska and 200 kg ha^{-1} urea. It was given in two stages: the first stage (40% of the recommended dosage) was applied at 14 DAC, followed by the second stage (60% of the recommended dosage) at 28 DAC.

Weeding was performed at 12 and 26 DAC to reduce weed competition. Pest and disease management was implemented according to integrated pest management (IPM) principles at 56

and 63 DAC, using Decis 25 EC as the control agent. Rice harvesting occurred at 75 DAC when the rice reached physiological maturity (95% yellowing). With the appropriate maintenance and fertilization practices, the rice yield in the SS was expected to be maximized.

Parameters

The research observations included the growth and yield components from each plot treatment. The plant growth components comprised the percentage of tillering stems (%), plant height (cm), the number of tillers clump⁻¹ (stems), and dry weight of shoots (leaves and stems) clump⁻¹ (g). In addition, the observed yield components included the grains dry weight clump⁻¹ (g), the weight of 1000-filled grains (g), and panicle length (cm).

Statistical Analysis

The observation data were analyzed using analysis of variance (ANOVA) at 5% significant level (Gomez & Gomez, 1984) with IBM SPSS Version of 23 software. If differences were found between treatment means, further testing was conducted using Duncan's new multiple range test (DMRT) at 5% significant level. The term of SD is derived from the abbreviation of standard deviation.

RESULTS

Component of the Rice Growth

Tillering Stems


Observations of the tillering stems were conducted on 64 clumps plot⁻¹ at 14 DAC. The results of the ANOVA (*Appendix 1*) indicate that the timing of NPK-base fertilizer application did not significantly affect the percentage of tillering stems that sprouted. The number of tillering stems is presented in *Table 1*.

Table 1. The number of tillering stems (%) at 14 DAC

The timing of NPK-base fertilizer application			
Control	At harvest	At stem cuttings	At 7 DAC
92.7 a	93.8 a	97.9 a	97.4 a

Remarks: The values in the same row with different letters are significantly different based on DMRT at 5% significant level. $SD_{\text{tillering stems}} = 2.617$.

Figure 3a shows shoot emergence from the basal stem of the parent at 7 DAC, while *Figure 3b* illustrates the tiller growth at 28 DAC.



a. The shoot emergence from the basal stem of the parents at 7 DAC



b. The tiller growth in the SS at 28 DAC

Figure 3. Photo of the shoot emergence from the basal stem of the parents at 7 DAC (a) and the tiller growth at 28 DAC (b)

Plant height

Plant height was observed on 10 clumps plot⁻¹ at regular intervals of 14, 28, 42, and 56 DAC. The results of the ANOVA (*Appendix 2*) find that the timing of NPK-base fertilizer application significantly affected the plant height at 28 and 42 DAC, while there was no significance at 14 and 56 DAC. The average plant height is presented in *Table 2*.

Table 2. The plant height at 14, 28, 42, and 56 DAC

Time of observations (DAC)	The timing of NPK-base fertilizer application			
	Control	At harvest	At stem cuttings	At 7 DAC
14	33.9 a	32.3 a	42.1 a	36.0 a
28	62.4 b	70.3 ab	73.3 a	63.3 b
42	87.0 a	86.9 a	84.3 ab	80.8 b
56	90.1 a	91.3 a	89.5 a	86.8 a

Remarks: The values in the same row with different letters are significantly different based on DMRT at 5% significant level. $SD_{14\text{ DAC}} = 4.577$, $SD_{28\text{ DAC}} = 4.322$, $SD_{42\text{ DAC}} = 2.211$, and $SD_{28\text{ DAC}} = 3.681$.

Table 2 shows that applying NPK-base fertilizer increased only plant height at the 28 DAC. The application of NPK-base fertilizer at stem cuttings of the parent plants resulted in taller plants compared to applying it at control and 7 DAC, although no different with at harvest. However, at the 42 DAC observation, it appeared that the application of NPK-base fertilizer at 7 DAC led to a decrease in plant height compared to control.

Tiller numbers

The number of tillers was also observed on 10 clumps plot⁻¹ at regular intervals of 14, 28, 42, and 56 DAC. The results of the ANOVA (*Appendix 3*) show that the timing of NPK-base fertilizer application had no significant effect on the number of tillers clump⁻¹ at 14, 28, 42, and 56 DAC. The average number of tillers is presented in *Table 3*.

Table 3. The number of tiller clump⁻¹ at 14, 28, 42, and 56 DAC

Time of observations (DAC)	The timing of NPK-base fertilizer application			
	Control	At harvest	At stem cuttings	At 7 DAC
14	7.3 a	6.4 a	7.9 a	6.0 a
28	14.6 a	15.9 a	16.9 a	13.6 a
42	17.5 a	18.2 a	19.0 a	15.2 a
56	16.2 a	15.9 a	16.2 a	13.8 a

Remarks: The values in the same row with different letters are significantly different based on DMRT at 5% significant level. $SD_{14\text{ DAC}} = 2.804$, $SD_{28\text{ DAC}} = 2.567$, $SD_{42\text{ DAC}} = 3.868$, and $SD_{28\text{ DAC}} = 3.636$.

Component of the Rice yield

The dry weight of shoots clump⁻¹ (g), grains clump⁻¹ (g), weight of 1000-filled grain (g), and panicle length (cm) were observed on 10 clumps plot⁻¹ at 75 DAC. The results of the ANOVA (Appendix 4) indicate that the timing of NPK-base fertilizer application significantly affected the dry weight of shoots, grains clump⁻¹, and 1000-filled grains, while panicle length showed no significant effect. The average of dry weight of shoots, grains, 1000-filled grains, and panicle length are presented in Table 4.

Table 4. The means of shoot dry weight, grain dry weight, the weight of 1000-filled grain, and panicle length at 75 DAC.

Parameters observed	The timing of NPK-base fertilizer application			
	Control	At harvest	At stem cuttings	At 7 DAC
Shoots dry weight clump ⁻¹ (g)	23.1 b	31.3 a	32.2 a	30.1 a
Grains dry weight clump ⁻¹ (g)	13.2 b	13.9 ab	15.3 a	14.6 ab
1000-filled grains (g)	13.3 b	19.0 a	19.3 a	17.3 a
Panicle length (cm)	18.9 a	18.6 a	20.5 a	19.8 a

Remarks: The values in the same row with different letters are significantly different based on DMRT at 5% significant level. $SD_{\text{shoot dry weight}} = 2.778$, $SD_{\text{grains dry weight}} = 0.703$, $SD_{\text{1000-filled grains}} = 1.803$, and $SD_{\text{panicle length}} = 1.682$.

Table 4 shows that the application of NPK-base fertilizer -can significantly increase the shoot dry weight clump⁻¹ (g), grain dry weight clump⁻¹ (g), and 1000-filled grains (g). However, the application of NPK-base fertilizer at stem cuttings significantly increased the grain dry weight clump⁻¹ and was not significantly different at harvest and 7 DAC. No significant differences were observed among the treatments at harvest, 7 DAC, and the control.

DISCUSSION

Components of Rice Growth

The application of NPK-base fertilizer in the SS did not influence the percentage of new shoots emerging from cut stems and tillers number. This may be because the new shoots primarily rely on the food reserve stored in the parent stems. After cutting, the emerging shoots utilize the resources available in the remaining stem for growth. Therefore, although NPK provides essential

nutrients, internal factors such as the energy stored in the parent stem may be more dominant in determining the success of new shoot emergence. While NPK contains essential nutrients, the growth of tillers is influenced by various factors, including the genetic traits of the rice variety, soil conditions, and management techniques. In the SS, which emphasizes effective resource use, the number of tillers may be more influenced by care practices and other cultivation techniques rather than by fertilizer alone. Thus, applying NPK as a fertilizer was insufficient to increase the number of tillers significantly.

In contrast, the application of NPK-base fertilizer significantly impacted plant height at 28 DAC in the SS, although this was not always the case for all plant height observations, particularly at 42 DAC. The delayed application of NPK base fertilizer at 7 DAC actually disrupted plant height growth during the maximum vegetative phase, with inhibition observed at 45 DAC. At 56 DAC, there was a tendency for plant height to be lower than control, although the difference was not significant.

The NPK provides nitrogen, phosphorus, and potassium, which are crucial for vegetative growth. Nitrogen plays a role in protein synthesis and leaf growth, directly affecting plant height. Additionally, phosphorus supports better root development, allowing the plant to absorb more nutrients from the soil. With optimal nutrient availability, the plant can grow taller, demonstrating that while NPK did not affect new shoots and tillers numbers, it was key in supporting plant height growth. According to Samira et al. (2012), the application of NPK can increase the plant height of rice.

Components of Rice Yield

In the early stages of growth, new shoots utilize the food reserves from the remains of the parent stems while forming new roots that replace the function of the parent plant's roots. New shoots were ready to utilize nutrients available in their environment 10 days after the stem cuttings. The application of NPK-base fertilizer at stem cuttings the parent enhanced the availability of essential nutrients, including nitrogen (N), phosphorus (P), and potassium (K), which were crucial for rice growth in the SS. The new shoot's roots quickly absorbed available nutrients to support their development. This base fertilizer helps provide the essential nutrients needed by the new shoots that emerge from the stem cuttings. New shoots will appear the day after cutting, although small and without leaves.

The application of NPK-base fertilizer significantly increased shoot dry weight compared to the control. This fertilizer contains essential nutrients such as nitrogen, phosphorus, and potassium, crucial for plant growth. Nitrogen plays a vital role in chlorophyll formation and leaf development. With the application of NPK, plants can develop denser leaves and wider shoots, enhancing their photosynthetic capacity. As a result, the shoots' dry weight was higher than those that did not receive the fertilizer. Yang et al. (2004) emphasized that the rice root system was vital in absorbing water and nutrients from the soil. According to Wang et al. (2006), root system formation was closely coordinated with shoot growth, where the shoot regulated the root development through the circulation of materials between the two. Root distribution determines the structure of the root system within the soil, affecting the acquisition of soil resources such as nutrients and water.

In addition to increasing shoots dry weight, it positively influenced grain dry weight clump⁻¹ higher than the control. The nutrients provided by NPK support the grain-filling process, enabling the plants to produce more and heavier grains. Applying nitrogen and phosphorus is

crucial for optimal grain development during the grain-filling phase. Plants receiving NPK-base fertilizer show a significant increase in the dry weight of grains clump^{-1} , indicating that a good nutrient supply contributes to better harvest yields. The product of photosynthesis in the leaves can be translocated in the seed-filling process if the plants are supplied with this nutrient. According to Rawat et al. (2022), phosphorus is the most abundant element after nitrogen and is involved in many physiological processes such as photosynthesis and translocation. Sarkar et al. (2021) stated that the capacity for source-sink dynamics and assimilate translocation can play an important role during grain formation. As explained by Wei et al. (2018), rice grain yield depends on the supply assimilates (source) and the grain capacity to utilize them (sink).

The application of NPK resulted in the resulting grains being heavier and of higher quality than the control. Adequate nutrients aid in grain-filling, allowing the grains to fill out properly and achieve greater dry weight. The application of NPK-base fertilizer not only increased the quantity of the harvest but also improved the quality of the rice grains, which is essential for determining the market value of the produce. According to Metwally et al. (2020), the grain quality characteristics can be enhanced by the NPK fertilizer application. Explain by Pasaribu et al. (2018), the parent plant's ability to generate new shoots highly depends on the carbohydrate content and phytohormones retained in the intercalary meristem tissues of the stubble after harvest. Additionally, increasing the rice yield with the SS can be achieved by enhancing fertilization, especially nitrogen.

Therefore, applying NPK-base fertilizer at stem cuttings was more effective in increasing grain dry weight clump^{-1} than using it at harvest or 7 DAC. The application of NPK-base fertilizer at stem cutting provided essential nutrients for the new shoots that emerged from the remaining parts of the parent plants. The fertilizer applied at this time can be directly utilized by the shoots after ten days, as nitrogen, phosphorus, and potassium are crucial for accelerating the formation of new tissues and photosynthesis. In contrast, applying NPK-base fertilizer at harvest or 7 DAC was less effective because the growth of new shoots had not yet started during that period. The availability of nutrients was not optimally utilized, leading to lower dry weight and harvest yields. Therefore, the timing of NPK-base fertilizer application significantly influenced nutrient absorption efficiency and the final yield of the plants. Moreover, the application of NPK-base fertilizer at stem cuttings optimally enhances the rice yield in the SS.

Conclusions

The results indicated that the application of NPK-base fertilizer can significantly increase the shoot dry weight clump^{-1} and 1000-filled grains. The applying of NPK-base fertilizer can enhance the plant height (cm) at 28 DAC, with no significant difference compared to the time of harvesting. Furthermore, this treatment resulted in a higher grains dry weight clump^{-1} (15.3 g) compared to the control (13.2 g), and was not significantly different from the treatments at harvest (13.9 g) or 7 DAC (14.6 g). No significant differences were observed among the treatments at harvest, 7 DAC, and the control. The research findings indicate that applying NPK-base fertilizer at stem cuttings of the parent plants optimally enhances rice yield in the SS. Future research should recommend determining the optimal dosage of NPK-base fertilizer to maximize rice yields.

Acknowledgements. We sincerely thank the --

REFERENCES

- [32] Abdulrachman, S., Suhartatik, E., Erdiman, Susilawati, Zaini, Z., Jamil, A., Mejaya, M. J., Sasmita, P., Abdulah, B., Suwarna, Baliadi, Y., Dhalimi, A., Sujinah, Suharna, Ningrum, E. S. (2015): Panduan teknologi budidaya padi salibu. Badan Penelitian dan Pengembangan Pertanian, Kementerian Pertanian. (in Indonesian)
- [33] Chauhan, B. S., Jabran, K., Mahajan, G. (2017): Fertilizer management in rice. – In: Rice Production Worldwide (Issue February 2017, pp. 1–563).
- [34] Chen, G., Duan, Q., Wu, C., He, X., Hu, M., Li, C., Ouyang, Y., Peng, L., Yang, H., Zhang, Q., Jiang, Q., Lan, Y., Li, T. (2024): Optimizing rice yield, quality, and nutrient use efficiency through combined application of nitrogen and potassium. – *Frontiers in Plant Science* 15: 1–16.
- [35] Fidiyawati, E., Latifah, E., Krismawati, A., Sihombing, D., Setyorini, D., Bakar, A., Sugiono. (2022): Effectivity of inorganic fertilizer NPK (15-15-6) to growth and yield of lowland rice (*Oryza sativa* L.) on alfisol soil. – In: IOP Conference Series: Earth and Environmental Science. The 4th International Conference on Food and Agriculture 980: 012014.
- [36] Fitri, R., Erdiman, Kusnadi, N., Yamaoka, K. (2019): Salibu technology in Indonesia: An alternative for efficient use of agricultural resources to achieve sustainable food security. – *Paddy and Water Environment* 17: 403–410.
- [37] Gomez, A. G., & Gomez, K. A. (1984): Statistical procedures for agricultural research (Second edition). John Wiley & Sons, Inc. New York.
- [38] Hartatik, W., Widowati, L. R. (2015): Effect of compound fertilizer NPKS and NPK on rice growth and yield on Inceptisol. – *Jurnal Penelitian Pertanian Tanaman Pangan* 34(3): 175–186. (in Indonesian)
- [39] He, X., Zhu, H., Shi, A., Wang, X. (2024): Optimizing nitrogen fertilizer management enhances rice yield, dry matter, and nitrogen use efficiency. – *Agronomy* 14: 919.
- [40] Hindersah, R., Kalay, A. M., Talahaturuson, A. (2022): Rice yield grown in different fertilizer combination and planting methods: Case study in Buru Island, Indonesia. – *Open Agriculture* 7: 871–881.
- [41] Hussain, T., Mulla, D. J., Hussain, N., Qin, R., Tahir, M., Liu, K., Harrison, M. T., Sinutok, S., Duangpan, S. (2023): Optimizing nitrogen fertilization to enhance productivity and profitability of upland rice using CSM–CERES–Rice. – *Plants* 12: 3685.
- [42] Irwanto. (2014): Pengaruh pemberian pupuk NPK terhadap pertumbuhan tanaman buah naga di Kecamatan Pemayang, Kabupaten Batanghari, Provinsi Jambi. Balai Pelatihan Pertanian Jambi. (in Indonesian)
- [43] Jahari, M., Sinaga, P. H. (2019): Overcome limited growing season with cultivation ratoon and salibu. – *Jurnal Dinamika Pertanian* 3: 65–72.
- [44] Juanda, R. (2016): Peningkatan produksi padi melalui potensi dan pengembangan wilayah produksi benih unggul di Provinsi Aceh. – *Agrosamudra* 3(2): 72–80. (in Indonesian)
- [45] Kaya, E. (2013): Effect of rice straw compost and fertilizer NPK on N-available of soil, N-uptake, growth and yield of paddy (*Oryza sativa* L.). – In: Prosiding FMIPA Universitas Pattimura 2013 (pp. 41–47). (in Indonesian)
- [46] Kurniadie, D. (2002): Pengaruh kombinasi dosis pupuk majemuk NPK phonska dan pupuk N terhadap pertumbuhan dan hasil tanaman padi sawah (*Oriza sativa* L.) varietas IR 64. – *Jurnal Bionatura* 4(3): 137–147. (in Indonesian)
- [47] Metwally, T. F., Mohamed, A. A. E., Sorour, S. G. R., Elsayed, G. A. (2020): Stimulating compounds affect the grain quality characteristics and the nutritional value of rice (*Oryza sativa*). – *Applied*

Ecology and Environmental Research 18(5): 6829–6840.

- [48] Pasaribu, P. O., Triadiati, Anas, I. (2018): Rice ratooning using the salibu system and the system of rice intensification method influenced by physiological traits. – *Pertanika* 41(2): 637–654.
- [49] Ratmini, N. P. S., Subiksa, I. G. M., Suprihatin, A., Suastika, I. W., Riyanto, D. (2023): Optimization of lowland productivity through fertilization as a food buffer against the impact of climate change. – *E3S Web of Conferences*, 9th ICCO 2023, 467: 01019.
- [50] Rawat, J., Pandey, N., Saxena, J. (2022): Role of potassium in plant photosynthesis, transport, growth, and yield. – In: *Role of Potassium in Abiotic Stress*. Springer Nature Singapore Ltd. (pp. 1–14).
- [51] Samira, D., Gani, A., Slavich, P., McLeod, M. (2012): Effect of NPK fertilizer and biochar residue on paddy growth and yield of second planting. – In: 2nd Annual International Conference Syiah Kuala University 2012 & The 8th IMT-GT Uninet Biosciences Conference, Banda Aceh, 22-24 November 2012, 2(1): 22–24.
- [52] Sarkar, S., Monshi, F. I., Uddin, M. R., Tabassum, R., Sarkar, M., Hasan, A. (2021): Source-sink manipulation influences the grain-filling characteristics associated with the grain weight of rice. – *Journal of Innovative Agriculture* 8: 20–29.
- [53] Shrestha, J., Shah, K. K., Timsina, K. P. (2020): Effects of different fertilizers on growth and productivity of rice (*Oryza sativa* L.): a review. – *International Journal of Global Science Research* 7(1): 1291–1301.
- [54] Siska, W., Ismon, L. (2019): Pemupukan NPK dan nitrogen pada tanaman padi di lahan sawah berstatus P tinggi di Sumatera Barat. – *Jurnal Pengkajian dan Pengembangan Teknologi Pertanian* 22(2): 175–184. (in Indonesian)
- [55] Srinivas, M. (2017): Response of rice (*Oryza sativa* L.) to different nutrient levels under machine planted conditions. – In: 5th International Conference on Agriculture, Ecology and Biological Engineering, Thailand, May 2-4, 2017 (pp: 1–5).
- [56] Tian-Yang, Z., Zhi-Kang, L., En-Peng, L., Wei-Lu, W., Li-Min, Y., Hao, Z., Li-Jun, L., Zhi-Qin, W., Jun-Fei, G., Jian-Chang, Y. (2022): Optimization of nitrogen fertilization improves rice quality by affecting the structure and physicochemical properties of starch at high yield levels. – *Journal of Integrative Agriculture* 21(6): 1576–1592.
- [57] Wang, H., Inukai, Y., Yamauchi, A. (2006). Root development and nutrient uptake. – *Critical Reviews in Plant Sciences* 25(3): 279–301.
- [58] Waty, R., Musyassir, Syamaun, Chairunnas. (2014): Effects of NPK fertilization and biochar residual to physical properties of soil, plant growth and result of rice field (*Oryza sativa* L.) second planting season. – *Jurnal Manajemen Sumber Daya Lahan* 3(1): 383–389.
- [59] Wei, H., Meng, L., Li, X., Dai, Q., Zhang, H., Yin, X. (2018): Sink-source relationship during rice grain filling is associated with grain nitrogen concentration. – *Field Crops Research* 2015: 23–38.
- [60] Yang, C., Yang, L., Yang, Y., Ouyang, Z. (2004): Rice root growth and nutrient uptake as influenced by organic manure in continuously and alternately flooded paddy soils. – *Agricultural Water Management* 70: 67–81.
- [61] Yang, G., Nabi, F., Sajid, S., Kaleri, A. R., Jakhar, A. M., Cheng, L., Raspor, M., Muhammad, N., Ma, J., Hu, Y. (2021): Response of root development and nutrient uptake of two Chinese cultivars of hybrid rice to nitrogen and phosphorus fertilization in Sichuan Province, China. – *Molecular Biology Reports* 48(12): 8009–8021
- [62] Zhuang, Y., Ruan, S., Zhang, L., Chen, J., Li, S., Wen, W., Liu, H. (2022): Effects and potential of optimized fertilization practices for rice production in China. – *Agronomy* 42(32): 1–16.

APPENDIX

Appendix 1. ANOVA of tillering stem

Source of variance	Degree of freedom	Sum of Squares	Mean square	F calc.	F table 5%
Block	2	37.028	18.514	2.703 ns	5.14
Treatment	3	60.832	20.277	2.960 ns	4.76
Error	6	41.097	6.850		
Total	11	138.957			

Remarks: ns = not significant at 5%

Appendix 2. ANOVA of plant height

Source of variance	Degree of freedom	Mean square				F table 5%
		14 DAC	28 DAC	42 DAC	56 DAC	
Block	2	9.027 ns	6.754 ns	5.127 ns	4.868 ns	5.14
Treatment	3	55.394 ns	84.675 *	25.636 *	10.640 ns	4.76
Error	6	20.947	18.678	4.934	13.549	
Total	11					

Remarks: ns = not significant at 5% and * = significance different at 5%

Appendix 3. ANOVA of tiller number

Source of variance	Degree of freedom	Mean square				F table 5%
		14 DAC	28 DAC	42 DAC	56 DAC	
Block	2	9.896 ns	17.453 ns	20.818 ns	12.766 ns	5.14
Treatment	3	2.200 ns	6.444 ns	8.132 ns	4.125 ns	4.76
Error	6	7.861	4.960	10.220	14.141	
Total	11					


Remarks: ns = not significant at 5%

Appendix 4. ANOVA of shoot dry weight, grain dry weight, 1000-filled grains, and panicle length


Source of variance	Degree of freedom	Mean square				F table 5%
		Shoot dry weight	Grain dry weight	1000-filled grains	Panicle length	
Block	2	13.898 ns	0.259 ns	0.250 ns	0.002 ns	5.14
Treatment	3	51.490 *	2.570 *	22.750 *	2.207 ns	4.76
Error	6	7.719	0.494	3.250	2.829	
Total	11					

Remarks: ns = not significant at 5% and * = significance different at 5%


Bukti Submit A Revised Paper with Highlights: 15 Januari 2025



Invoice Pembayaran: 24 Januari 2025



URL: <https://mail.google.com/mail/u/0/#sent/FMfcgzQZSsCxdCdZBNTJHMGbSMGgmVCW>



PAYMENT DETAILS:

Name of Beneficiary (Recipient): Aloki Institute Ltd.
Postal Address of Beneficiary: Kassa utca 118., 1185 Budapest, Hungary
IBAN Code: HU53
International Bank Account Number (IBAN): HU53 1201 1265 0143 7534 0020 0001


Name of Bank: Raiffeisen Bank Zrt.
Postal Address of Bank: Váci út 116-118., 1133 Budapest, Hungary
SWIFT (BIC) Code: UBRTHUHB

Amount of Payment: 950 EUR

*Payment should be made by bank transfer with Details of Charges: „OUR”:
All transfer fees are covered by payer.*

Please indicate the reference number of your manuscript.

Pembayaran Publikasi Artikel: 26 Mei 2025



Berikut ada tiga bukti pembayaran artikel.



Formulir Kiriman Uang

Formulir Kiriman Uang

Pererima/Beneficiary

[illegible]☐ Religion/Personal☐ : Financially/Government ☐ Remittance

Name/Name: Alvin

Alamat/Address : Karesidenan 118

Telephone/Phone:

Kota/City: *Indore*

Negara/Country: Thailand

Bank Penerima/Beneficiary Bank: Bank of China

Kota/City: 5
Negara/Country: Thailand

No. Rek./Acc. No.: 11162/1962-73

Pengirim/Remitter

Resident	Non Resident
1	1
2	2
3	3
4	4
5	5
6	6
7	7
8	8
9	9
10	10
11	11
12	12
13	13
14	14
15	15
16	16
17	17
18	18
19	19
20	20
21	21
22	22
23	23
24	24
25	25
26	26
27	27
28	28
29	29
30	30
31	31
32	32
33	33
34	34
35	35
36	36
37	37
38	38
39	39
40	40
41	41
42	42
43	43
44	44
45	45
46	46
47	47
48	48
49	49
50	50
51	51
52	52
53	53
54	54
55	55
56	56
57	57
58	58
59	59
60	60
61	61
62	62
63	63
64	64
65	65
66	66
67	67
68	68
69	69
70	70
71	71
72	72
73	73
74	74
75	75
76	76
77	77
78	78
79	79
80	80
81	81
82	82
83	83
84	84
85	85
86	86
87	87
88	88
89	89
90	90
91	91
92	92
93	93
94	94
95	95
96	96
97	97
98	98
99	99
100	100

100

☐ Perorangan/Personal☐ Pemerintah/Government ☐ Remittance

Nama/Name: Ali Kurnia Nofal

Nama Alias/*Alias Name*

No. ID: 330 41623 64910661

KIP/SIM/Passpor

Marshall Business: 662 702 8178

Kota/City: XX
Nepara/Country: Nepal

Tujuan Transaksi (*Transaction Purpose*): *Artist Payment*

Berita (Message): *Thank you for your's*

Journal of the
Cattle

Biaya dari Bank koresponden dibebankan ke rekening/

☐ Correspondent bank charges are for account of:

☐ Informing/Convincing ☐ Persuading/Remitter ☐ Sharing

Sah jika ada cetakan data komputer atau tanda

- Transaksi oleh bukan penduduk di atas USD 10.000 atau ekuivalen wajib mengisi Form LDD1/ *Transactional by non-resident amounting over US\$ 10,000 or the equivalent*

Transactions by non-resident amounting over US\$ 10,000 or its equivalent must fill in LBD Form

REFERENCE : S10YGY00041725 /



NO. TRX. : 84976 901213 96963 TRAN 26/05/2025 11:11:35
NO. RPK. : 000000723567241 Sdr ART KUSUMA WARDANA
JMLAH : TDR 35,000 - 1568
004 - YOGYAKARTA

NO. TRX. : 84976 901213 96963 TRAN 26/05/2025 11:11:35
NO. RPK. : 004360420801001 PENDAPATAN PROPTIST KTI
JMLAH : TDR 35,000 1568
004 - YOGYAKARTA

NO. TRX. : 84976 901213 96963 TRAN 26/05/2025 11:11:35
NO. RPK. : 000000723567241 Sdr ART KUSUMA WARDANA
JMLAH : TDR 17,785,900 - 1568
004 - YOGYAKARTA

NO. TRX. : 84976 901213 96963 TRAN 26/05/2025 11:11:35
NO. RPK. : 004360420801001 KTI YAKIR KUB 11:50:44
JMLAH : TDR 950 1568
004 - YOGYAKARTA

NO. TRX. : 84976 901213 96963 TRAN 26/05/2025 11:50:44
NO. RPK. : 000000723567241 KUSUMA WARDANA 11:50:44
JMLAH : TDR 561.660,00 - TRANSFER KR
4 - YOGYAKARTA

NO. TRX. : 84976 142755 021031 BEMI 26/05/2025 11:20:44
NO. RPK. : 4333482010001 Pendapatan Restitusi Biaya ORK
JMLAH : RTR 30,00 NPR MTSG. DR
4 - YOGYAKARTA



Biaya dari Bank koresponden dibebankan ke rekening/
Correspondent bank charges are for account of :
☐ Penerima/Beneficiary ☐ Pengirim/Remitter ☐ Sharing

Pejabat Bank/Bank Officer

Settle ada cetakan data komputer atau tanda tangan yang bermaterai. The applicant form will be void if there is a computerized validation or the authorized signature.
Tersedia oleh Bank Indonesia di atas USD 10.000 atau ekuivalen wajib mengisi Form LDD. Transaction by non-resident amounting over US \$ 10,000 or its equivalent must fill in the DD Form.
Pemohon/Applicant



PT. BANK NEGARA INDONESIA (Persero), Tbk
CABANG : YOGYAKARTA

IBOC - Maintenance (S10)

Teller ID : 84976
Date : 26/05/2025
Time : 11:18:06

Sender's Reference:
:20:S10YGY00041725
Bank Operation Code:
:23B:CRED
Value Date/Currency/Interbank Settled Amount:
:32A:250526EUR950,
Ordering Customer:
:50K:/0000000723567241
SDR ARI KUSUMA WARDANA
BINTANG FAMILY RESIDENCE NO D 1
BANGUNJIWO KASIHAN YOGYAKARTA
INDONESIA
Ordering Institution:
:52A:BNINIDJAXXX
Account With Institution:
:57A:UBRTHUHBXXX
Beneficiary Customer:
:59:/HU53120112650143753400200001
ALOKI INSTITUTE LTD
KASSA UTCA 118 1185
BUDAPEST
HUNGARY
Remittance Information:
:70:ARTICLE PAYMENT
PAYMENT FOR PAIMANS JOURNAL ARTICLE
Details Of Charges:
:71A:OUR





Correspondent bank charges are for account of:
☐ Penerima/Beneficiary
☐ Pengirim/Remitter


☐ Sharing

Pejabat Bank/Bank Officer


Sila isi sah sekiranya data komputer atau tanda tangan yang berlawanan/If there is a computerized validation or the authorized signature
Tanda tangan pejabat bank atau tanda tangan yang berlawanan/If there is a computerized validation or the authorized signature
KANTOR CABANG YOGYAKARTA
Pemohon/Applicant
Lampiran 3 u/ Nasabah

Galley proof for manuscript: 2 Juni 2025





Bukti submit Final Correction: 2 Juni 2025



RESPONSE OF RICE GROWTH AND YIELD UNDER DIFFERENT TIMING OF NPK-BASE FERTILIZERS APPLICATION IN THE SALIBU SYSTEM

PAIMAN* – RAKASIWI, N. – DELZA – MUADIM, N. – SRIWATI, E. – WIBOWO, W. T. – LAILI, L. N. – MUSTAKIM – KUSUMASTUTI, C. T. – ARDIYANTA

Agrotechnology Department, Agriculture Faculty, Universitas PGRI Yogyakarta, Yogyakarta 55182, Indonesia

(e-mail/phone: paiman@upy.ac.id/+62-821-3439-1616; riskiraka46@gmail.com/+62-821-8035-8795; delzawess@gmail.com/+62-813-9970-6506; nurmuadim@gmail.com/+62-822-9880-0503; endangsriwati949@gmail.com/+62-812-3751-0996; wahyutriwibowo281201@gmail.com/+62-858-7715-2262; latifahlaili8@gmail.com/+62-882-1645-4912; mustakimagro@gmail.com/+62-812-4685-7797; kusumastuti@upy.ac.id/+62-821-3439-1664; ir.ardiyanta@upy.ac.id/+62-812-275-7813)

**Corresponding author*

e-mail: paiman@upy.ac.id; phone: +62-821-3439-1616; fax: +62-274-376-808

(Received ; accepted)

Abstract. In the Salibu system (SS), rice plants have a shorter growth cycle than their parent plants. Applying NPK-base fertilizer can promote optimal vegetative growth and yield. This study aimed to determine the optimal timing of NPK-base fertilizer application to enhance rice yield in the SS. The experiment was arranged in a randomized complete block design (RCBD) with three replications, where each replication acted as a block. The timing of NPK-base fertilizers application included four treatments: no application (control), at harvest, stem cuttings (seven days after harvest, or 7 DAH), and seven days after stem cuttings (7 DAC). The results indicated that the application of NPK-base fertilizers could significantly increase the shoot dry weight clump⁻¹ and 1000-filled grains. The Application of NPK-base fertilizer at stem cuttings enhanced plant height at 28 DAC, with no significant difference compared to harvest. Furthermore, this treatment resulted in a higher grains dry weight clump⁻¹ (15.3 g) than the control (13.2 g), and was not significantly different from the treatments at harvest (13.9 g) or 7 DAC (14.6 g). No significant differences were observed among the treatments at harvest, 7 DAC, and the control. The conclusion shows that applying NPK-base fertilizer at stem cuttings of the parent plants optimally enhances the rice yield in the SS. Future research should recommend determining the optimal dosage of NPK-base fertilizer to maximize rice growth and yield.

Keywords: *harvest, main stem, parent plant, ratoon, stem cutting*

Introduction

As Indonesia's population grows, the demand for food, particularly rice, is increasing. Various measures have been implemented to achieve national rice self-sufficiency, including adapting the SS. The term of Salibu is derived from the Indonesian phrase "Salin ibu", which translates to "change of the parent plants" in English.

The ratoon rice technique is a rice cultivation method that utilizes the rice stems remaining after harvest (ratoon) to regrow and produce the next rice crop. After the main harvest, the remaining rice stems are left to regenerate, and new roots and shoots develop to make rice in a

shorter time frame. In the SS, after the main rice harvest, the remaining stems are not regenerated as in the conventional ratoon technique but also undergo special management practices such as cutting the main stems, fertilization, weed control, and others to ensure higher and more stable yields.

The SS features a shorter life cycle than parent plants, enabling increased productivity by applying NPK-base fertilizer. Furthermore, using NPK-base fertilizer can accelerate the growth of rice shoots during the early stages, thereby enhancing the potential for higher yields.

The SS is a rice cultivation method based on local wisdom, originally implemented in West Sumatra, Indonesia. The technology has been tested in various regions across Indonesia. Although farmers have not widely adopted it due to perceptions of lower yields, this technology can enhance rice harvests and improve land-use efficiency with proper management. According to Abdurachman et al. (2015), the SS offers advantages such as faster harvest times, more efficient water usage, and reduced production costs by up to 45% compared to the transplanting method. This method reduces land preparation, seedling production, and planting expenses. Therefore, the SS has the potential to improve the welfare of farmers.

The success of the SS technology is influenced by various factors, one of which is the timing of NPK-base fertilizer application. The NPK fertilizer plays a crucial role in supporting the growth of the new shoots after the parent plants are cut. Applying the fertilizer can provide the essential nutrients the new shoots need and promotes their development. Since the SS has a shorter harvest cycle than its parent plants, it is important to determine the optimal fertilizer application timing to accelerate the growth of the new shoots. However, research on the timing of NPK-base fertilizer in the SS remains limited. Therefore, further studies are necessary to identify the optimal timing for fertilizer application in the SS.

In Indonesia, rice cultivation in irrigated land typically occurs one to three times annually, depending on the irrigation system employed. In rainfed fields, rice can be planted only once a year, while fully irrigated fields can support up to three planting cycles each year. Each planting season involves various activities, including land preparation, seedling production, planting, plant maintenance, and harvesting. Conventional rice cultivation often incurs relatively high costs, lowering profit margins. In contrast, the SS offers greater advantages due to its land and resource utilization efficiency.

The SS can enhance rice production and land productivity annually. This method utilizes the shoots that emerge from the basal stems after cuttings. To maximize the number of tillers, apply the optimal dosage of NPK-base fertilizer. New shoots emerge from the nodes of the remaining stem, followed by the emergence of new roots. Eventually, these shoots grow into independent plants. The formed shoots subsequently develop into new tillers, similar to the rice planting system used in transplanting techniques (Juanda, 2016). The SS can enhance food availability in Indonesia and contribute to food security and resilience. It is more effectively implemented on land that is not always flooded (Jahari and Sinaga, 2019). Rice productivity in tropical regions primarily relies on fertilizers, as soils often lack nitrogen and phosphorus (Hindersah et al., 2022).

Nitrogen (N), phosphorus (P), and potassium (K) are applied in large quantities as fertilizers in rice fields, and a deficiency of any of these nutrients can reduce crop yields (Shrestha et al., 2020). Nitrogen plays a crucial role in the synthesis of chlorophyll and protein. Phosphorus is involved in energy storage and transfer, and it is a key element in nucleic acids, coenzymes, nucleotides, phosphoproteins, phospholipids, and phosphate sugars. Meanwhile, potassium supports starch formation, activates enzymes, and acts as a catalyst in storing photosynthetic products.

Nitrogen is an essential element that forms plants' amino acids, proteins (enzymes), nucleic acids, and chlorophyll. A nitrogen deficiency in rice cultivation leads to reduced plant height, lower tiller formation, decreased leaf area index, and diminished photosynthesis rates (Bijay-Singh and Singh, 2017). Nitrogen plays a crucial role in promoting growth, enhancing yields, and improving grain quality by improving the number of tillers, expanding leaf area, and facilitating protein synthesis. Plants absorb nitrogen through nitrate ions (NO_3^-) and ammonium ions (NH_4^+) (Kaya, 2013). A lack of nitrogen can limit rice productivity, while excessive nitrogen can result in agronomic and economic losses (Hussain et al., 2023). The yield **increases** when nitrogen was applied at $202.15 \text{ kg ha}^{-1}$. The fertilization frequency **is** four times (He et al., 2024). Optimal nitrogen fertilization can effectively compensate for the negative impacts of N fertilizer, enhancing grain quality and nitrogen use efficiency (NUE) while sustaining high yields (Tian-Yang et al., 2022).

Phosphorus (P) is essential in photosynthesis, respiration, energy transfer, storage, cell division, and enlargement. Plants absorb phosphorus in the form of primary orthophosphate ions, H_2PO_4^- (Irwanto, 2014). Additionally, phosphorus is a key component of high-energy compounds such as adenosine triphosphate (ATP) and **is involved in the** synthesizes phospholipids, nucleotides, and glycoproteins. Phosphorus deficiency can hinder growth and yield in rice, which is evident through reduced tiller numbers and dry matter accumulation, along with symptoms such as stunted growth and upright leaves that exhibit a dark green color (Bijay-Singh and Singh, 2017).

Potassium (K) activates various enzymes in plant metabolic processes and is absorbed as potassium ions (K^+) (Irwanto, 2014). This nutrient is essential for enhancing root growth, promoting plant vigor, and increasing resistance to pests and diseases. Symptoms of potassium deficiency in rice include stunted growth with fewer tillers, dark green upper leaves, and chlorosis in the areas between leaf veins (Bijay-Singh and Singh, 2017). Although chemical fertilizers can improve rice yield, the response highly depends on optimal fertilization management in paddy fields (Yang et al., 2021). Using a combination of $135\text{--}210 \text{ kg ha}^{-1}$ N and $115\text{--}137 \text{ kg ha}^{-1}$ K **can** achieve relatively higher yield and improve quality in rice production (Chen et al., 2024).

The NPK Phonska (15-15-15) fertilizer is complex inorganic NPK with composition of 15% N, 15% P_2O_5 , 15% K_2O , and 10% S. This fertilizer is widely utilized by farmers due to its effectiveness in enhancing rice yields and grain quality. According to Shrestha et al. (2020), nitrogen (N), phosphorus (P), and potassium (K) are applied as fertilizers in large quantities to rice fields. Research conducted by Kurniadie (2002), which used a dosage of 300 kg ha^{-1} NPK Phonska along with 333 kg ha^{-1} ZA, resulted in the highest performance for the IR 64 rice variety, including parameters such as the number of panicle clump $^{-1}$, the number of grain panicle $^{-1}$, the harvest index, the weight of 1000 grains, as well as dry grain yields ($7.50 \text{ tons ha}^{-1}$) and milled rice yields ($6.17 \text{ tons ha}^{-1}$).

Optimized fertilization practices represent a promising management approach for achieving sustainable rice production (Zhuang et al., 2022). The application of NPK Phonska fertilizer at a rate of 300 kg ha^{-1} , combined with 200 kg ha^{-1} urea, significantly increased plant height, the number of tillers, and yield up to $7.448 \text{ tons ha}^{-1}$, with relative agronomic effectiveness (RAE) value of 106.46 (Fidiyawati et al., 2022). Combining NPK fertilizer with urea is essential to meet nitrogen nutrient requirements during various rice growth **stages**. Without urea addition, higher dosages of NPK were needed; specifically, 300 kg ha^{-1} NPK Phonska (15-15-15) required an additional 150 kg ha^{-1} urea (Hartatik and Widowati, 2015). The greatest yield of $5.81 \text{ tons ha}^{-1}$ was achieved with the combined treatment of 250 kg ha^{-1} NPK + 150 kg ha^{-1} urea + 1000 kg ha^{-1} dolomite (Ratmini et al., 2023).

Applying NPK fertilizer can enhance plant height and yield by 3%, although it may reduce the weight of 1000 grains. On the other hand, nitrogen fertilizer significantly contributed to an increase in productive tillers and boosted yield by 6%. The most effective combination for rice fields with high phosphorus levels was 300 kg ha⁻¹ NPK and 100 kg ha⁻¹ urea (Siska and Ismon, 2019). The NPK application also enhanced the absorption of nutrients N, P, and K, and improved grain yield (Shrestha et al., 2020). Additionally, Srinivas (2017) noted a significant increase in the number of tillers and panicles m⁻² with the application of 120-60-60 kg ha⁻¹ NPK.

The application of NPK fertilizer significantly influenced plant height at 35, 45, and 90 days after planting (DAP), the number of tillers at 35 and 45 DAP, and various production parameters such as the number of panicles clump⁻¹ and 1000-filled grains of rice (Waty et al., 2014). However, a dosage of 50 kg ha⁻¹ NPK Phonska did not show significant differences in rice production across various agroecological zones (highlands, midlands, and lowlands) using the SS (Fitri et al., 2019). In the SS, fertilization **was applied** to the parent plants: the first application **consisted** of 40% **of the** recommended dosage at 15-20 DAP and the second consisting of 60% at 30-35 DAP (Abdulrachman et al., 2015).

Based on the background and literature review, NPK fertilizer is more commonly used under the transplanting system in rice cultivation. In contrast, its application as base fertilizer in the SS has not been thoroughly investigated. Therefore, the study aimed to determine the optimal timing of NPK-base fertilizer application to enhance rice yield in the SS.

Materials and methods

Study site

The research was conducted from January to June 2024 in Tirtonirmolo Village, Kasihan District, Bantul Regency, Special Region of Yogyakarta, Indonesia. A map of the study site can be seen in *Figure 1*.




Figure 1. A map showing the study site in Tirtonirmolo Village, Kasihan District, Bantul Regency, Special Region of Yogyakarta, Indonesia

The site was located at an elevation of 70 m above sea level, with an average air temperature of 26.6 °C and humidity of 83%. Kasihan District is 110°20'40" East Longitude and 7°48'42" South Latitude.

Experiment design

The experiment was arranged in an RCBD with three replications, where each replication acted as a block. The timing of NPK-base fertilizer application included four treatments: no application (control), at harvest, at stem cuttings, or seven days after harvest (7 DAH), and at seven days after cutting the stems (7 DAC). This study required 12 treatment plots. The size of each treatment plot can be seen in *Figure 2*.




Figure 2. The size of the treatment plot for experimental

Each treatment plot size was 1.84 m (length) $\times$ 1.84 m (width). Plant spacing within the rows was 0.23 m, with 8 plants per row, making the row length 1.84 m. The spacing between rows was 0.23 m, with a total of 8 rows, resulting in a row width of 1.84 m. The number of rice clumps in each plot was 64. The research flow is shown in *Figure 3*.




Figure 3. The flow diagram in research activity

Research procedures

Parent plants

In the SS, the parent plants cultivated by farmers were utilized. The rice variety used was Inpari 42, which has a harvesting period of 87 DAP. The spacing was set at 23×23 cm. The land used for cultivation was technically irrigated, ensuring a consistent water supply, and consisted of alluvial soil. As the parent plant, harvesting was done manually by cutting the stems (along with the panicle) at a height of 15-20 cm above the soil surface using a sickle.

Salibu system

Then, 7 days after the harvest, the remaining stems above the soil were cut again at a height of 3 cm from the soil surface to stimulate new shoot growth in the SS technology. The new shoots emerged from the remaining cut of parent stems and were carefully maintained. Plant care included irrigation, fertilization, and the management of pests, diseases, and weeds to ensure optimal growth.

Water irrigation was applied to the experimental plots until it reached a height of 3 cm above the soil surface. Once it was evenly distributed across all plots, the water channels were closed. This was done every 7 days until the plants reached 70 DAC.

The application of NPK-base fertilizer occurred according to the treatment schedule, which took place at harvest, at stem cuttings (7 DAH), and 7 DAC. The dosage of NPK-base fertilizer used was 60 g plot^{-1} , equivalent to 150 kg ha^{-1} NPK Phonska, except in the control plots. After applying NPK-base fertilizer, a subsequent fertilizer was recommended with 150 kg ha^{-1} NPK Phonska and 200 kg ha^{-1} urea. It was given in two stages: the first stage (40% of the recommended dosage) was applied at 14 DAC, followed by the second stage (60% of the recommended dosage) at 28 DAC.

Weeding was performed at 12 and 26 DAC to reduce weed competition. Pest and disease management was implemented according to integrated pest management (IPM) principles at 56 and 63 DAC, using Decis 25 EC as the control agent. Rice harvesting occurred at 75 DAC when the rice reached physiological maturity (95% yellowing). With the appropriate maintenance and fertilization practices, the rice yield in the SS is expected to be maximized.

Parameters

The research observations included the growth and yield components from each plot treatment. The plant growth components comprised the percentage of tillering stems (%), plant height (cm), the number of tillers clump⁻¹ (stems), and dry weight of shoots (leaves and stems) clump⁻¹ (g). In addition, the observed yield components included the grains dry weight clump⁻¹ (g), the weight of 1000-filled grains (g), and panicle length (cm).

Statistical analysis

The observation data were analyzed using analysis of variance (ANOVA) at 5% significant level (Gomez and Gomez, 1984) with IBM SPSS Version 23 software. If differences were found between treatment means, further testing was conducted using Duncan's new multiple range test (DMRT) at 5% significant level. The term SD is derived from the abbreviation of standard deviation.

Results

Component of the rice growth

Tillering stems

Observations of the tillering stems were conducted on 64 clumps plot⁻¹ at 14 DAC. The results of the ANOVA (*Appendix 1*) indicated that the timing of NPK-base fertilizer application did not significantly affect the percentage of tillering stems that sprouted. The number of tillering stems is presented in *Table 1*.

Table 1. The number of tillering stems (%) at 14 DAC

The timing of NPK-base fertilizer application			
Control	At harvest	At stem cuttings	At 7 DAC
92.7 a	93.8 a	97.9 a	97.4 a

The values in the same row with different letters are significantly different based on DMRT at 5% significant level.
SD_{tillering stems} = 2.617

Figure 4a shows shoot emergence from the basal stem of the parent at 7 DAC, while *Figure 4b* illustrates the tiller growth at 28 DAC.




Figure 4. Photo of the shoot emergence from the basal stem of the parents at 7 DAC (a) and the tiller growth at 28 DAC (b)

Plant height

Plant height was observed on 10 clumps plot⁻¹ at regular intervals of 14, 28, 42, and 56 DAC. The results of the ANOVA (*Appendix 2*) found that the timing of NPK-base fertilizer application significantly affected the plant height at 28 and 42 DAC, while there was no significance at 14 and 56 DAC. The average plant height is presented in *Table 2*.

Table 2 shows that applying NPK-base fertilizer increased only plant height at the 28 DAC. The application of NPK-base fertilizer at stem cuttings of the parent plants resulted in taller plants compared to applying it at control and 7 DAC, although no different from at harvest. However, at

the 42 DAC observation, it appeared that the application of NPK-base fertilizer at 7 DAC led to a decrease in plant height compared to control.

Table 2. The plant height at 14, 28, 42, and 56 DAC

Time of observations (DAC)	The timing of NPK-base fertilizer application			
	Control	At harvest	At stem cuttings	At 7 DAC
14	33.9 a	32.3 a	42.1 a	36.0 a
28	62.4 b	70.3 ab	73.3 a	63.3 b
42	87.0 a	86.9 a	84.3 ab	80.8 b
56	90.1 a	91.3 a	89.5 a	86.8 a

The values in the same row with different letters are significantly different based on DMRT at 5% significant level. $SD_{14\text{ DAC}} = 4.577$, $SD_{28\text{ DAC}} = 4.322$, $SD_{42\text{ DAC}} = 2.211$, and $SD_{56\text{ DAC}} = 3.681$

Tiller numbers

The number of tillers was also observed on 10 clumps plot⁻¹ at regular intervals of 14, 28, 42, and 56 DAC. The results of the ANOVA (Appendix 3) showed that the timing of NPK-base fertilizer application had no significant effect on the number of tillers clump⁻¹ at 14, 28, 42, and 56 DAC. The average number of tillers is shown in Table 3.

Table 3. The number of tiller clump⁻¹ at 14, 28, 42, and 56 DAC

Time of observations (DAC)	The timing of NPK-base fertilizer application			
	Control	At harvest	At stem cuttings	At 7 DAC
14	7.3 a	6.4 a	7.9 a	6.0 a
28	14.6 a	15.9 a	16.9 a	13.6 a
42	17.5 a	18.2 a	19.0 a	15.2 a
56	16.2 a	15.9 a	16.2 a	13.8 a

The values in the same row with different letters are significantly different based on DMRT at 5% significant level. $SD_{14\text{ DAC}} = 2.804$, $SD_{28\text{ DAC}} = 2.567$, $SD_{42\text{ DAC}} = 3.868$, and $SD_{56\text{ DAC}} = 3.636$

Component of the rice yield

The dry weight of shoots clump⁻¹ (g), grains clump⁻¹ (g), weight of 1000-filled grain (g), and panicle length (cm) were observed on 10 clumps plot⁻¹ at 75 DAC. The results of the ANOVA (Appendix 4) indicated that the timing of NPK-base fertilizer application significantly affected the dry weight of shoots, grains clump⁻¹, and 1000-filled grains, while panicle length showed no significant effect. The average of dry weight of shoots, grains, 1000-filled grains, and panicle length are presented in Table 4.

Table 4. The means of shoot dry weight, grain dry weight, the weight of 1000-filled grain, and panicle length at 75 DAC

Parameters observed	The timing of NPK-base fertilizer application			
	Control	At harvest	At stem cuttings	At 7 DAC
Shoots dry weight clump ⁻¹ (g)	23.1 b	31.3 a	32.2 a	30.1 a
Grains dry weight clump ⁻¹ (g)	13.2 b	13.9 ab	15.3 a	14.6 ab
1000-filled grains (g)	13.3 b	19.0 a	19.3 a	17.3 a

Panicle length (cm)	18.9 a	18.6 a	20.5 a	19.8 a
---------------------	--------	--------	--------	--------

The values in the same row with different letters are significantly different based on DMRT at 5% significant level. $SD_{\text{shoot dry weight}} = 2.778$, $SD_{\text{grains dry weight}} = 0.703$, $SD_{\text{1000-filled grains}} = 1.803$, and $SD_{\text{panicle length}} = 1.682$

Table 4 shows that the application of NPK-base fertilizer significantly increase the shoot dry weight clump⁻¹ (g), grain dry weight clump⁻¹ (g), and 1000-filled grains (g). However, the application of NPK-base fertilizer at stem cuttings led to a significantly increase in grain dry weight clump⁻¹ and were not significantly different from those at harvest and 7 DAC. No significant differences were observed among the treatments at harvest, 7 DAC, and the control.

Discussion

Components of rice growth

The application of NPK-base fertilizer in the SS did not influence the percentage of new shoots emerging from cut stems and tillers number. This might have been because the new shoots primarily relied on the food reserve stored in the parent stems. After cutting, the emerging shoots utilized the resources available in the remaining stem for growth. Therefore, although NPK provided essential nutrients, internal factors such as the energy stored in the parent stem might have been more dominant in determining the success of new shoot emergence.

While NPK contained essential nutrients, the growth of tillers was influenced by various factors, including the genetic traits of the rice variety, soil conditions, and management techniques. In the SS, which emphasizes effective resource use, the number of tillers might have been more influenced by care practices and other cultivation techniques rather than by fertilizer alone. Thus, applying NPK as a fertilizer was insufficient to increase the number of tillers significantly.

In contrast, the application of NPK-base fertilizer significantly impacted plant height at 28 DAC in the SS, although this was not always the case for all plant height observations, particularly at 42 DAC. The delayed application of NPK base fertilizer at 7 DAC actually disrupted plant height growth during the maximum vegetative phase, with inhibition observed at 45 DAC. At 56 DAC, there was a tendency for plant height to be lower than control, although the difference was not significant.

The NPK provided nitrogen, phosphorus, and potassium, which were crucial for vegetative growth. Nitrogen played a role in protein synthesis and leaf growth, directly affecting plant height. Additionally, phosphorus supported better root development, allowing the plant to absorb more nutrients from the soil. With optimal nutrient availability, the plant could grow taller, demonstrating that while NPK did not affect new shoots and tillers numbers, it was key in supporting plant height growth. According to Samira et al. (2012), the application of NPK could increase the plant height of rice.

Components of rice yield

In the early stages of growth, new shoots utilized the food reserves from the remains of the parent stems while forming new roots that replaced the function of the parent plant's roots. New shoots were ready to utilize nutrients available in their environment 10 days after the stem cuttings. The application of NPK-base fertilizer at stem cuttings the parent enhanced the availability of essential nutrients, including nitrogen (N), phosphorus (P), and potassium (K), which were crucial for rice growth in the SS. The new shoot's roots quickly absorbed available nutrients to support their development. This base fertilizer helped provide the essential nutrients needed by the new

shoots that emerged from the stem cuttings. New shoots will appear the day after cutting, although small and without leaves.

The application of NPK-base fertilizer significantly increased shoot dry weight compared to the control. This fertilizer contained essential nutrients such as nitrogen, phosphorus, and potassium, which were crucial for plant growth. Nitrogen played a vital role in chlorophyll formation and leaf development. With the application of NPK, plants can develop denser leaves and wider shoots, enhancing their photosynthetic capacity. As a result, the shoots' dry weight was higher than those that did not receive the fertilizer. Yang et al. (2004) emphasized that the rice root system was vital in absorbing water and nutrients from the soil. According to Wang et al. (2006), root system formation was closely coordinated with shoot growth, where the shoot regulated the root development through the circulation of materials between the two. Root distribution determined the structure of the root system within the soil, affecting the acquisition of soil resources such as nutrients and water.

In addition to increasing shoots dry weight, it also increased grain dry weight clump⁻¹ higher than the control. The nutrients provided by NPK supported the grain-filling process, enabling the plants to produce more and heavier grains. Applying nitrogen and phosphorus was crucial for optimal grain development during the grain-filling phase. Plants receiving NPK-base fertilizer showed a significant increase in the dry weight of grains clump⁻¹, indicating that a good nutrient supply contributed to better harvest yields. The product of photosynthesis in the leaves could be translocated in the seed-filling process if the plants were supplied with this nutrient.

According to Rawat et al. (2022), phosphorus is the most abundant element after nitrogen and is involved in many physiological processes such as photosynthesis and translocation. Sarkar et al. (2021) stated that the capacity for source-sink dynamics and assimilate translocation can play an important role during grain formation. As explained by Wei et al. (2018), rice grain yield depends on the supply assimilates (source) and the grain capacity to utilize them (sink).

The application of NPK resulted in the resulting grains being heavier and of higher quality than the control. Adequate nutrients aided in grain-filling, allowing the grains to fill out properly and achieve greater dry weight. The application of NPK-base fertilizer not only increased the quantity of the harvest but also improved the quality of the rice grains, which was essential for determining the market value of the produce. According to Metwally et al. (2020), the grain quality characteristics can be enhanced by the NPK fertilizer application.

As explain by Pasaribu et al. (2018), the parent plant's ability to generate new shoots depends highly on the carbohydrate content and phytohormones retained in the intercalary meristem tissues of the stubble after harvest. Additionally, increasing the rice yield with the SS can be achieved by improving fertilization, especially nitrogen.

Therefore, applying NPK-base fertilizer at stem cuttings was more effective in increasing grain dry weight clump⁻¹ than using it at harvest or 7 DAC. The application of NPK-base fertilizer at stem cutting provided essential nutrients for the new shoots that emerged from the remaining parts of the parent plants. The fertilizer applied at this time could be directly utilized by the shoots after ten days, as nitrogen, phosphorus, and potassium are crucial for accelerating the formation of new tissues and photosynthesis. In contrast, applying NPK-base fertilizer at harvest or 7 DAC was less effective because the growth of new shoots had not yet started during that period. The availability of nutrients was not optimally utilized, leading to lower dry weight and harvest yields. Therefore, the timing of NPK-base fertilizer application significantly influenced nutrient absorption efficiency and the final yield of the plants. Moreover, the application of NPK-base fertilizer at stem cuttings optimally enhanced the rice yield in the SS.

Conclusions

The results indicated that the application of NPK-base fertilizer could significantly increase the shoot dry weight clump⁻¹ and 1000-filled grains. The application of NPK-base fertilizer could enhance the plant height (cm) at 28 DAC, with no significant difference compared to the time of harvesting. Furthermore, this treatment resulted in a higher grains dry weight clump⁻¹ (15.3 g) compared to the control (13.2 g), and was not significantly different from the treatments at harvest (13.9 g) or 7 DAC (14.6 g). No significant differences were observed among the treatments at harvest, 7 DAC, and the control. The conclusion indicate that applying NPK-base fertilizer at stem cuttings of the parent plants optimally enhances rice yield in the SS. Future research should recommend determining the optimal dosage of NPK-base fertilizer to maximize rice yields.

Acknowledgements. We sincerely thank the Institute for Research and Community Service at Universitas PGRI Yogyakarta for funding and supporting this research. We also wish to thank the Faculty of Agriculture, Universitas PGRI Yogyakarta, for providing the necessary financial support for greenhouse facilities and equipment used in this study.

REFERENCES

- [1] Abdulrachman, S., Suhartatik, E., Erdiman, Susilawati, Zaini, Z., Jamil, A., Mejaya, M. J., Sasmita, P., Abdulah, B., Suwarna, Baliadi, Y., Dhalimi, A., Sujinah, Suharna, Ningrum, E. S. (2015): Panduan teknologi budidaya padi salibu. – Badan Penelitian dan Pengembangan Pertanian, Kementerian Pertanian (in Indonesian).
- [2] Bijay-Singh, Singh, V.K. (2017): Fertilizer Management in Rice. – In: Chauhan, B., Jabran, K., Mahajan, G. (eds.) Rice Production Worldwide. Springer, Cham.
- [3] Chen, G., Duan, Q., Wu, C., He, X., Hu, M., Li, C., Ouyang, Y., Peng, L., Yang, H., Zhang, Q., Jiang, Q., Lan, Y., Li, T. (2024): Optimizing rice yield, quality, and nutrient use efficiency through combined application of nitrogen and potassium. – *Frontiers in Plant Science* 15: 1-16.
- [4] Fidiyawati, E., Latifah, E., Krismawati, A., Sihombing, D., Setyorini, D., Bakar, A., Sugiono (2022): Effectivity of inorganic fertilizer NPK (15-15-6) to growth and yield of lowland rice (*Oryza sativa* L.) on alfisol soil. – In: IOP Conference Series: Earth and Environmental Science. The 4th International Conference on Food and Agriculture 980: 012014.
- [5] Fitri, R., Erdiman, Kusnadi, N., Yamaoka, K. (2019): Salibu technology in Indonesia: an alternative for efficient use of agricultural resources to achieve sustainable food security. – *Paddy and Water Environment* 17: 403-410.
- [6] Gomez, A. G., Gomez, K. A. (1984): Statistical Procedures for Agricultural Research. Second Ed. – John Wiley & Sons, Inc., New York.
- [7] Hartatik, W., Widowati, L. R. (2015): Effect of compound fertilizer NPKS and NPK on rice growth and yield on Inceptisol. – *Jurnal Penelitian Pertanian Tanaman Pangan* 34(3): 175-186 (in Indonesian).
- [8] He, X., Zhu, H., Shi, A., Wang, X. (2024): Optimizing nitrogen fertilizer management enhances rice yield, dry matter, and nitrogen use efficiency. – *Agronomy* 14: 919.
- [9] Hindersah, R., Kalay, A. M., Talahaturuson, A. (2022): Rice yield grown in different fertilizer combination and planting methods: case study in Buru Island, Indonesia. – *Open Agriculture* 7: 871-881.
- [10] Hussain, T., Mulla, D. J., Hussain, N., Qin, R., Tahir, M., Liu, K., Harrison, M. T., Sinutok, S., Duangpan, S. (2023): Optimizing nitrogen fertilization to enhance productivity and profitability of upland rice using CSM–CERES–Rice. – *Plants* 12: 3685.

- [11] Irwanto (2014): Pengaruh pemberian pupuk NPK terhadap pertumbuhan tanaman buah naga di Kecamatan Pemayung, Kabupaten Batanghari, Provinsi Jambi. – Balai Pelatihan Pertanian Jambi (in Indonesian).
- [12] Jahari, M., Sinaga, P. H. (2019): Overcome limited growing season with cultivation ratoon and salibu. – Jurnal Dinamika Pertanian 3: 65-72.
- [13] Juanda, R. (2016): Peningkatan produksi padi melalui potensi dan pengembangan wilayah produksi benih unggul di Provinsi Aceh. – Agrosamudra 3(2): 72-80 (in Indonesian).
- [14] Kaya, E. (2013): Effect of rice straw compost and fertilizer NPK on N-available of soil, N-uptake, growth and yield of paddy (*Oryza sativa* L.). – In: Prosiding FMIPA Universitas Pattimura 2013, pp. 41-47 (in Indonesian).
- [15] Kurniadie, D. (2002): Pengaruh kombinasi dosis pupuk majemuk NPK phonska dan pupuk N terhadap pertumbuhan dan hasil tanaman padi sawah (*Oriza sativa* L.) varietas IR 64. – Jurnal Bionatura 4(3): 137-147 (in Indonesian).
- [16] Metwally, T. F., Mohamed, A. A. E., Sorour, S. G. R., Elsayed, G. A. (2020): Stimulating compounds affect the grain quality characteristics and the nutritional value of rice (*Oryza sativa*). – Applied Ecology and Environmental Research 18(5): 6829-6840.
- [17] Pasaribu, P. O., Triadiati, Anas, I. (2018): Rice ratooning using the salibu system and the system of rice intensification method influenced by physiological traits. – Pertanika 41(2): 637-654.
- [18] Ratmini, N. P. S., Subiksa, I. G. M., Suprihatin, A., Suastika, I. W., Riyanto, D. (2023): Optimization of lowland productivity through fertilization as a food buffer against the impact of climate change. – E3S Web of Conferences, 9th ICCO 2023, 467: 01019.
- [19] Rawat, J., Pandey, N., Saxena, J. (2022): Role of Potassium in Plant Photosynthesis, Transport, Growth, and Yield. – In: Iqbal, N., Umar, S. (eds.) Role of Potassium in Abiotic Stress. Springer Nature, Singapore, pp. 1-14.
- [20] Samira, D., Gani, A., Slavich, P., McLeod, M. (2012): Effect of NPK fertilizer and biochar residue on paddy growth and yield of second planting. – In: 2nd Annual International Conference Syiah Kuala University 2012 & The 8th IMT-GT Uninet Biosciences Conference, Banda Aceh, 22-24 November 2012, 2(1): 22-24.
- [21] Sarkar, S., Monshi, F. I., Uddin, M. R., Tabassum, R., Sarkar, M., Hasan, A. (2021): Source-sink manipulation influences the grain-filling characteristics associated with the grain weight of rice. – Journal of Innovative Agriculture 8: 20-29.
- [22] Shrestha, J., Shah, K. K., Timsina, K. P. (2020): Effects of different fertilizers on growth and productivity of rice (*Oryza sativa* L.): a review. – International Journal of Global Science Research 7(1): 1291-1301.
- [23] Siska, W., Ismon, L. (2019): Pemupukan NPK dan nitrogen pada tanaman padi di lahan sawah berstatus P tinggi di Sumatera Barat. – Jurnal Pengkajian dan Pengembangan Teknologi Pertanian 22(2): 175-184 (in Indonesian).
- [24] Srinivas, M. (2017): Response of rice (*Oryza sativa* L.) to different nutrient levels under machine planted conditions. – In: 5th International Conference on Agriculture, Ecology and Biological Engineering, Thailand, May 2-4, 2017, pp: 1-5.
- [25] Tian-Yang, Z., Zhi-Kang, L., En-Peng, L., Wei-Lu, W., Li-Min, Y., Hao, Z., Li-Jun, L., Zhi-Qin, W., Jun-Fei, G., Jian-Chang, Y. (2022): Optimization of nitrogen fertilization improves rice quality by affecting the structure and physicochemical properties of starch at high yield levels. – Journal of Integrative Agriculture 21(6): 1576-1592.
- [26] Wang, H., Inukai, Y., Yamauchi, A. (2006): Root development and nutrient uptake. – Critical Reviews in Plant Sciences 25(3): 279-301.
- [27] Waty, R., Musyassir, Syamaun, Chairunnas (2014): Effects of NPK fertilization and biochar residual to physical properties of soil, plant growth and result of rice field (*Oryza sativa* L.) second planting season. – Jurnal Manajemen Sumber Daya Lahan 3(1): 383-389.
- [28] Wei, H., Meng, L., Li, X., Dai, Q., Zhang, H., Yin, X. (2018): Sink-source relationship during rice grain filling is associated with grain nitrogen concentration. – Field Crops Research 2015: 23-38.

- [29] Yang, C., Yang, L., Yang, Y., Ouyang, Z. (2004): Rice root growth and nutrient uptake as influenced by organic manure in continuously and alternately flooded paddy soils. – *Agricultural Water Management* 70: 67-81.
- [30] Yang, G., Nabi, F., Sajid, S., Kaleri, A. R., Jakhar, A. M., Cheng, L., Raspor, M., Muhammad, N., Ma, J., Hu, Y. (2021): Response of root development and nutrient uptake of two Chinese cultivars of hybrid rice to nitrogen and phosphorus fertilization in Sichuan Province, China. – *Molecular Biology Reports* 48(12): 8009-8021.
- [31] Zhuang, Y., Ruan, S., Zhang, L., Chen, J., Li, S., Wen, W., Liu, H. (2022): Effects and potential of optimized fertilization practices for rice production in China. – *Agronomy* 42(32): 1-16.

APPENDIX

Appendix 1. ANOVA of tillering stem

Source of variance	Degree of freedom	Sum of squares	Mean square	F calc.	F table 5%
Block	2	37.028	18.514	2.703 ns	5.14
Treatment	3	60.832	20.277	2.960 ns	4.76
Error	6	41.097	6.850		
Total	11	138.957			

ns = not significant at 5%

Appendix 2. ANOVA of plant height

Source of variance	Degree of freedom	Mean square				F table 5%
		14 DAC	28 DAC	42 DAC	56 DAC	
Block	2	9.027 ns	6.754 ns	5.127 ns	4.868 ns	5.14
Treatment	3	55.394 ns	84.675 *	25.636 *	10.640 ns	4.76
Error	6	20.947	18.678	4.934	13.549	
Total	11					

ns = not significant at 5% and * = significance different at 5%

Appendix 3. ANOVA of tiller number

Source of variance	Degree of freedom	Mean square				F table 5%
		14 DAC	28 DAC	42 DAC	56 DAC	
Block	2	9.896 ns	17.453 ns	20.818 ns	12.766 ns	5.14
Treatment	3	2.200 ns	6.444 ns	8.132 ns	4.125 ns	4.76
Error	6	7.861	4.960	10.220	14.141	
Total	11					



ns = not significant at 5%

Appendix 4. ANOVA of shoot dry weight, grain dry weight, 1000-filled grains, and panicle length

Source of variance	Degree of freedom	Mean square				F table 5%
		Shoot dry weight	Grain dry weight	1000-filled grains	Panicle length	
Block	2	13.898 ns	0.259 ns	0.250 ns	0.002 ns	5.14
Treatment	3	51.490 *	2.570 *	22.750 *	2.207 ns	4.76
Error	6	7.719	0.494	3.250	2.829	
Total	11					

ns = not significant at 5% and * = significance different at 5%

OnlineFirst of AEER: 1 Juli 2025





As Indonesia's population grows, the demand for food, particularly rice, is increasing. Various measures have been implemented to achieve national rice self-sufficiency, including adapting the SS. The term of Salibu is derived from the Indonesian phrase "Salin ibu" which translates to "change of the parent plants" in English.