

jurnal agussalim (2)

by Agussalim Masulili

Submission date: 20-Dec-2022 10:57AM (UTC+0700)

Submission ID: 1984843956

File name: 11._787-794.docx (51.91K)

Word count: 5084

Character count: 26364

Role of biochar amendments in improving the properties of acid sulphate soil

AGUSALIM MASULILI*, SUTIKARINI, RINI SURYANI, IDA AYU SUCI, ISMAIL ASTAR, HARDI DOMINIKUS BANCIN AND PAIMAN¹

⁴
Department of Agrotechnology, Faculty of Agriculture
Panca Bhakti University, Pontianak 78113, West Kalimantan, Indonesia
*(e-mail : agusalim@upb.ac.id)

(Received : October 03, 2022/Accepted : November 28, 2022)

ABSTRACT

Acid sulfate soils have potential for agricultural development. However, this soil has various properties problems that can inhibit plant growth. One way to improve soil properties is through the application of biochar amendments. The study aimed at knowing the role of rice husk biochar combined with amendments to *Chromolaena odorata*, rice straw, and rice husk ash to improve the properties of acid sulphate soil. The study was conducted in a greenhouse and arranged in a complete randomized design (CRD) with three replications. Biochar amendments consisted of nine kinds, namely, without amendments (control), *C. odorata* (10 t/ha), rice straw (10 t/ha), rice husk biochar (10 t/ha), rice husk ash (10 t/ha), *C. odorata* (10 t/ha)+rice husk biochar (5 t/ha), rice straw (10 t/ha)+rice husk biochar (5 t/ha), *C. odorata* (10 t/ha)+rice husk ash (5 t/ha), rice straw (10 t/ha)+rice husk ash (5 t/ha). The results showed that the amendment of rice husk biochar combined with amendments to *C. odorata* and rice straw, respectively, had a good effect on improving soil physical properties, namely, total soil pores, reducing soil content weight and soil strength. Its effect on soil chemical properties was that it could increase pH, C-organic, P-available, cation exchange capacity (CEC), lower Al-dd and Fe soluble. The research findings showed that applying biochar and organic amendments to *Chromolaena odorata* and rice straw could potentially improve the properties of acid sulphate soil. In future research, we recommend increasing the carrying capacity of acid sulphate soil against plant growth, and it can be done by improving soil properties through the organic amendments' application of *C. odorata* or straw enriched with rice husk biochar.

Key words : *Chromolaena odorata*, rice husk ash, rice husk biochar, rice straw, soil properties

INTRODUCTION

Acidic sulphate soils are the common name for soils and sediments containing iron sulfide, namely, pyrite (Das and Das, 2015). In

agricultural development, efforts to reclaim acid sulphate soils do not always result in soil productivity since improvements in soil properties after reclamation are not achieved.

So far, efforts have been encouraged to overcome existing obstacles after reclamation, including inorganic fertilizers (urea, TSP and KCl) and liming (dolomite). However, rice crop productivity in this field has not been optimal.

Inorganic fertilizers increase over time

and can even decrease land quality if inputs are high and intensive. Liming can only cope with short-term pH and is often done repeatedly. As a result, high costs are needed, which becomes a burden for farmers and will threaten the sustainability of land productivity. Therefore, it is necessary to manage land based on land amendments that can improve the nature of sustainable soils.

Reclamation can result in changes in soil properties and is related to the potential for pyrite oxidation, affecting the increase in soil packaging and the solubility of ions such as Al, Fe, Mn and sulphate. On the other hand, this soil has a high clay content which affects

²⁶
¹Department of Agrotechnology, Faculty of Agriculture, Universitas PGRI Yogyakarta, Yogyakarta 55182, Indonesia.

the increased weight of the contents. Therefore, the development of acid sulphate soil for agriculture after reclamation requires proper management of soil amendments to achieve the formation of steady cultivated land and sustainable farming. Das and Das (2015) suggested that acid sulphate soil having a high pH could inhibit plant growth. In addition, when this soil is drained, it can affect an increase in sulfuric acid and the release of iron, aluminum, and other heavy metals. Shamshuddin *et al.* (2014) stated that plants grown in acidic sulphate soils could not grow well due to low pH and high Al content.

Biochar is one of the soil amendment materials with the long-term potential to maintain soil fertility (Ogunleye *et al.*, 2019; Jaswal *et al.*, 2022). The results of experts' investigations into the phenomenon of terra preta show that soil fertility is stable in the Amazon basin due to the presence of black carbon, the result of the management of the Amerindian nation 500-2,500 years ago. This carbon black is nothing but biochar immersed in the soil. Murtaza *et al.* (2021) found that 4.0 t/ha cotton biochar improved physical, chemical, soil biological soil fertility properties and peanut productivity in Alfisol. Zhang *et al.* (2022) have proven that biochar provides effects such as liming and providing nutrients that lead to soil improvement. Kocsis *et al.* (2022) stated that biochar had a high porosity and could improve the physico-chemical and biological properties of the soil. In line, Sovova *et al.* (2021) found that biochar had a good influence on improving the soil properties of regosol. Further, Das *et al.* (2021) suggest the application of biochar to the soil provide agronomic benefits, support crop yield improvement, and improve soil quality and health. In addition, biochar has been shown to act as a fertilizer and soil conditioner.

Nutrient retention and higher nutrient availability were found in the soil after biochar addition, associated with higher CEC, surface acreage and direct nutrient addition (Glaser *et al.*, 2002). In line with opinions, Murtaza *et al.* (2021) suggested that biochar was a recalcitrant carbon product of biomass produced through the pyrolysis process. The recalcitrant properties of biochar are distinctive and important to the long-term improvement of soil properties through the effects of their residues that can

persist for a long time in the soil. Beusch (2021), biochar can contribute to soil fertility and crop productivity, and Kalu *et al.* (2021) increase water availability and lower soil pH.

Applying biochar can improve soil properties and long-term crop productivity compared to other organic amendments. In other organic amendments, research results by Suri and Yudono (2020) showed that giving

C. odorata as compost can increase plants' nutrition and nutrient uptake. Agustina *et al.* (2019) argued that using *C. odorata* and rice straw could maintain soil quality. Mahmoud *et al.* (2009) also showed rice straw compost could improve soil quality. The same was discovered by Gewaily (2019), that straw compost can increase potassium availability.

Furthermore, the results of the study by Yin *et al.* (2022) showed the weight of the soil contents decreased, and the water-holding capacity and total porosity increased in the soil with rising rice husk ash. Adding rice husk ash neutralizes soil acidity and increases plant nutrient availability. Research by Lestari and Rachmawa (2020) showed that applying rice husk ash (4 t/ha) could increase rice growth and reduce salinity stress.

Previously, several studies on the effect of biochar singularly have been carried out, including on the P-availability (Zhang *et al.*, 2022), improvement of soil physical properties (Aslam *et al.*, 2014; Yin *et al.*, 2018), chemical properties of the soil (Bista *et al.*, 2019), some soil characteristics (Yadav *et al.*, 2018), soil aggregate stability (Ebido *et al.*, 2021) and to the development of rice roots (Kartika *et al.*, 2021). However, no research has been found on how the role of biochar was given together with other organic amendments in acid sulphate soil. Therefore, it is necessary to study how the effect of biochar is given together with organic amendments to *C. odorata* and rice straw. This research is expected to contribute to improving the nature of acid sulphate soil in West Kalimantan, which has been one of the types of soil widely used as an agricultural activity for rice crop. In connection with this background and literature review above, this study aimed at knowing the role of rice husk biochar combined with amendments to *C. odorata*, rice straw and rice husk ash in improving the properties of acid sulphate soil.

14 MATERIALS AND METHODS

Study Area

This research was conducted from September to November 2021. The research site was carried out at the Laboratory of the Faculty of Agriculture, Universitas Panca Bhakti, Pontianak, West Kalimantan Province, Indonesia. The height of the study site is one m from above sea level, with an average temperature and humidity of 27.6°C and 82.8%, respectively. The research position is located at a latitude of 2°05' N-3° 05'S and longitude of 108°30' E-144°10' W.

51 Experimental Design

The study was conducted in a greenhouse and arranged in a complete randomized design (CRD) with three replications. Biochar amendments consisted of nine kinds, namely, without amendment (control), *C. odorata* (10 t/ha), rice straw (10 t/ha), rice husk biochar (10 t/ha), rice husk ash (10 t/ha), *C. odorata* (10 t/ha)+rice husk biochar (5 t/ha), rice straw (10 t/ha)+rice husk biochar (5 t/ha), *C. odorata* (10 t/ha)+rice husk ash (5 t/ha), and rice straw (10 t/ha)+rice husk ash (5 t/ha). In all, 27 polybags were needed.

Research Procedures

The material in this study used acid sulphate soil taken on the experimental land of the West Kalimantan, Sungai Kakap, Agricultural Technology Assessment Center, and land farmers' work on rice crops. This area included the development of swampland agriculture which had built irrigation canals since 1980 and was currently designated as an Integrated Farmer Business Area. Soil picking was carried out randomly at a soil depth of 0-20 cm, then mixed, air-dried, and cleaned. Then put in polybags was as much as 8 kg.

Biochar was made by taking rice husk from grain harvested from rice plants grown directly in the acid sulphate soil of the Kakap River, West Kalimantan, Indonesia. First, the rice mill was taken the grain to obtain rice husk. Furthermore, the rice husk was placed on the pyrolysis device to be processed into biochar by incomplete combustion (limited air

in a reactor made of Pertamina drums.

Organic amendments to *C. odorata* were taken from around the field, then chopped to a size of 1-2 cm. Rice straw was taken from waste harvested rice grown and grown by farmers in acid sulphate soil and chopped within 1-2 cm. Rice husks were taken from waste from rice milling, then used as ash through burning in open areas to produce grayish-white ash.

Prepared acid sulphate soil introduced into 27 polybags until it reached 8 kg. Further, the dose of soil amendment was calculated based on the soil weight in the polybag area, mixed with the soil in each polybag to a soil depth of 20 cm, after which it was incubated with moisture content until it was closed to the field capacity for 30 days.

At 30 days after incubation, from each study unit, soil sampling of about 50 g was undisturbed to determine the weight of the soil moisture content accident. In addition, a disturbed soil sample of 100 g was taken from each research unit for analysis of chemical properties.

Observation Parameters

The soil bulk density (BD) was determined by the clod method described by Blake & Harke. The total pore was calculated from the soil moisture content (v/v) at the matrix potential of 0 kPa. The available groundwater was estimated by reducing the soil moisture content in the matrix potential by -33 kPa (field capacity) with the soil moisture content in the matrix potential, which was -15 Mpa (wilting point). The soil moisture content at this potential matrix was determined using a pressure plate device. The strength of the soil was measured with a hand penetrometer (Daikie) at a soil depth of 15 cm.

The pH level was measured in a soil solution at a ratio of 1 : 2.5 (with deionized water), using a pH meter (Jenway 3305). Walkley and Black's wet oxidation method was used to determine organic C levels (Soil Survey Laboratory Staff). Total N levels were measured by the Kjeldhal (Bremner and Mulvaney) method. Al₃ and Fe₂ were extracted with 1 M KCl (Barnhisel and Bertsch). CEC was extracted with 1 M NH₄Oac (buffer at pH 7.0), and the concentration of alkaline cations was measured using AAS (Shimadzu), P-available with Bray I.

Statistical Analysis

Statistical analysis was carried out on each observation parameter data to determine the effect of the amendments on the nature of acid sulphate soil. The data obtained were analyzed by analysis of variance (ANOVA) at significant level. In addition, the test of least significant difference (LSD) at 5% significant level was used to determine the average value between treatments.

RESULTS AND DISCUSSION

Effect of Various Soil Amendments on Some Physical Soil Properties

Applying biochar improved the soil's physical properties (Aslam *et al.*, 2014). Applying biochar and organic amendments influenced the changes in some of the observed soil physical properties. The statistical test is shown in Table 1. The soil BD was lower on all amendments granted. For example, in soil fed a single rice husk biochar (10 t/ha), soil BD decreased from 1.27 mg/m³ in the control soil to 1.15 mg/cm³. However, not significantly different from soils was given straw (10 t/ha) and a combination of straw+biochar, *C. odorata*+rice husk ash, and straw+rice husk ash. Furthermore, in the combination of *C. odorata* (10 t/ha)+rice husk biochar (5 t/ha), a BD of 1.13 mg/m³ was obtained and was the lowest soil BD. Still, it was not significantly different from the soil, given a combination of straw (10 t/ha) + rice husk biochar (5 t/ha) produced soil BD of 1.14 mg/m³. These results indicated an

improvement in the physical soil properties if biochar was given together with organic amendments to *C. odorata*.

Applying various biochar amendments and their combinations increased the soil's total pore. The statistical test results are shown in Table 1. There was a significant difference in the total pore of the soil between the various amendments. Control soils had the lowest total pore (44.43%), contrasting with soils that provided various amendments. On soils with rice husk biochar (10 t/ha), the total soil pore increased to 54.21%, significantly different from other single amendments. Furthermore, a combination of *C. odorata* (10 t/ha)+rice husk biochar (5 t/ha) caused the total pore to increase to 56.73% and was the highest total pore.

The application of various amendments, both single and combined, affected the decrease in soil strength as measured as soil penetration resistance (N/cm²) at conditions of pF 0, pF 2 and pF 2.5, where the higher the soil matric suction (pF), the higher the soil penetration resistance. The under saturated conditions (pF 0), the various soil amendments did not offer a noticeable difference. In pF 2 showed that the application of rice husk biochar (10 t/ha) could reduce soil penetration resistance from 310.00 N/cm² (control soil) to 223.33 N/cm², but not significant difference between *C. odorata* (10 t/ha)+rice straw or a combination of straw (10 t/ha)+rice husk biochar (5 t/ha). Furthermore, soils given a combination of *Cromolaena odorata* (10 t/ha)+biochar (5 t/ha) obtained the lowest penetration resistance (220.00 N/cm²), and this treatment was not significantly different

Table 1. Effect of rice husk biochar with soil amendments on some physical properties of acid sulphate soil at one month after incubation.

Soil amendments	BD (mg/m ³)	Total pores (%)	Penetration resistance (N/cm ²)		
			pF 0	pF 2	pF 2.5
Control	1.24e	44.43a	36.67b	310.00e	500.00d
<i>C. odorata</i> (10 t/ha)	1.16c	52.17e	20.00a	230.00abc	403.33bc
Rice straw (10 t/ha)	1.17bc	53.27f	16.67a	243.33bc	393.33b
Rice husk biochar (10 t/ha)	1.15b	54.21g	40.33a	223.33ab	390.00b
Rice husk ash (42 t/ha)	1.19d	47.30b	20.00a	270.00d	403.33bc
<i>C. odorata</i> (10 t/ha)+rice husk ash (5 t/ha)	1.13a	56.73i	10.00a	220.00a	340.00a
Rice straw (10 t/ha)+rice husk biochar (5 t/ha)	1.14ab	55.57h	13.33a	223.00ab	360.00a
<i>C. odorata</i> (10 t/ha)+rice husk ash (5 t/ha)	1.15b	48.90c	16.67a	240.00bc	420.00c
Rice straw (10 t/ha)+rice husk ash (5 t/ha)	1.16bc	50.30d	20.00a	246.67c	403.33bc

Remarks : The average numbers followed by the same letter in the same column show no significant difference in the LSD test at 5% significant level.

from soils have given *C. odorata* (10 t/ha), rice husk biochar (10 t/ha), and a combination of rice straw (10 t/ha)+rice husk biochar (5 t/ha).

Similarly, in pF 2.5, the application of rice husk biochar (10 t/ha) could reduce the value of soil penetration resistance from 500.00 N/cm² (control soil) to 390.00 N/cm², and this treatment not significantly different from soils had given *C. odorata* (10 t/ha), rice straw (10 t/ha), rice husk ash (8 t/ha), and a combination of rice straw (10 t/ha)+rice husk ash (5 t/ha) furthermore, the combination of *C. odorata* (10 t/ha)+rice husk biochar (5 t/ha) achieved the lowest penetration resistance (340.00 N/cm²). Also, it was not significantly different from the combination of rice straw (10 t/ha)+biochar (5 t/ha) (360.00 N/cm²). It indicated that rice husk biochar had the potential to control soil strength. In line with Das *et al.* (2021), applying biochar into the soil could reduce soil strength.

22 Effect of Rice Husk Biochar on Some Soil Chemical Properties

The application of several soil amendments and their combinations could influence some chemical properties of acid sulphate soil. The results of statistical tests in Table 2 show that soil reaction (pH) of the soil increased in all treatments given compared to controls. For example, in the application of rice husk biochar showed an increase in pH from 3.36 (control soil) to 4.40 on soils given biochar (10 t/ha) but did not significantly differ from other single amendments. Furthermore, the combined application of *C. odorata* (10 t/ha)+biochar (5

t/ha) obtained the highest increase in soil pH (4.58), in marked contrast to soils given rice husk ash but not significantly different from single amendments or other combinations.

Applying single and combined soil amendments affected the increase in soil C-organics. The statistical test results in Table 2 show that the control soil had the lowest C-organic (1.94%), in stark contrast to other amended soils, except that the soil had given rice husk ash (10 t/ha) was not significantly different. The soil given a single amendment of rice straw (10 t/ha) obtained 4.58% C-organic, in stark contrast to the control soil and the soil given rice husk ash, but not significantly different from other soil amendments. Similarly, soils given a combination of *C. odorata* (10 t/ha)+biochar husk (5 t/ha) achieved the highest soil C-organic (4.93%) in marked contrast to control soils and soils fed with rice husk ash but did not significantly differ from other soil amendments.

Applying various amendments and combinations to acid sulphate soil generally affected the increase in P-available. The results of statistical tests (Table 2) show that control soils had the lowest P-available content (0.21%), significantly different from soils given various amendments. In soils fed with a rice husk biochar (10 t/ha), the P-available increased to 0.32% . Still, it was not significantly different from other combined or single-amended soils, except with soils that were given rice husk ash with a lower P-available (0.27%). The highest P-available was obtained on soils given a combination of *C. odorata* (10 t/ha)+biochar (5 t/ha) (0.34%), in stark contrast to control soils and those topped

1 Table 2. Effect of rice husk biochar with soil amendments on some chemical properties of acid sulphate soil at one month after incubation.

Soil amendments	17 pH H ₂ O	C-organic (%)	P-available (%)	Al-dd (me/100 g)	Soluble Fe (%)	CEC (me/100 g)
Control	3.36a	1.94a	0.21a	3.84e	3.61d	6.64a
<i>C. odorata</i> (10 t/ha)	4.06bc	4.22b	0.29bc	3.31c	3.28c	7.15ab
Rice straw (10 t/ha)	4.28bc	4.58b	0.30bc	3.42cd	3.34cd	7.32abc
Rice husk biochar (10 t/ha)	4.40bc	4.09b	0.32cd	3.16b	3.10abc	8.03c
Rice husk ash (10 t/ha)	3.98b	2.78a	0.28b	3.51d	3.34cd	7.76bc
<i>C. odorata</i> (10 t/ha)+rice husk biochar (5 t/ha)	4.58c	4.93b	0.34d	3.02a	2.94ab	10.04e
Rice straw (10 t/ha)+rice husk biochar (5 t/ha)	4.48bc	4.75b	0.33d	3.08ab	2.91a	9.67de
<i>C. odorata</i> (10 t/ha)+rice husk ash (5 t/ha)	4.26bc	4.70b	0.31cd	3.37c	3.25c	9.17d
Rice straw (10 t/ha)+rice husk ash (5 t/ha)	4.26bc	4.73b	0.31cd	3.47d	3.22bc	9.20d

7 Remarks : The average numbers followed by the same letter in the same column show no significant difference in the LSD test at 5% significant level.

with rice husk ash but not significantly different from single amendments or other combinations.

The decrease in Al-dd and Fe-dissolve in acid sulphate soil also occurred due to applying various soil amendments and their combinations. Statistical tests (Table 2) showed that the highest Al-dd was obtained on control soil (3.84%), further decreasing and differing markedly from soils given various single and combined amendments. On single-amendment soils of rice husk biochar (10 t/ha), Al-dd decreased to 3.16%, significantly different from other single-amended soils. Furthermore, the soil given the combination of *C. odorata* (10 t/ha)+rice husk biochar (5 t/ha) obtained the lowest Al-dd content (3.02%). It was not significantly different from the soil given a combination of rice straw (10 t/ha)+rice husk biochar (5 t/ha), but it was significantly different from other amended soils. In soluble Fe content, the application of rice husk biochar (10 t/ha) could reduce soluble Fe from 3.61% on control soil to 3.10% and was significantly different from other single amendment. And when the soil was given a combination of rice straw (10 t/ha)+rice husk biochar (5 t/ha) caused the lowest decrease in Al-dd (2.91%). Still, it was not significantly different from soil that was given a combination of *C. odorata* (10 t/ha)+rice husk biochar (5 t/ha).

The CEC of land has also increased due to the application of various land amendments and their combinations. The statistical test shown in Table 2 shows that the control soil had the lowest CEC (6.64 Cmol/kg), not significantly different from soils fed with *C. odorata* and rice straw. On the other hand, on soils provided with rice husk biochar (10 t/ha), the soil CEC increased to 8.03 Cmol/kg, but it was not significantly different from soil given rice straw and rice husk ash.

In line with the results that have been put forward, Yamato *et al.* (2006) found that the application of biochar from *Acacia nigium* bark into the soil in Sumatra led to changes in soil chemical properties through increased pH, N-total and P-available, CEC and lowered interchangeable Al. Liang *et al.* (2006) found that two mechanisms could create a larger CEC. First, a higher charged density per unit of surface acreage means a higher degree of oxidation of soil organic matter. Second, the

presence of a higher surface area for cation absorption or the combined effect of both. Glaser *et al.* (2002) posit that oxidation of aromatic C and carboxyl formation were the main reasons for the high CEC. Liang *et al.* (2006) also found that the CEC per unit C was higher, and the load density was higher in carbon-black-rich Anthrosol compared to soils that were poor in carbon black. In addition, Anthrosol showed a higher surface acreage due to its higher carbon black concentration.

From the results, the great potential of rice husk biochar against improving soil properties increased when given together with the biomass *C. odorata*. It could happen because *C. odorata* contained higher humate and fulvic organic acids, so that it could encourage the complex mechanisms of organo-mineral organs and soil aggregation. In addition, *C. odorata* can highly suppress Al and Fe, thereby increasing the P availability in the soil due to the presence of organic acids released during decomposition. Research results by Suri and Yudono (2020) also found that using *C. odorata* compost could improve soil quality and nutrient uptake of lettuce plants.

C. odorata can contribute to the care of soil C-organic and physical properties due to a reasonably high polyphenol content that could inhibit the decomposition and mineralization of N organic matter, even though it had a low C/N ratio.

CONCLUSION

In conclusion, the amendment of rice husk biochar combined with amendments to *C. odorata* and rice straw improved soil physical properties, namely, total soil pores, reducing soil content weight and strength. Its impact on soil chemical properties was that it could increase pH, C-organic, P-available, CEC, lower Al-dd and Fe soluble. Therefore, applying biochar and organic amendments to *C. odorata* and rice straw could improve the properties of acid sulphate soil. Consequently, we recommend increasing the carrying capacity of acid sulphate soil against plant growth. It can be done by enhancing soil properties through the organic amendments application of *C. odorata* or straw enriched with rice husk biochar.

ACKNOWLEDGEMENTS

6

The authors thank the Institute for Research and Community Service, Universitas Panca Bhakti, which has given support for research funds. In addition, they thank the Faculty of Agriculture, Universitas Panca Bhakti, which has provided loans for facilities in the form of equipment for research.

REFERENCES

- Agustina, R., Jumadi, R., Firmani, U. and Abdul Faisal, R. H. (2019). Effects of decomposition rate of *Chromolaena odorata* and straw rice in fresh and compost form to the growth and yield of rice. *IOP Conf. Ser. Earth Environ. Sci.* **250** : 1-5.
- Aslam, Z., Khalid, M. and Aon, M. (2014). Impact of biochar on soil physical properties. *Sch. J. Agric. Sci.* **4** : 280-84.
- Beusch, C. (2021). Biochar as a soil ameliorant : How biochar properties benefit soil fertility-A review. *J. Geosci. Environ. Prot.* **9** : 28-46.
- Bista, P., Ghimire, R., Machado, S. and Pritchett, L. (2019). Biochar effects on soil properties and wheat biomass vary with fertility management. *Agronomy* **9** : 1-10.
- Das, S. K. and Das, S. K. (2015). Acid sulphate soil : Management strategy for soil health and productivity. *Pop. Kheti* **3** : 3-8.
- Das, S., Mohanty, S., Sahu, G., Rana, M. and Pilli, K. (2021). Biochar : A sustainable approach for improving soil health and environment. In : *Soil Erosion-Current Challenges and Future Perspectives in a Changing World*. pp. 1-18.
- Ebido, N. E., Edeh, I. G., Unagwu, B. O., Nnadi, A. L., Ozongwu, O. V., Obalum, S. E. and Igwe, C. A. (2021). Rice-husk biochar effects on organic carbon, aggregate stability and nitrogen fertility of coarse-textured Ultisols evaluated using *Celosia argentea* growth. *Sains. Tanah.* **18** : 177-87.
- Gewaily, E. (2019). Impact of compost rice straw and rice straw as organic fertilizer with potassium treatments on yield and some grain quality of Giza 179 rice variety. *J. Plant Prod.* **10** : 143-51.
- Glaser, B., Lehmann, J. and Zech, W. (2002). Ameliorating physical and chemical properties of highly weathered soils in the tropics with charcoal-A review. *Biol. Fertil. Soils* **35** : 219-30.
- Jaswal, A., Mehta, C. M. and Singh, A. (2022). Probing the impact of biochar combined with organic and inorganic amendments on soil carbon pools of rice (*Oryza sativa*)-wheat (*Triticum aestivum*) cropping system. *Res. Crop* **23** : 508-15.
- Kalu, S., Simojoki, A., Karhu, K. and Tammeorg, P. (2021). Long-term effects of softwood biochar on soil physical properties, greenhouse gas emissions and crop nutrient uptake in two contrasting boreal soils. *Agric. Ecosyst. Environ.* **316** : doi.org/10.1016/j.agee.2021.107454.
- Kartika, K., Sakagami, J. I., Lakitan, B., Yabuta, S., Akagi, I., Widuri, L. I., Siaga, E., Iwanaga, H. and Nurrahma, A. H. I. (2021). Rice husk biochar effects on improving soil properties and root development in rice (*Oryza glaberrima* Steud.) exposed to drought stress during early reproductive stage. *AIMS Agric. Food* **6** : 737-51.
- Kocsis, T., Ringer, M. and Biró, B. (2022). Characteristics and applications of biochar in soil-plant systems : A short review of benefits and potential drawbacks. *Appl. Sci.* **12** : 1-16.
- Lestari, M. F. and Rachmawati, D. (2020). The effect of rice husk ash on growth of rice (*Oryza sativa* (L.) 'Sembada Merah') on salinity stress. *AIP Conf. Proc.* **2260** : 1-6.
- Liang, B., Lehmann, J., Solomon, D., Kinyangi, J., Grossman, J., O'Neill, B., Skjemstad, J. O., Thies, J., Luizão, F. J., Petersen, J. and Neves, E. G. (2006). Black carbon increases cation exchange capacity in soils. *Soil Sci. Soc. Am. J.* **70** : 1719-30.
- Mahmoud, E., Ibrahim, M., Robin, P., Akkal-Corfini, N. and El-Saka, M. (2009). Rice straw composting and its effect on soil properties. *Compost Sci. Util.* **17** : 146-50.
- Murtaza, G., Ahmed, Z., Usman, M., Tariq, W., Ullah, Z., Shareef, M., Iqbal, H., Waqas, M., Tariq, A., Wu, Y., Zhang, Z. and Ditta, A. (2021). Biochar induced modifications in soil properties and its impacts on crop growth and production. *J. Plant Nutr.* **44** : 1677-91.
- Ogunleye, K. S., Okeke, O. S., Umego, M. O. and Ogbonnaya, O. U. (2019). Extraction behaviour of cadmium in biochar amended soils. *Crop Res.* **54** : 59-64.
- Shamshuddin, J., Elisa Azura, A., Shazana, M. A. R. S., Fauziah, C. I., Panhwar, Q. A. and Naher, U. A. (2014). Properties and management of acid sulfate soils in South-east Asia for sustainable cultivation of rice, oil palm and cocoa. *Adv. Agron.* **124** : 91-142.
- Sovova, S., Enev, V., Smilek, J., Kubikova, L., Trudicova, M., Hajzler, J. and Kalina, M. (2021). The effect of biochar application on soil properties and growth of the model plant *Zea mays*. *Ecocycles* **7** : 46-54.

- Suri, A. M. and Yudono, P. (2020). Effects of *Chromolaena odorata* compost on soil and nutrient uptake of lettuce (*Lactuca sativa*). *Planta Trop. J. Agro. Sci.* **8** : 33-38.
- Yadav, N. K., Kumar, V., Sharma, K. R., Choudhary, R. S., Butter, T. S., Singh, G., Kumar, M. and Kumar, R. (2018). Biochar and their impacts on soil properties and crop productivity : A review. *J. Pharmacogn. Phytochem.* **7** : 49-54.
- Yamato, M., Okimori, Y., Wibowo, I. F., Anshori, S. and Ogawa, M. (2006). Effects of the application of charred bark of *Acacia mangium* on the yield of maize, cowpea and peanut, and soil chemical properties in South Sumatra, Indonesia. *Soil Sci. Plant Nutr.* **52** : 489-95.
- Yin, M., Li, X., Liu, Q. and Tang, F. (2022). Rice husk ash addition to acid red soil improves the soil property and cotton seedling growth. *Sci. Rep.* **12** : 1-9.
- Yin, X., Huang, M. and Zou, Y. (2018). Potential use of biochar in raising machine-transplanted rice seedlings. *Farm. Manage.* **3** : 87-92.
- Zhang, Y., Chen, H., Xiang, J., Xiong, J., Wang, Y., Wang, Z. and Zhang, Y. (2022). Effect of rice-straw biochar application on the acquisition of rhizosphere phosphorus in acidified paddy soil. *Agronomy* **12** : 1-13.

jurnal agussalim (2)

ORIGINALITY REPORT

25%

SIMILARITY INDEX

20%

INTERNET SOURCES

23%

PUBLICATIONS

4%

STUDENT PAPERS

PRIMARY SOURCES

1	docplayer.net Internet Source	3%
2	repository.upy.ac.id Internet Source	2%
3	Submitted to Nasarawa State University Student Paper	2%
4	S Andayani, E S Hayat, R Hayati, H M Kurniawan, E Noerhartati, A T P Irianti. "Compost of oil palm empty fruit bunches with coastal mud and rice husk biochar to improve the acid sulphate soil fertility", Journal of Physics: Conference Series, 2019 Publication	2%
5	opus4.kobv.de Internet Source	1%
6	Submitted to Universitas Wijaya Kusuma Surabaya Student Paper	1%
7	N Arfarita. "The effect of VP3 Biofertilizer and its carrying materials on the germination of	1%

six plants", IOP Conference Series: Earth and Environmental Science, 2021

Publication

8	Rr Ernawati, Meidaliyantisyah, J Barus. "The response of Inpago 8 rice to the provision of compost and Biochar in dryland", IOP Conference Series: Earth and Environmental Science, 2020 Publication	1 %
9	link.springer.com Internet Source	1 %
10	www.iiste.org Internet Source	1 %
11	Agusalim Masulili, Wani Hadi Utomo, Erwin I Wisnubroto. "Growing Rice (<i>Oriza sativa</i> L.) in the Sulphate Acid Soils of West Kalimantan, Indonesia", International Journal of Agricultural Research, 2016 Publication	<1 %
12	JAS, Editor. "Journal of Agricultural Science, Vol. 2, No. 1, March 2010, all in two files, part I", Journal of Agricultural Science, 2010. Publication	<1 %
13	www.cropresearch.org Internet Source	<1 %
14	PAIMAN, ARDIYANTA, SUBENI, KHARISUN, S.F. YUSOFF. "EFFECT OF WATERLOGGING ON	<1 %

WEED SEED GERMINATION AND GROWTH IN LOWLAND RICE", Applied Ecology and Environmental Research, 2022

Publication

15

Titiek Islami, Erwin Ismu Wisnubroto, Waego Hadi Nugroho. "Biochar Derived from Sugarcane Industry Waste Increasing Productivity of Degraded Land", International Journal of Soil Science, 2016

Publication

<1 %

16

G. C. Paul, M. A. Mannan. "An integrated nutrient management approach to improve sugar productivity", Sugar Tech, 2018

Publication

<1 %

17

Y Lestari, W Annisa. "Phosphate-solubilizing bacterial activity and their effect to rice growth in acid sulphate soil", IOP Conference Series: Earth and Environmental Science, 2019

Publication

<1 %

18

docobook.com

Internet Source

<1 %

19

www.frontiersin.org

Internet Source

<1 %

20

www.ncbi.nlm.nih.gov

Internet Source

<1 %

21

Ghulam Murtaza, Zeeshan Ahmed, Muhammad Usman, Waseem Tariq et al.

<1 %

"Biochar induced modifications in soil properties and its impacts on crop growth and production", Journal of Plant Nutrition, 2021

Publication

22

Sanat Kumar Dwibedi, Vimal Chandra Pandey, Donakonda Divyasree, Omesh Bajpai.

"Biochar - based land development", Land Degradation & Development, 2022

Publication

23

Tingqiang Li, Hong Jiang, Xiaoe Yang, Zhenli He. "Competitive sorption and desorption of cadmium and lead in paddy soils of eastern China", Environmental Earth Sciences, 2012

Publication

24

www.nature.com

Internet Source

25

C N Ichsan, Bakhtiar, Sabaruddin, Efendi. "Morpho-agronomic traits and balance of sink and source of rice planted on upland rainfed", IOP Conference Series: Earth and Environmental Science, 2021

Publication

26

www.aloki.hu

Internet Source

27

Diogo José Horst, Jhon Jairo Ramírez Behainne, Pedro Paulo de Andrade Junior.

<1 %

<1 %

<1 %

<1 %

<1 %

<1 %

"Lignin as alternative fuel: an estimate of the thermal energy generation potential from brazilian crops", Environmental Progress & Sustainable Energy, 2017

Publication

28

Mohammad Babla, Utsab Katwal, Miing-Tiem Yong, Soheil Jahandari, Maroun Rahme, Zhong-Hua Chen, Zhong Tao. "Value-added products as soil conditioners for sustainable agriculture", Resources, Conservation and Recycling, 2022

Publication

<1 %

29

Qurban Ali Panhwar, Umme Aminun Naher, Shamshuddin Jusop, Radziah Othman, Md Abdul Latif, Mohd Razi Ismail. "Biochemical and Molecular Characterization of Potential Phosphate-Solubilizing Bacteria in Acid Sulfate Soils and Their Beneficial Effects on Rice Growth", PLoS ONE, 2014

Publication

<1 %

30

muir.massey.ac.nz

Internet Source

<1 %

31

psasir.upm.edu.my

Internet Source

<1 %

32

www.uea.ac.uk

Internet Source

<1 %

33 Masahide Yamato, Yasuyuki Okimori, Irhas Fredy Wibowo, Saifuddin Anshori, Makoto Ogawa. " Effects of the application of charred bark of on the yield of maize, cowpea and peanut, and soil chemical properties in South Sumatra, Indonesia ", Soil Science and Plant Nutrition, 2006
Publication

34 academicjournals.org
Internet Source

35 journal.umy.ac.id
Internet Source

36 krishikosh.egranth.ac.in
Internet Source

37 staff.najah.edu
Internet Source

38 ui.adsabs.harvard.edu
Internet Source

39 www.omicsonline.org
Internet Source

40 British Food Journal, Volume 111, Issue 4 (2009-04-19)
Publication

41 Honghong Lyu, Siyu Xu, Yaxuan Liu, Wenzhu Zhang, Qiuzhi Duan, Minjie Zhu, Jingchun Tang. "Effect of biochar on the emission of

greenhouse gas in farmland", Elsevier BV,
2022

Publication

42

Israel Dzomeku, Osman Illiasu, Isaac Amegbor. "Effects of Biochar, Rice Husk and Rice Straw on Productivity of Maize (*Zea mays* L.) and Sustainable Soil Fertility Restoration", *Journal of Experimental Agriculture International*, 2018

Publication

<1 %

43

Recycling of Biomass Ashes, 2011.

Publication

<1 %

44

Smilja Lambert, Hussin bin Purung, Syawaluddin, Peter McMahon. "Growth and flowering of young cocoa plants is promoted by organic and nitrate-based fertiliser amendments", *Experimental Agriculture*, 2020

Publication

<1 %

45

T. I. Moiseenko, M. M. Bazova, M. I. Dinu, N. A. Gashkina, L. P. Kudryavtseva. "Changes in the Geochemistry of Land Waters at Climate Warming and a Decrease in Acid Deposition: Recovery of the Lakes or Their Evolution?", *Geochemistry International*, 2022

Publication

<1 %

46

Veronika Jílková, Gerrit Angst. "Biochar and compost amendments to a coarse-textured

<1 %

temperate agricultural soil lead to nutrient leaching", Applied Soil Ecology, 2022

Publication

47

depositonce.tu-berlin.de

Internet Source

<1 %

48

etd.ohiolink.edu

Internet Source

<1 %

49

mdpi-res.com

Internet Source

<1 %

50

"Engineered Biochar", Springer Science and Business Media LLC, 2022

Publication

<1 %

51

"Halt soil salinization, boost soil productivity", Food and Agriculture Organization of the United Nations (FAO), 2022

Publication

<1 %

52

"Sustainable Intensification for Agroecosystem Services and Management", Springer Science and Business Media LLC, 2021

Publication

<1 %

53

Jyrki Hytönen. "Effect of peat ash fertilization on the nutrient status and biomass production of short-rotation willow on cut-away peatland area", Biomass and Bioenergy, 1998

Publication

<1 %

Exclude quotes On

Exclude matches < 1 words

Exclude bibliography On