

## RIWAYAT KORESPONDENSI

In Journal: **Research on Crops (Q3), ISSN:0972-3226**  
Publisher: Gaurav Society of Agricultural Research Information Centre

<https://www.scopus.com/sourceid/19900191751>

The screenshot shows the Scopus Source details page for 'Research on Crops'. The page includes the following information:


- Research on Crops**
- Scopus coverage years: from 2008 to 2022
- Publisher: Gaurav Society of Agricultural Research Information Centre
- ISSN: 0972-3226
- Subject area: Agricultural and Biological Sciences: Agronomy and Crop Science, Agricultural and Biological Sciences: Soil Science
- Source type: Journal
- Metrics: CiteScore 2021 (1.7), SJR 2021 (0.272), SNIP 2021 (1.056)
- Actions: View all documents, Set document alert, Save to source list
- Navigation: CiteScore, CiteScore rank & trend, Scopus content coverage
- Table: Year, Documents published, Actions

<https://www.scimagojr.com/journalsearch.php?q=19900191751&tip=sid&clean=0>


The screenshot shows the ScimagoJR journal search results for 'Research on Crops'. The page includes the following information:

- Research on Crops**
- COUNTRY:** India
- SUBJECT AREA AND CATEGORY:** Agricultural and Biological Sciences, Agronomy and Crop Science, Soil Science
- PUBLISHER:** Gaurav Society of Agricultural Research Information Centre
- H-INDEX:** 15
- PUBLICATION TYPE:** Journals
- ISSN:** 09723226


Link Cover Web : [https://indianjournals.com/ijor.aspx?target=ijor:rcr&type=for\\_authors](https://indianjournals.com/ijor.aspx?target=ijor:rcr&type=for_authors)




3 Oktober 2022 Submit Manuscript



**4 Oktober 2022 Penerimaan Manuscript oleh Jurnal Research on crops**



**20 Oktober 2022\_Saran Perbaikan Manuscript dan  
Tgl 21 Oktober 2022\_Respon untuk menerima saran perbaikan**



# RESEARCH ON CROPS

*Gaurav Publications (Regd.), Hisar, India*

Ref. No. GP-ROC/2022/RC/ ROC-460  
19/10/2022

Date:

## REFEREE'S COMMENTS

**Title of the Article/note:** Effect of harvest age and storage duration on viability and vigor of shallot tubers

Please tick (√) in the brackets of appropriate word (s) along with comments if any.

- 1. Title**

Suitable	( <input checked="" type="checkbox"/> )	S. Name to be included
Need modification	( <input type="checkbox"/> )	
  
- 2. Abstract**

Adequate	( <input type="checkbox"/> )	Novelty statement missing
Inadequate	( <input checked="" type="checkbox"/> )	
Should be rewritten	( <input type="checkbox"/> )	
  
- 3. Introduction**

Adequate	( <input type="checkbox"/> )	√
Inadequate	( <input type="checkbox"/> )	
Should be rewritten	( <input type="checkbox"/> )	
  
- 4. Materials and Methods**

Adequate	( <input type="checkbox"/> )	Coordinates of site missing.
Inadequate	( <input type="checkbox"/> )	
Should be rewritten	( <input type="checkbox"/> )	
  
- 5. Statistical analysis**

Adequate	( <input type="checkbox"/> )	√
Inadequate	( <input type="checkbox"/> )	
Erroneous	( <input type="checkbox"/> )	
Not analysed	( <input type="checkbox"/> )	
  
- 6. Interpretation of results and discussion**

Adequate	( <input type="checkbox"/> )	Well written
Not supported by data	( <input type="checkbox"/> )	
Suffers from omissions	( <input type="checkbox"/> )	
Vague and generalized	( <input type="checkbox"/> )	
Too brief for clarity	( <input type="checkbox"/> )	
Too comprehensive, must be condensed	( <input type="checkbox"/> )	
Result and Discussion need to be merged	( <input type="checkbox"/> )	
  
- 7. Conclusion**

Adequate	( <input type="checkbox"/> )	√
Need modification	( <input type="checkbox"/> )	
To be rewritten	( <input type="checkbox"/> )	

- 8. Language**
- |                             |        |   |
|-----------------------------|--------|---|
| Grammatically correct       | (    ) |   |
| Spelling / grammar mistakes | (    ) | √ |
| Need revision               | (    ) |   |
- 9. Tables**
- |  |        |   |
|--|--------|---|
| Merge table (s)                              | (    ) |   |
| Rearrange table (s)                          | (    ) |   |
| Delete table (s)                             | (    ) | √ |
| Delete table(s) and incorporate data in text | (    ) |   |
| Furnish table (s) for                        | (    ) |   |
- 10. Illustrations**
- |   |        |   |
|---|--------|---|
| Not required                              | (    ) |   |
| Combine Figures                           | (    ) |   |
| Figure(s) duplicates the data of table(s) | (    ) | √ |
| Figure is desirable for                   | (    ) |   |
- 11. References**
- |   |             |            |
|---|-------------|------------|
| Adequate                                      | (    √    ) |            |
| Omit very old/ irrelevant references          | (    )      |            |
| Inadequate                                    | (    )      | Sufficient |
| References in the text are missing at the end | (    )      |            |
| References at the end are missing in the text | (    )      |            |
| Not as per Journal format                     | (    )      |            |
- 12. Article Rating (1: Poor; 2: Fair; 3: Good; 4: Excellent)**
- |                           |             |
|---------------------------|-------------|
| Originality               | (    4    ) |
| Contribution to the Field | (    4    ) |
| Technical Quality         | (    4    ) |
| Clarity of Presentation   | (    3    ) |
| Depth of Research         | (    4    ) |
- 13. Remarks for Publication**
- |                  |             |
|------------------|-------------|
| May be Published | (    √    ) |
| Minor Revision   | (    √    ) |
| Major Revision   | (    )      |
| Not suitable     | (    )      |
- 14. Are you willing to review revision of this article**
- |                            |             |
|----------------------------|-------------|
| Yes, Required              | (    )      |
| No, to be resubmitted      | (    √    ) |
| Not required; Good as such | (    )      |

**15. Specific comments**

The manuscript is well written. However, it must be revised as per the comments suggested and all the corrections made are to be incorporated before resubmission.

**REVIEWER**

**The manuscript is very well written and may be considered for publication.**

*Viability and vigor of shallot tubers*

**Effect of harvest age and storage duration on viability and vigor of shallot tubers**

**Commented [SH1]:** Include scientific name...

MUHAMMAD ANSAR<sup>1,\*</sup>, BHRUDIN<sup>1</sup>, MAEMUNAH<sup>1</sup> AND PAIMAN<sup>2</sup>

*<sup>1</sup>Department of Agrotechnology, Faculty of Agriculture*

*Universitas Tadulako, Palu 94118, Center Sulawesi, Indonesia*

*\*e-mail: [ansharpasigai@gmail.com](mailto:ansharpasigai@gmail.com)*

<sup>2</sup>Department of Agrotechnology, Faculty of Agriculture, Universitas PGRI Yogyakarta, Yogyakarta 55182, Indonesia.

#### **ABSTRACT**

**Commented [SH2]:** Start abstract with novel background of research on why the present investigation was conducted.

Therefore, this research examined the Lembah Palu variety of shallots (LPVS) as the best raw material for the fried shallot industry. The main problem is that tubers have no standard harvest age and storage duration to support these plants' germination and early growth. Therefore, shallots' growth and productivity are influenced by tubers' viability and vigor. Therefore, this study aimed to investigate the harvest age and storage duration to give the best viability and vigor of shallot tubers. This research was arranged in a randomized complete block design (RCBD) factorial and three replications. The first factor was the harvest age of shallots, which consisted of four levels: 60, 65, 70, and 75 days after planting (DAP). The second factor was the storage duration of tubers, which consisted of four levels: 30, 40, 50, and 60 days after harvest (DAH). The harvesting age of 60-65 DAP significantly affected the parameter of germination power, tuber weight loss, and seedling dry weight. Likewise, the storage duration significantly affected the parameter of germination time,

germination rate, tuber weight loss, and seedling dry weight. These research findings that the harvest age of 60-65 DAP and the storage duration of 30 DAH give the best viability and vigor of shallot tuber. Therefore, we suggest that it is necessary to pay attention to tubers' harvest age and storage duration to find their maximal growth and yield of shallots.

**Key words:** Germination, shallot, tuber, viability, vigor

## INTRODUCTION

Shallots are generally one of the oldest tuber crops used in human life worldwide. The Lembah Palu variety of shallots (LPVS) only exists in the Central Sulawesi Province of Indonesia and has been developed by farming communities since 30 years ago. LPVS tubers are well-known as the main raw material of the highest quality for the fried shallot industry. LPVS is a type of wakegi shallot named *Allium cepa* (L.) *Aggregatum* group (Sulistyaningsih *et al.*, 2008)

LPVS generally can only be developed in lowlands < 400 m ASL. In the Palu Valley area of Central Sulawesi Province, Indonesia. LPVS can grow up to 800 m ASL; if the microenvironment was modified, using a plastic hood and mulch to increase the growth and yields were not significantly different at 100 and 400 m ASL (Pasigai *et al.*, 2016).

Fried shallots produced from LPVS raw materials are one of the leading horticultural products that have the opportunity to become an export commodity. Market demand for fried shallot products is quite high but has not yet been fulfilled due to limited raw materials for LPVS (Bahrudin *et al.*, 2019). It is because the productivity of LPVS is still very low, only 3.5-4.5 tons/ha, while its productivity can reach 9.7 tons/ha (Anonymous, 2011).

One of the problems in increasing LPVS production is using tubers for seedlings that do not yet have a standard harvest age and storage time for tubers before planting to obtain high tuber viability and vigor. Seed vigor is important because several factors can influence it: genetic composition, environmental conditions during production, parent plant nutrition, ripening and harvesting stages, seed size and mass, seed age, mechanical integrity, and environmental conditions during storage (Ignácio *et al.*, 2015). Seeds play an important role as an early determinant of plant growth and development and an important factor in increasing crop yields. Brewster (1994) in Rabinowitch and Currah (2002) describes a complex interaction among factors contributing to tuber quality in post-harvest storage, including cultivar, tuber development stage, early defoliation, skin integrity, and conditions

Commented [SH3]: Scientific name with author.

during ripening, harvesting, and treatment. Saving seeds is an important crop production process to avoid unfavorable environmental conditions and accelerated damage, which starts after harvesting. Storage conditions play an important role in maintaining high seed quality, which is directly related to environmental conditions (especially temperature and relative humidity) (Suriyong *et al.*, 2015).

Seed size is an important physical indicator of seed quality that influences vegetative growth and is often linked to yields, market quality factors, and crop efficiency (Ambika *et al.*, 2014). Seed damage was associated with genotype, seed history, and physiological and chemical composition (Copeland and McDonald, 1999). The shelf life of shallot bulbs is a genetic trait that can be improved by efficient harvesting, post-harvest management, and good tuber storage conditions. The storage quality of shallot bulbs can be influenced by several factors, such as thiosulfonic acid, pyruvic acid, dissolved solids, sugars, and many other biological compounds (Sekara *et al.*, 2017). The power of seeds plays a role in producing agronomic and horticultural crops (Khan *et al.*, 2017). The use of high-quality seeds is one of the essential elements for increasing agricultural production in any agricultural system (Elias, 2018).

Seed viability and viability can predict seed performance under various environmental conditions and strategies to increase seed viability to allow normal growth. However, the sensitivity of seeds to high temperatures depends on the moisture content; the high of water content, the lower the viability (Suriyong *et al.*, 2015). The high physiological potential of seeds with rapid and uniform germination is a fundamental requirement for forming a good plant, especially in adverse environmental conditions. Windauer *et al.* (2012) showed that germination was influenced by temperature and high sensitivity to water deficiency.

Shallots are slow-growing, shallow-rooted plants with no shade habitus. Therefore, their productivity highly depends on water availability in the soil, proper fertilization, and weed control. Storage of shallot nutrient requirements is highest during vegetative growth (Sekara *et al.*, 2017). During the transition from endo-dormancy to ecto dormancy and subsequent growth, shallot bulbs transition from storage to source organs to maintain cell division in meristematic tissue. The biochemical and physiological components were analyzed by identifying the ratio of monosaccharides (fructose and glucose) to disaccharides (sucrose) and the concentration of zeatin riboside to differentiate between germinated and non-germinated tubers. However, this has not been widely used (Chope *et al.*, 2012).



The growth and productivity of shallots are influenced by the viability and strength of the tubers. These characteristic changes are most likely related to the respiration and remobilization of carbohydrates to provide energy for growth. As a result, the tuber receives all the nutrients needed for sprout growth (Chope *et al.*, 2012). Harvest age and storage time determine tuber viability and strength before planting. However, the exact age of harvest and storage of tubers is unknown before planting to obtain higher growth and productivity of shallots. The germination of shallot tubers is strongly influenced by the age of harvest, storage time, and storage area of shallot bulbs. The speed of germination of shallots is determined by the size of the tubers, the content of carbohydrate reserves, and natural growth hormones (Priya *et al.*, 2014).

Handling seeds is different from handling them for consumption. Seed handling starts from production, harvesting, processing, drying, cleaning, packaging, and storage. One postharvest action of shallots is drying to remove excess moisture from the harvested shallot bulb tissue's outer skin, roots, and neck (Wright and Grant, 1997). This aimed to improve and maintain the quality of shallot tubers and reduce the possibility of infection by organisms that cause tuber disease in the warehouse. Seed production was an agronomic study considering genetic, environmental, and technological aspects of land agroecosystem conditions. Therefore, it is important to focus on vigor and viability to produce shallot seeds. Sadjad *et al.* (1999) stated that seed vigor is the ability of seeds to grow and develop into ordinary seeds in optimal growth conditions or powerful seeds that produce above-normal products if planted in optimal health conditions. It was further argued that two pieces of information about viability reflect seed vigors, such as growth strength and storage capacity. Seed viability is determined by pre-harvest conditions, including soil fertility, method and age of harvest, and post-harvest, including drying, seed care, packaging and storage. Ideally all seeds should have high growth strength so that if planted in various field conditions they will still grow healthy and strong and produce high quality with good quality. This study aimed to investigate the harvest age and storage duration to give the best viability and vigor of shallot tubers.

## **MATERIALS AND METHODS**

### **Study site**

The research was conducted in April-September 2017. Observation of the harvest age of LPVS tubers was carried out on farmers' land in Bulupontou Jaya Village, Sigi-Biromaru

**Commented [SH4]:** Avoid old citations. Citations after 2010 are only accepted.

District, Sigi Regency, Central Sulawesi Province, Indonesia. The study was conducted on dry land at 120 m above sea level (ASL). The average daily temperature is 29-30 °C and the humidity was 60-65%. The storage place for tubers was at the Central Sulawesi Horticultural Seed Center (CSSCH) in Sidera Village, Sigi-Biromaru District, Sigi Regency. The storage time for tubers was carried out in the storage room of shallots at 100 m ASL; the average temperature of the daily storage room was between 28 - 29 °C, and humidity ranged from 60-70%.

Commented [SH5]: Provide the coordinates of the study area.

### Experimental design

This research was arranged in RCBD factorial and three replications. The first factor was the harvest age of shallots, which consisted of four levels: 60, 65, 70, and 75 DAP. The second factor was the storage duration of the tubers, which consisted of four levels: 30, 40, 50, and 60 DAH. The effect of harvesting age was carried out by harvesting LPVS tubers from the experimental field in stages according to the predetermined harvest age treatment and followed by the treatment of the storage duration of tubers in a special room for storing shallot bulbs at the CSSCH location. The length of time to store the tubers is adjusted to each predetermined treatment.

Commented [SH6]: ???

### Research procedures

Each experimental plot was given organic fertilizer (bokashi from goat manure) at a dose of 15 tons/ha. The organic fertilizer was evenly mixed on the top of the bed. The experimental plot was 1.05 m (width) × 2.55 m (length) × 0.25 m (height). The distance between treatment beds was 50 cm, while between replicates was 75 cm. Plant spacing was 15 cm × 15 cm. Each hole was planted with one shallot bulb sterilized with a fungicide solution (Dithane M-45), a concentration of 0.02%.

Plant maintenance included providing irrigation water every three days and using a pinwheel (sprinkles) for 1-1.5 hours or until it reaches field capacity. Inorganic fertilizers were given after planting. The first fertilizer was applied at the age of 7 DAP, 100 kg/ha N, 100 kg/ha ZA, 150 kg/ha P, and 100 kg/ha K. At 30 DAP, additional fertilizer was given, 100 kg N/ha. Fertilizer application through a hole of about 5-7 cm on the side of the plant with a depth of 7-10 cm. Pest and disease control used insecticides and fungicides to spray on plants if there were symptoms of the attack. In addition, physical pest control was also carried out by killing pests, especially leaf worms that attack shallots.

## Parameters

The observation parameters of this experiment were germination power (GP), germination time (GT), germination rate (GR), tuber weight loss (TW), and seedling dry weight according to Sadjad *et al.* (1999) formulation as follows:

$$\text{GP (\%)} = \frac{\text{the number of seeds germinated}}{\text{the number of seeds germinated}} \times 100\%$$

(1)

$$\text{GT} = \frac{N_1 T_1 + N_2 T_2 + \dots + N_n T_n}{\text{the number of seeds germinated}} \times 100\%$$

(2)

$$\text{GR} = \frac{\% \text{NS}_1}{\text{etmal}_1} + \frac{\% \text{NS}_2}{\text{etmal}_2} + \dots + \frac{\% \text{NS}_n}{\text{etmal}_n}$$

(3)

$$\text{TW (\%)} = \frac{\text{Weights before storage}}{\text{Weight after storage}} \times 100\%$$

(4)

The seedling dry weight (g) biomass was dried in an oven at a temperature of 70-80 °C for 24 hours or until its weight was constant.

**Note:** N = Number of seeds germinated; T = Time (days); NS = Normal Sprouts

## Statistical analysis

The observed data were analyzed using analysis of variance (ANOVA) (Gomez and Gomez, 1984). The treatment with a significantly different response was followed by Duncan's new multiple range tests (DMRT) at 5% significant level.

## RESULTS AND DISCUSSION

The treatment of harvest age significantly affected germination power, tuber weight loss, and seedlings dry weight of LPVS, but not significantly on germination time and rate. The DMRT at 5% significant level on germination power, tuber weight loss, and seedlings dry weight can be seen in Table 1.

### Germination power

The results showed that the harvest age significantly affected the germination of LPVS tubers. Harvest age 60-65 DAP resulted in higher germination and significantly differed from the harvest age of 75 DAP (Table 1). This was because the tubers that were slow to

harvest had passed physiological maturity. High-quality seeds were inseparable from high viability and vigor. Therefore, one seed viability and vigor determinant was the right harvest time to reach physiological maturity (Suriyong *et al.*, 2015). This condition allowed for rapid and uniform seed germination and better seed formation and development (Cardoso *et al.*, 2015). Differently treated seeds become an effective technique to improve seedling growth and survival percentage compared to conventional seed germination (Kamlesh *et al.*, 2022).

### **Tuber weight loss**

The analysis results showed that the tubers' harvest age and storage time had a major impact on the reduction in tuber weight during storage. Tuber harvesting age of 60-65 DAP resulted in higher tuber weight loss values, namely 14.00% and 13.70%, respectively, significantly different from the 70 and 75 DAP harvest ages. Furthermore, the smallest tuber weight reduction was obtained at the harvest age of 75 DAP (5.60%) (Table 1).

### **Seedlings dry weight**

The analysis showed that the harvesting age and tuber storage time significantly differed from the dry weight of LPVS seedlings. Harvest age 60 DAP produced higher seedling dry weight (3.02 g/plant), significantly different from the harvest age of 75 DAP. Still, it was not significantly different from the harvest age of 65 and 70 DAP (Table 1). It shows that the longer the tuber harvest age will reduce the initial growth of shallot seeds. The dry matter content increased during the period of tuber development. Tubers harvested at 70 DAP will lose most of their leaves because they have withered and died. Harvesting more than 80-90% wilted leaves reduced tuber dry matter content and storage capacity.

The results of the quadratic regression analysis showed that the effect of harvesting age on germination was obtained through the equation:  $y = -0.0195 x^2 + 2.0115 x - 36.03$  and the reduction in tuber weight with the equation:  $y = -0.0066 x^2 + 0.8834 x - 23.462$ . Therefore, based on quadratic regression can be represented in Fig. 1.

Fig. 1 shows that tuber harvesting age affected germination time and weight loss value. After 65 DAS, the harvest age caused the tuber shrinkage value to decrease, and the lowest occurred at the harvest age of 75 DAS. On the other hand, the longer the harvesting age tended to increase the germination time of shallot bulbs, but the increase was relatively small or not significantly different. The reduction in tuber weight at harvest age after 65 DAS

from (13.7%) to the lowest at 75 DAS (5.6%) was caused by the enlargement and filling of the tubers with maximum photosynthetic carbohydrates so that the tubers were denser and had a slight decrease in tuber weight. There were many changes in physiological and biochemical characteristics during the storage of shallot bulbs, including moisture content and concentrations of flavoring compounds, carbohydrates, minerals, and growth regulators (Chope *et al.*, 2012).

The treatment of storage duration significantly affected germination time, germination rate, tuber weight loss, and seedlings dry weight of LPVS, but not significantly on germination power. The DMRT at 5% significant level on germination time, germination rate, tuber weight loss, and seedlings dry weight can be seen in Table 2.

#### **Germination time**

The results showed that the treatment of tuber storage time significantly affected the germination time of LPVS tubers. The 60 DAH storage time resulted in a faster germination time (3.97 days), significantly different from the storage time of 30, 40, and 50 DAH. Tubers stored for 30-40 days required longer germination time, namely 7.10 and 7.85 days, respectively (Table 2).

#### **Germination rate**

The analysis showed that the storage duration for LPVS tubers significantly affected the germination rate. For example, the storage duration for tubers of 60 DAH resulted in the highest germination rate (26.57%/etmal) and significantly differed from the storage duration for 30, 40, and 50 DAH tubers. On the other hand, storage duration of 30-40 DAH resulted in the lowest germination rates of 15.73 and 13.88%/etmal, respectively (Table 2). These results were supported by Grohs *et al.* (2017), stating that environmental conditions greatly determine the storage capacity of seeds during the production process to store seeds.

The storability of seeds was a quantitative trait influenced by environmental factors during seed formation, harvesting, and storage. Sekara *et al.* (2017) stated that the process of seed maturity included morphological and physiological changes that occur from the beginning of planting until the seeds mature into seeds ready for harvest. During the seed ripening process, certain changes occur in the characteristics of the seeds and ovules, including changes in seed size, moisture content, dry weight, and seed vigor. Chemical energy is synthesized during photosynthesis when carbohydrates, lipids, and proteins accumulate in the seeds, providing carbohydrate reserves for germination (Nonogaki, 2006).

The strength of shallot seeds was also greatly influenced by the dormancy period and the tuber storage length. Kapoor *et al.* (2011) stated that there was an increase in gibberellin, auxins, cytokinins, and abscisic acid at the end of tuber dormancy.

### **Tuber weight loss**

Water loss leads to an increase in tuber weight during the storage process. In addition, the length of storage for tubers significantly affected the reduction in tuber weight during storage. The storage time for tubers was up to 60 DAP, resulting in the highest reduction in tuber weight (13.55%) and was significantly different from the storage time for tubers of 30, 40, and 50 DAP (Table 2).

### **Seedlings dry weight**

Furthermore, the storage duration of 30 DAH tubers resulted in higher seedling dry weight (3.14 g/plant) and significantly differed from the storage duration of 60 DAH 1.89 g/plant. However, it was not significantly different from the storage duration for tubers of 40 and 50 DAH (Table 2). These results also indicated that the longer the storage duration would reduce the initial growth of shallot seeds. Germination and initial growth of plants were influenced by environmental factors as well as carbohydrate content, dry weight, and growth hormones in the tubers. Henriksen and Hansen (2001) tuber moisture and carbohydrate content were in ideal conditions to support faster germination with higher dry weight growth. The high dry weight of the seeds indicated that the vigor and viability of the seeds were increasing.

Shallot tubers with high dry weight were more suitable for long-term storage and tended to contain higher fructan concentrations (Chope *et al.*, 2012). However, low seed vigor causes low seed emergence, especially in less-than-ideal soil conditions. Therefore, seeds should be stored in a conducive environment (optimal temperature) so that seed quality remains high until the end of storage. Germination can be increased by adjusting the substrate for higher surface contact and the presence of light and temperature between 25-35 °C (Cardoso *et al.*, 2015).

The result of the quadratic regression analysis showed that the effect of storage duration on germination time was obtained equation:  $y = -0.0085x^2 + 1.6005x - 58.752$  and on tuber weight loss with equation:  $y = -0.0197x^2 + 2.4185x - 66.848$ . Based on quadratic regression can be represented in Fig. 2.

Fig. 2 shows that the storage time for tubers will affect the germination time and the reduction in tuber weight. The length of time it takes for shallot bulbs to germinate was shorter (3.97 days) at 60 DAH. On the other hand, the reduction in tuber weight increased with a longer tuber storage duration (13.55%) at 60 DAH. The increase in tuber weight loss during the longer storage time of the tubers was caused by the ongoing process of tuber respiration, which caused changes in the carbohydrate reserves found in shallot tubers. The results were consistent with the statement of Eshel *et al.* (2014) that the tubers storage for a long time, the tubers began to sprout, the roots began to form, and the weight of the tubers decreased. The results of the analysis showed that there was no significant interaction between harvesting age and storage duration on the vigor and viability of LPVS tubers. Harvesting the tubers at the right level of maturity (physiological maturity) was very important to get a high level of tuber quality. The right time to store the tubers will also result in high vigor and viability. Seeds that have high viability and vigor are shown to have the ability to grow above 80%. The germination percentage increased linearly with the seed age (Ruiz and Parera, 2017).

## **CONCLUSION**

Based on the analysis and discussion of the result, it can be concluded that the harvesting age of 60-65 DAP significantly affected the parameter of germination power, tuber weight loss, and seedling dry weight. Likewise, the storage duration significantly affected the parameter of germination time, germination rate, tuber weight loss, and seedling dry weight. These research findings that the harvest age of 60-65 DAP and the storage duration of 30 DAH give the best viability and vigor of shallot tuber. Therefore, we suggest that it is necessary to pay attention to tubers' harvest age and storage duration to find their maximal growth and yield of shallots.

## **ACKNOWLEDGEMENTS**

This research was carried out with financial support from the Higher Education Excellence Research Scheme for Fiscal Year 2017. For this reason, the authors would like to thank the Directorate General of Higher Education, Ministry of Research, Technology and Higher Education for all the facilities provided so that this research could be carried out properly and this scientific article could be published in scientific journals.

## **REFERENCES**

- Ambika, S., Manonmani, V. and Somasundaram, G. (2014). Review on effect of seed size on seedling vigour and seed yield. *Res J Seed Sci.* **7**(2) : 31–8. <https://doi.org/10.3923/rjss.2014.31.38>
- Anonymous. (2011). *Lampiran Keputusan Menteri Pertanian Nomor: 1843/Kpts/SR.120/4/2011 tentang deskripsi bawang merah varietas Lembah Palu* (pp: 1–1).
- Bahrudin, B., Ansar, M. and Thaha, A. R. (2019). Organic fertilizers are effective in increasing growth and productivity of shallot Lembah Palu varieties. *Agrol Agric Sci J.* **6**(1) : 13–9. <https://doi.org/10.22487/j24077593.2019.v6.i1.12145>
- Cardoso, A. Á., Obolari, A. de M. M., Borges, E. E. de L. e, Silva, C. J. da, and Rodrigues, H. S. (2015). Environmental factors on seed germination, seedling survival and initial growth of sacha inchi (*Plukenetia volubilis* L.). *J Seed Sci.* **37**(2) : 111–6. <https://doi.org/10.1590/2317-1545v37n2145054>
- Chope, G. A., Cools, K., Hammond, J. P., Thompson, A. J. and Terry, L. A. (2012). Physiological, biochemical and transcriptional analysis of onion bulbs during storage. *Ann Bot.* **109** : 819–31. <https://doi.org/10.1093/aob/mcr318>
- Copeland, L. O. and McDonald, M. B. (1999). Principles of seed science and technology. In: *Principles of Seed Science and Technology* (3<sup>rd</sup> ed.). Kluwer academic Publishers. <https://doi.org/10.1007/978-1-4615-1783-2>
- Elias, S. G. (2018). The importance of using high quality seeds in agriculture systems. *Agric Res Technol Open Access J.* **15**(4) : 1–2. <https://doi.org/10.19080/artoaj.2018.14.555961>
- Eshel, D., Teper-Bamnlker, P., Vinokur, Y., Saad, I., Zutahy, Y. and Rodov, V. (2014). Fast curing: A method to improve postharvest quality of onions in hot climate harvest. *Postharvest Biology and Technology.* **88** : 34–9. <https://doi.org/10.1016/j.postharvbio.2013.09.002>
- Gomez, A. G. and Gomez, K. A. (1984). *Statistical procedures for agricultural research* (Sec. Ed.). John Wiley & Sons, Inc.
- Grohs, M., Marchesan, E., Tomazetti, M. B., Roso, R. and Formentini, T. C. (2017). Physiological potential of *Oryza sativa* seeds treated with growth regulators at low



- temperatures. *Rev Cienc Agron.* **48**(3) : 473–9. <https://doi.org/10.5935/1806-6690.20170055>
- Henriksen, K. and Hansen, S. L. (2001). Increasing the dry matter production in bulb onions (*Allium cepa* L.). *Proc. II IS Edible Allianceae Ed. J. Armstrong Acta Hort.* **555** : 147–52. <https://doi.org/10.17660/actahortic.2001.555.20>
- Ignácio, V. L., Nava, I. A., Malavasi, M. de M. and Gris, E. P. (2015). Influence of foliar fertilization with manganese on germination, vigor and storage time of RR soybean seeds. *Revista Ceres.* **62**(5) : 446–52. <https://doi.org/10.1590/0034-737X201562050004>
- Kamlesh, J., Pratima, K., Akansha, S., Gaurav, S. and Esha, R. (2022). Effect of different pre-treatments on seed germination and seedling growth of *Adenanthera pavonina*. *Res Crop.* **23** : 229–34. <https://doi.org/10.31830/2348-7542.2022.032>
- Kapoor, N., Arya, A., Siddiqui, M. A., Kumar, H. and Amir, A. (2011). Physiological and biochemical changes during seed deterioration in aged seeds of rice (*Oryza sativa* L.). *American Journal of Plant Physiology.* **6**(1) : 28–35. <https://doi.org/10.3923/ajpp.2011.28.35>
- Khan, F. A., Bhat, S. A., Narayan, S., Maqbool, R., Murtuza, I. and Khan, F. U. (2017). Seed deterioration and priming - An Overview. *SKUAST J Res.* **19**(1) : 12–21.
- Nonogaki, H. (2006). Seed germination - The biochemical and molecular mechanisms. In: *Breeding Science* (pp: 93–105). <https://doi.org/10.1270/jsbbs.56.93>
- Pasigai, M. A., Thaha, A. R., Nasir, B., Lasmini, S. A., Maemunah, and Bahrudin. (2016). *Teknologi budidaya bawang merah varietas Lembah Palu*. Untad Press.
- Priya, E. P. B., Sinja, V. R., Alice, R. P. J. S., Shanmugasundaram, S. and Alagusundaram, K. (2014). Storage of onions – A review. *Agric Rev.* **35**(4) : 239–249. <https://doi.org/10.5958/0976-0741.2014.00911.8>
- Rabinowitch, H. D. and Currah, L. (2002). Allium crop science: recent advances. In: H. D. R. and L. Currah (Ed.), *CABI Publishing*. CABI Publishing is a division of CAB International. <https://doi.org/10.1079/9780851995106.0000>
- Ruiz, M. B. and Parera, C. A. (2017). Effect of harvesting time on seed quality of two bell

pepper cultivars (*Capsicum annuum*). *Rev la Fac Ciencias Agrar.* **49**(2) : 67–77.

Sadjad, S., Murniati, E. and Ilyas, S. (1999). *Parameter pengujian vigor benih dari komparatif ke simulatif*. (D. Herfan (ed.)). Grasindo. (in Indonesia).


Sekara, A., Pokluda, R., Del Vacchio, L., Somma, S. and Caruso, G. (2017). Interactions among genotype, environment and agronomic practices on production and quality of storage onion (*Allium cepa* L.) - A review. In: *Horticultural Science*. <https://doi.org/10.17221/92/2015-HORTSCI>

Sulistyaningsih, E., Takatori, Y., Isshiki, S. and Tashiro, Y. (2008). *Identification of wakegi onion Indonesia by GISH* (p. 333).

Suriyong, S., Krittigamas, N., Pinmanee, S., Punyalue, A. and Vearasilp, S. (2015). Influence of storage conditions on change of hemp seed quality. *Agric Agric Sci Procedia.* **5** : 170–176. <https://doi.org/10.1016/j.aaspro.2015.08.026>

Windauer, L. B., Martinez, J., Rapoport, D., Wassner, D. and Benech-Arnold, R. (2012). Germination responses to temperature and water potential in *Jatropha curcas* seeds: A hydrotime model explains the difference between dormancy expression and dormancy induction at different incubation temperatures. *Ann Bot.* **109**(1) : 265–73. <https://doi.org/10.1093/aob/mcr242>

Wright, P. J. and Grant, D. G. (1997). Effects of cultural practices at harvest on onion bulb quality and incidence of rots in storage. *New Zeal J Crop Hortic Sci.* **25**(4) : 353–8. <https://doi.org/10.1080/01140671.1997.9514026>



**Lampiran Manuscript yang telah direvisi:**

**The manuscript is very well written and may be considered for publication.**

*Viability and vigor of shallot tubers*

**Effect of harvest age and storage duration on viability and vigor of shallot (*Allium cepa* L.) tubers**

MUHAMMAD ANSAR<sup>1,\*</sup>, BAHRUDIN<sup>1</sup>, MAEMUNAH<sup>1</sup> AND PAIMAN<sup>2</sup>

*<sup>1</sup>Department of Agrotechnology, Faculty of Agriculture  
Universitas Tadulako, Palu 94118, Center Sulawesi, Indonesia*

*\*(e-mail: [ansharpasigai@gmail.com](mailto:ansharpasigai@gmail.com))*

*<sup>2</sup>Department of Agrotechnology, Faculty of Agriculture, Universitas PGRI Yogyakarta,  
Yogyakarta 55182, Indonesia.*

**ABSTRACT**

Shallot is one of the tuber crops used in human life worldwide. Lembah Palu variety of shallot (LPVS) is the best raw material for the fried shallot industry. The main problem is that tubers have no standard harvest age and storage duration to support these plants' germination and early growth. Therefore, shallots' growth and productivity are influenced by tubers' viability and vigor. This study aimed to investigate the harvest age and storage duration to give the best viability and vigor of shallot tubers. This research was arranged in a randomized complete block design (RCBD) factorial and three replications. The first factor was the harvest age of shallots, which consisted of four levels: 60, 65, 70, and 75 days after planting (DAP). The second factor was the storage duration of tubers, which consisted of four levels: 30, 40, 50, and 60 days after harvest (DAH). The harvesting age of 60-65 DAP significantly affected the parameter of germination power, tuber weight loss, and seedling dry weight. Likewise, the storage duration significantly affected the parameter of germination time, germination rate, tuber weight loss, and seedling dry weight. These research findings that the harvest age of 60-65 DAP and the storage duration of 30 DAH give the best viability and vigor of shallot tuber. Therefore, we suggest that it is necessary to pay attention to tubers' harvest age and storage duration to find their maximal growth and yield of shallots.

**Key words:** Germination, shallot, tuber, viability, vigor

## INTRODUCTION

Shallot (*Allium cepa* L.) is generally one of the oldest tuber crops used in human life worldwide. The Lembah Palu variety of shallots (LPVS) only exists in the Central Sulawesi Province of Indonesia and has been developed by farming communities since 30 years ago. LPVS tubers are well-known as the main raw material of the highest quality for the fried shallot industry.

LPVS generally can only be developed in lowlands < 400 m above sea level (ASL). In the Palu Valley area of Central Sulawesi Province, Indonesia. LPVS can grow up to 800 m ASL; if the microenvironment was modified, using a plastic hood and mulch to increase the growth and yields were not significantly different at 100 and 400 m ASL (Pasigai *et al.*, 2016).

Fried shallots produced from LPVS raw materials are one of the leading horticultural products that have the opportunity to become an export commodity. Market demand for fried shallot products is quite high but has not yet been fulfilled due to limited raw materials

for LPVS (Bahrudin *et al.*, 2019). The productivity of LPVS is still very low, only 3.5-4.5 tons/ha, while its productivity can reach 9.7 tons/ha (Anonymous, 2011).

One of the problems in increasing LPVS production is using tubers for seedlings that do not yet have a standard harvest age and storage duration for tubers before planting to obtain high tuber viability and vigor. Seed vigor is important because several factors can influence it: genetic composition, environmental conditions during production, parent plant nutrition, ripening and harvesting stages, seed size and mass, seed age, mechanical integrity, and environmental conditions during storage (Ignácio *et al.*, 2015). Tubers play an important role as an early determinant of plant growth and development and an important factor in increasing crop yields. Saving tubers are an important crop production process to avoid unfavorable environmental conditions and accelerated damage, which starts after harvesting. Storage conditions play an important role in maintaining high seed quality, which is directly related to environmental conditions (especially temperature and relative humidity) (Suriyong *et al.*, 2015).

Seed size is an important physical indicator of seed quality that influences vegetative growth and is often linked to yields, market quality factors, and crop efficiency (Ambika *et al.*, 2014). Seed damage was associated with genotype, seed history, and physiological and chemical composition (Copeland and McDonald, 1999). The shelf life of shallot bulbs is a genetic trait that can be improved by efficient harvesting, post-harvest management, and good tuber storage conditions. The storage quality of shallot bulbs can be influenced by several factors, such as thiosulfonic acid, pyruvic acid, dissolved solids, sugars, and many other biological compounds (Sekara *et al.*, 2017). The power of seeds plays a role in producing agronomic and horticultural crops (Khan *et al.*, 2017). The use of high-quality seeds is one of the essential elements for increasing agricultural production in any agricultural system (Elias, 2018).

Seed viability and viability can predict seed performance under various environmental conditions and strategies to increase seed viability to allow normal growth. However, the sensitivity of seeds to high temperatures depends on the moisture content; the high of water content, the lower the viability (Suriyong *et al.*, 2015). The high physiological potential of seeds with rapid and uniform germination is a fundamental requirement for forming a good plant, especially in adverse environmental conditions. Windauer *et al.* (2012) showed that germination was influenced by temperature and high sensitivity to water deficiency.

Shallots are slow-growing, shallow-rooted plants with no shade habitus. Therefore, their productivity highly depends on water availability in the soil, proper fertilization, and weed control. Storage of shallot nutrient requirements is highest during vegetative growth (Sekara *et al.*, 2017). During the transition from endo-dormancy to ecto dormancy and subsequent growth, shallot bulbs transition from storage to source organs to maintain cell division in meristematic tissue. The biochemical and physiological components were analyzed by identifying the ratio of monosaccharides (fructose and glucose) to disaccharides (sucrose) and the concentration of zeatin riboside to differentiate between germinated and non-germinated tubers. However, this has not been widely used (Chope *et al.*, 2012).

The growth and productivity of shallots are influenced by the viability and strength of the tubers. These characteristic changes are most likely related to the respiration and remobilization of carbohydrates to provide energy for growth. As a result, the tuber receives all the nutrients needed for sprout growth (Chope *et al.*, 2012). Harvest age and storage time determine tuber viability and strength before planting. However, the exact age of harvest and storage of tubers is unknown before planting to obtain higher growth and productivity of shallots. The germination of shallot tubers is strongly influenced by the age of harvest, storage time, and storage area of shallot bulbs. The speed of germination of shallots is determined by the size of the tubers, the content of carbohydrate reserves, and natural growth hormones (Priya *et al.*, 2014).

Handling tubers is different from handling them for consumption. Handling tubers starts from production, harvesting, processing, drying, cleaning, packaging, and storage. One postharvest action of shallots is drying to remove excess moisture from the harvested shallot bulb tissue's outer skin, roots, and neck. It aimed to improve and maintain the quality and reduce the possibility of infection by organisms that cause tuber disease in the warehouse. Tubers production were an agronomic study considering genetic, environmental, and technological aspects of land agroecosystem conditions. Therefore, it is important to focus on vigor and viability to produce shallot tubers.

Vigor is the ability of tubers to grow and develop into ordinary in optimal growth conditions or powerful that produce above-normal products if planted in optimal health conditions. It was further argued that two pieces of information about viability reflect tubers vigors, such as growth strength and storage capacity. tubers viability are determined by pre-harvest conditions, including soil fertility, method and age of harvest, and post-harvest,

including drying, seed care, packaging and storage. Ideally all seeds should have high growth strength so that if planted in various field conditions they will still grow healthy and strong and produce high quality with good quality. This study aimed to investigate the harvest age and storage duration to give the best viability and vigor of shallot tubers.

## **MATERIALS AND METHODS**

### **Study site**

The research was conducted in April-September 2017. Observation of the harvest age of LPVS tubers was carried out on farmers' land in Bulupontou Jaya Village, Sigi-Biromaru District, Sigi Regency, Central Sulawesi Province, Indonesia. Central Sulawesi located between 2°22 North Latitude and 3°48 South Latitude and between 119°22' and 124°22 East Longitude. The study was conducted on dry land at 120 m above sea level (ASL). The average daily temperature is 29-30 °C and the humidity was 60-65%. The storage place for tubers was at the Central Sulawesi Horticultural Seed Center (CSSCH) in Sidera Village, Sigi-Biromaru District, Sigi Regency. The storage time for tubers was carried out in the storage room of shallots at 100 m ASL; the average temperature of the daily storage room was between 28 - 29 °C, and humidity ranged from 60-70%.

### **Experimental design**

This research was arranged in RCBD factorial and three replications. The first factor was the harvest age of shallots, which consisted of four levels: 60, 65, 70, and 75 DAP. The second factor was the storage duration of the tubers, which consisted of four levels: 30, 40, 50, and 60 DAH. The effect of harvesting age was carried out by harvesting LPVS tubers from the experimental field in stages according to the predetermined harvest age treatment and followed by the treatment of the storage duration of tubers in a special room for storing shallot bulbs at the CSSCH location. The length of time to store the tubers is adjusted to each predetermined treatment.

### **Research procedures**

Each experimental plot was given organic fertilizer (bokashi from goat manure) at a dose of 15 tons/ha. The organic fertilizer was evenly mixed on the top of the bed. The experimental plot was 1.05 m (width) × 2.55 m (length) × 0.25 m (height). The distance between treatment beds was 50 cm, while between replicates was 75 cm. Plant spacing was

15 cm × 15 cm. Each hole was planted with one shallot bulb sterilized with a fungicide solution (Dithane M-45), a concentration of 0.02%.

Plant maintenance included providing irrigation water every three days and using a pinwheel (sprinkles) for 1-1.5 hours or until it reaches field capacity. Inorganic fertilizers were given after planting. The first fertilizer was applied at the age of 7 DAP, 100 kg/ha N, 100 kg/ha ZA, 150 kg/ha P, and 100 kg/ha K. At 30 DAP, additional fertilizer was given, 100 kg N/ha. Fertilizer application through a hole of about 5-7 cm on the side of the plant with a depth of 7-10 cm. Pest and disease control used insecticides and fungicides to spray on plants if there were symptoms of the attack. In addition, physical pest control was also carried out by killing pests, especially leaf worms that attack shallots.

### Parameters

The observation parameters of this experiment were germination power (GP), germination time (GT), germination rate (GR), tuber weight loss (TW), and seedling dry weight can be seen in Eq. 1, 2, 3, and 4.

$$GP (\%) = \frac{\text{the number of seeds germinated}}{\text{the number of seeds germinated}} \times 100\%$$

(1)

$$GT = \frac{N_1 T_1 + N_2 T_2 + \dots + N_n T_n}{\text{the number of seeds germinated}} \times 100\%$$

(2)

$$GR = \frac{\% NS_1}{\text{etmal}_1} + \frac{\% NS_2}{\text{etmal}_2} + \dots + \frac{\% NS_n}{\text{etmal}_n}$$

(3)

$$TW (\%) = \frac{\text{Weights before storage}}{\text{Weight after storage}} \times 100\%$$

(4)

The seedling dry weight (g) biomass was dried in an oven at a temperature of 70-80 °C for 24 hours or until its weight was constant.

**Note:** N = Number of seeds germinated; T = Time (days); NS = Normal Sprouts

### Statistical analysis

The observed data were analyzed using analysis of variance (ANOVA) (Gomez and Gomez, 1984). The treatment with a significantly different response was followed by Duncan's new multiple range tests (DMRT) at 5% significant level.



## RESULTS AND DISCUSSION

The treatment of harvest age significantly affected germination power, tuber weight loss, and seedlings dry weight of LPVS, but not significantly on germination time and rate. The DMRT at 5% significant level on germination power, tuber weight loss, and seedlings dry weight can be seen in Table 1.

### Germination power

The results showed that the harvest age significantly affected the germination of LPVS tubers. Harvest age 60-65 DAP resulted in higher germination and significantly differed from the harvest age of 75 DAP (Table 1). This was because the tubers that were slow to harvest had passed physiological maturity. High-quality seeds were inseparable from high viability and vigor. Therefore, one seed viability and vigor determinant was the right harvest time to reach physiological maturity (Suriyong *et al.*, 2015). This condition allowed for rapid and uniform seed germination and better seed formation and development (Cardoso *et al.*, 2015). Differently treated seeds become an effective technique to improve seedling growth and survival percentage compared to conventional seed germination (Kamlesh *et al.*, 2022).

### Tuber weight loss

The analysis results showed that the tubers' harvest age and storage time had a major impact on the reduction in tuber weight during storage. Tuber harvesting age of 60-65 DAP resulted in higher tuber weight loss values, namely 14.00% and 13.70%, respectively, significantly different from the 70 and 75 DAP harvest ages. Furthermore, the smallest tuber weight reduction was obtained at the harvest age of 75 DAP (5.60%) (Table 1).

### Seedlings dry weight

The analysis showed that the harvesting age and tuber storage time significantly differed from the dry weight of LPVS seedlings. Harvest age 60 DAP produced higher seedling dry weight (3.02 g/plant), significantly different from the harvest age of 75 DAP. Still, it was not significantly different from the harvest age of 65 and 70 DAP (Table 1). It shows that the longer the tuber harvest age will reduce the initial growth of shallot seeds. The dry matter content increased during the period of tuber development. Tubers harvested at 70 DAP will lose most of their leaves because they have withered and died. Harvesting more than 80-90% wilted leaves reduced tuber dry matter content and storage capacity.

The results of the quadratic regression analysis showed that the effect of harvesting age on germination was obtained through the equation:  $y = -0.0195 x^2 + 2.0115 x - 36.03$  and the reduction in tuber weight with the equation:  $y = -0.0066 x^2 + 0.8834 x - 23.462$ . Therefore, based on quadratic regression can be represented in Fig. 1.

Fig. 1 shows that tuber harvesting age affected germination time and weight loss value. After 65 DAS, the harvest age caused the tuber shrinkage value to decrease, and the lowest occurred at the harvest age of 75 DAS. On the other hand, the longer the harvesting age tended to increase the germination time of shallot bulbs, but the increase was relatively small or not significantly different. The reduction in tuber weight at harvest age after 65 DAS from (13.7%) to the lowest at 75 DAS (5.6%) was caused by the enlargement and filling of the tubers with maximum photosynthetic carbohydrates so that the tubers were denser and had a slight decrease in tuber weight. There were many changes in physiological and biochemical characteristics during the storage of shallot bulbs, including moisture content and concentrations of flavoring compounds, carbohydrates, minerals, and growth regulators (Chope *et al.*, 2012).

The treatment of storage duration significantly affected germination time, germination rate, tuber weight loss, and seedlings dry weight of LPVS, but not significantly on germination power. The DMRT at 5% significant level on germination time, germination rate, tuber weight loss, and seedlings dry weight can be seen in Table 2.

#### **Germination time**

The results showed that the treatment of tuber storage time significantly affected the germination time of LPVS tubers. The 60 DAH storage time resulted in a faster germination time (3.97 days), significantly different from the storage time of 30, 40, and 50 DAH. Tubers stored for 30-40 days required longer germination time, namely 7.10 and 7.85 days, respectively (Table 2).

#### **Germination rate**

The analysis showed that the storage duration for LPVS tubers significantly affected the germination rate. For example, the storage duration for tubers of 60 DAH resulted in the highest germination rate (26.57%/etmal) and significantly differed from the storage duration for 30, 40, and 50 DAH tubers. On the other hand, storage duration of 30-40 DAH resulted in the lowest germination rates of 15.73 and 13.88%/etmal, respectively (Table 2). These

results were supported by Grohs *et al.* (2017), stating that environmental conditions greatly determine the storage capacity of seeds during the production process to store seeds.

The storability of seeds was a quantitative trait influenced by environmental factors during seed formation, harvesting, and storage. Sekara *et al.* (2017) stated that the process of seed maturity included morphological and physiological changes that occur from the beginning of planting until the seeds mature into seeds ready for harvest. During the seed ripening process, certain changes occur in the characteristics of the seeds and ovules, including changes in seed size, moisture content, dry weight, and seed vigor. Chemical energy is synthesized during photosynthesis when carbohydrates, lipids, and proteins accumulate in the seeds, providing carbohydrate reserves for germination. The strength of shallot seeds was also greatly influenced by the dormancy period and the tuber storage length. Kapoor *et al.* (2011) stated that there was an increase in gibberellin, auxins, cytokinins, and abscisic acid at the end of tuber dormancy.

#### **Tuber weight loss**

Water loss leads to an increase in tuber weight during the storage process. In addition, the length of storage for tubers significantly affected the reduction in tuber weight during storage. The storage time for tubers was up to 60 DAP, resulting in the highest reduction in tuber weight (13.55%) and was significantly different from the storage time for tubers of 30, 40, and 50 DAP (Table 2).

#### **Seedlings dry weight**

Furthermore, the storage duration of 30 DAH tubers resulted in higher seedling dry weight (3.14 g/plant) and significantly differed from the storage duration of 60 DAH 1.89 g/plant. However, it was not significantly different from the storage duration for tubers of 40 and 50 DAH (Table 2). These results also indicated that the longer the storage duration would reduce the initial growth of shallot seeds. Germination and initial growth of plants were influenced by environmental factors as well as carbohydrate content, dry weight, and growth hormones in the tubers. Tuber moisture and carbohydrate content were in ideal conditions to support faster germination with higher dry weight growth. The high dry weight of the seeds indicated that the vigor and viability of the seeds were increasing.

Shallot tubers with high dry weight were more suitable for long-term storage and tended to contain higher fructan concentrations (Chope *et al.*, 2012). However, low seed vigor

causes low seed emergence, especially in less-than-ideal soil conditions. Therefore, seeds should be stored in a conducive environment (optimal temperature) so that seed quality remains high until the end of storage. Germination can be increased by adjusting the substrate for higher surface contact and the presence of light and temperature between 25-35 °C (Cardoso *et al.*, 2015).

The result of the quadratic regression analysis showed that the effect of storage duration on germination time was obtained equation:  $y = -0.0085x^2 + 1.6005x - 58.752$  and on tuber weight loss with equation:  $y = -0.0197x^2 + 2.4185x - 66.848$ . Based on quadratic regression can be represented in Fig. 2.

Fig. 2 shows that the storage time for tubers will affect the germination time and the reduction in tuber weight. The length of time it takes for shallot bulbs to germinate was shorter (3.97 days) at 60 DAH. On the other hand, the reduction in tuber weight increased with a longer tuber storage duration (13.55%) at 60 DAH. The increase in tuber weight loss during the longer storage time of the tubers was caused by the ongoing process of tuber respiration, which caused changes in the carbohydrate reserves found in shallot tubers. The results were consistent with the statement of Eshel *et al.* (2014) that the tubers storage for a long time, the tubers began to sprout, the roots began to form, and the weight of the tubers decreased. The results of the analysis showed that there was no significant interaction between harvesting age and storage duration on the vigor and viability of LPVS tubers. Harvesting the tubers at the right level of maturity (physiological maturity) was very important to get a high level of tuber quality. The right time to store the tubers will also result in high vigor and viability. Seeds that have high viability and vigor are shown to have the ability to grow above 80%. The germination percentage increased linearly with the seed age (Ruiz and Parera, 2017).

## CONCLUSION

Based on the analysis and discussion of the result, it can be concluded that the harvesting age of 60-65 DAP significantly affected the parameter of germination power, tuber weight loss, and seedling dry weight. Likewise, the storage duration significantly affected the parameter of germination time, germination rate, tuber weight loss, and seedling dry weight. These research findings that the harvest age of 60-65 DAP and the storage duration of 30 DAH give the best viability and vigor of shallot tuber. Therefore, we suggest that it is

necessary to pay attention to tubers' harvest age and storage duration to find their maximal growth and yield of shallots.

#### ACKNOWLEDGEMENTS

This research was carried out with financial support from the Higher Education Excellence Research Scheme for Fiscal Year 2017. For this reason, the authors would like to thank the Directorate General of Higher Education, Ministry of Research, Technology and Higher Education for all the facilities provided so that this research could be carried out properly and this scientific article could be published in scientific journals.

#### REFERENCES

- Ambika, S., Manonmani, V. and Somasundaram, G. (2014). Review on effect of seed size on seedling vigour and seed yield. *Res J Seed Sci.* **7**: 31–8.
- Anonymous. (2011). *Lampiran Keputusan Menteri Pertanian Nomor: 1843/Kpts/SR.120/4/2011 tentang deskripsi bawang merah varietas Lembah Palu.* Kementrian Pertanian, Indonesia. pp. 1–1.
- Bahrudin, B., Ansar, M. and Thaha, A. R. (2019). Organic fertilizers are effective in increasing growth and productivity of shallot Lembah Palu varieties. *Agrol Agric Sci J.* **6**: 13–9.
- Cardoso, A. Á., Obolari, A. D. M. M., Borges, E. E. D. L. E, Silva, C. J. D. and Rodrigues, H. S. (2015). Environmental factors on seed germination, seedling survival and initial growth of sacha inchi (*Plukenetia volubilis* L.). *J Seed Sci.* **37**: 111–6.
- Chope, G. A., Cools, K., Hammond, J. P., Thompson, A. J. and Terry, L. A. (2012). Physiological, biochemical and transcriptional analysis of onion bulbs during storage. *Ann Bot.* **109**: 819–31.
- Copeland, L. O. and McDonald, M. B. (1999). *Seed viability and viability testing.* In: Principles of Seed Science and Technology (3<sup>rd</sup> ed.). Springer Nature Switzerland AG. Part of [Springer Nature](#). pp. 124-39.
- Elias, S. G. (2018). The importance of using high quality seeds in agriculture systems. *Agric Res Technol Open Access J.* **15**: 1–2.
- Eshel, D., Teper-Bamnolker, P., Vinokur, Y., Saad, I., Zutahy, Y. and Rodov, V. (2014). Fast curing: A method to improve postharvest quality of onions in hot climate harvest.

*Postharvest Biol Technol.* **88**: 34–9.

- Gomez, A. G. and Gomez, K. A. (1984). *Statistical procedures for agricultural research* (Sec. Ed.). John Wiley & Sons, Los Bannos, Phillipines. pp. 680.
- Grohs, M., Marchesan, E., Tomazetti, M. B., Roso, R. and Formentini, T. C. (2017). Physiological potential of *Oryza sativa* seeds treated with growth regulators at low temperatures. *Rev Cienc Agron.* **48**: 473–9.
- Ignácio, V. L., Nava, I. A., Malavasi, M. de M. and Gris, E. P. (2015). Influence of foliar fertilization with manganese on germination, vigor and storage time of RR soybean seeds. *Rev Ceres.* **62**: 446–52.
- Kamlesh, J., Pratima, K., Akansha, S., Gaurav, S. and Esha, R. (2022). Effect of different pre-treatments on seed germination and seedling growth of *Adenanthera pavonina*. *Res Crop.* **23**: 229–34.
- Kapoor, N., Arya, A., Siddiqui, M. A., Kumar, H. and Amir, A. (2011). Physiological and biochemical changes during seed deterioration in aged seeds of rice (*Oryza sativa* L.). *Am J Plant Physiol.* **6**: 28–35.
- Khan, F. A., Bhat, S. A., Narayan, S., Maqbool, R., Murtuza, I. and Khan, F. U. (2017). Seed deterioration and priming - An Overview. *SKUAST J Res.* **19**: 12–21.
- Pasigai, M. A., Thaha, A. R., Nasir, B., Lasmini, S. A., Maemunah, and Bahrudin. (2016). *Teknologi budidaya bawang merah varietas Lembah Palu*. Untad Press, Palu, Indonesia. pp. 130.
- Priya, E. P. B., Sinja, V. R., Alice, R. P. J. S., Shanmugasundaram, S. and Alagusundaram, K. (2014). Storage of onions – A review. *Agric Rev.* **35**: 239–49.
- Ruiz, M. B. and Parera, C. A. (2017). Effect of harvesting time on seed quality of two bell pepper cultivars (*Capsicum annuum*). *Rev la Fac Ciencias Agrar.* **49**: 67–77.
- Sekara, A., Pokluda, R., Del Vacchio, L., Somma, S. and Caruso, G. (2017). Interactions among genotype, environment and agronomic practices on production and quality of storage onion (*Allium cepa* L.) - A review. *Hort. Sci.* **44**: 21-42.
- Suriyong, S., Krittigamas, N., Pinmanee, S., Punyalue, A. and Vearasilp, S. (2015). Influence of storage conditions on change of hemp seed quality. *Agric Agric Sci*

*Procedia*. **5**: 170–6.

Windauer, L. B., Martinez, J., Rapoport, D., Wassner, D. and Benech-Arnold, R. (2012). Germination responses to temperature and water potential in *Jatropha curcas* seeds: A hydrotime model explains the difference between dormancy expression and dormancy induction at different incubation temperatures. *Ann Bot.* **109**: 265–73.

Table 1. The effect of harvest age on germination power, tuber weight loss, and seedlings dry weight

Harvest age (DAP)	Germination power (%)	Tuber weight loss (%)	Seedling dry weight (g/plant)
60	100.00 a	14.00 a	3.02 a
65	99.17 a	13.70 a	2.82 ab
70	97.50 ab	7.85 b	2.15 ab
75	94.17 b	5.60 c	2.02 b

Note: The average number followed by the same letter in the same column shows no significant difference based on the DMRT at 5% significance level.

Table 2. The effect of storage duration on germination time, germination rate, tuber weight loss, and seedling dry weight.

Bulb storage duration (DAH)	Germination time (days)	Germination rate (%/etmal)	Tuber weight loss (%)	Seedling dry weight (g/plant)
30	7.10 a	15.73 c	6.60 d	3.14 a
40	7.85 a	13.88 c	9.60 c	2.77 ab
50	5.19 b	20.23 b	11.40 b	2.21 ab
60	3.97 c	26.57 a	13.55 a	1.89 b

Note: The average number followed by the same letter in the same column shows no significant difference based on the DMRT at 5% significance level.






Figure 1. The effect of harvest age on germination time and the tuber weight loss.





Figure 2. The effect of storage duration on germination time and tuber weight loss.

## Tanggal 28 Oktober 2022: Penerimaan Paper untuk di Publikasi



Print ISSN: 0972-3226  
Online ISSN: 2348-7534

# RESEARCH ON CROPS

[Indexed by Scopus Q3/Elsevier, EBSCO, CrossRef, Google Scholar, DRJI, CABL, ICI etc.]

Ref. No.: GP-2022/ROC/11146/912

Dated: October 28, 2022

### Subject: Acceptance of paper for publication

Dear Ansar,

It is to inform you that your following paper/article has been peer reviewed by the expert reviewers and found suitable for publication. The paper is accepted for publication as full-length paper in RESEARCH ON CROPS journal Vol. 23, No. 4 (December) 2022.

**Effect of harvest age and storage duration on viability and vigor of shallot (*Allium cepa* L.) tubers - Muhammad Ansar, Bahrudin, Maemunah and Paiman**

#### Read the following:

1. The above article will cover 6 printed pages of the journal including 2 tables and 2 figures..
2. Indexed in Scopus, EBSCO, CiteFactor, Scientific Indexing Services Products and Services, etc.
3. Scopus Q3: SciMago Journal Ranking (SJR) 2021: 0.27
4. Scopus Impact Score 2021: 1.70
5. ResearchGate journal Impact 2019/2020: 0.17
6. NAAS, New Delhi rating/scoring for 2022: 4.56
7. Impact Factor (CiteFactor) 2020-2021: 1.30
8. Approved (SIS), CrossRef, DRJI, CABL, ICI, Google Scholar, ResearchGate, UDL-EDGE by UGC (Included in CARE list of journals)
9. Each published paper is assigned with DOI number.
10. The pdf copy of published paper will be sent to corresponding author free of cost.
11. The pdf of galley paper for correction will be sent to author in last week of December 2022.
12. The print copy of journal is available on sale @ US \$ 100 per copy.


Yours Sincerely,

Sd/-  
(Managing Editor)

### **GAURAV PUBLICATIONS, REGD.**

Systematic Printers, Udayapuria Street, Video Market Hisar-125 001  
Haryana, India

[www.gauravpublications.com](http://www.gauravpublications.com) ; [www.cropresearch.org](http://www.cropresearch.org)




Tel: + 91 9354 324922  
email: info@gauravpublications.com  
researchoncrops@gmail.com

## GAURAV PUBLICATIONS, REGD.

Systematic Printers, Udayapuria Street, Near Video Market, Hisar-125 001, India

Muhammad Ansar Department of Agrotechnology Faculty of Agriculture Universitas Tadulako Palu 94118, Center Sulawesi Indonesia		INVOICE	
		<b>Invoice No. :</b> ROC-2022/11146	
		<b>Date :</b> October 28, 2022	
		<b>Your Ref. No:</b> Your email dated October 04, 2022	
Sr. No.	Particulars	Rate (US \$)	Amount (US \$)
1	Publication charges of 6 printed pages including 2 tables and 2 figures of your paper ID No. GP 2022/ROC-11146/912 entitled "Effect of harvest age and storage duration on viability and vigor of shallot ( <i>Allium cepa</i> L.) tubers - Muhammad Ansar, Bahrudin, Maemunah and Paiman" accepted for publication in RESEARCH ON CROPS journal Vol. 23, No. 4 (December) 2022.	US \$ 100/- (Per printed page)	600.00
<b>In words:</b> Six hundred US dollars only		<b>Total (US \$)</b>	<b>600.00</b>
<b>Note:</b> Payment should be sent through Bank Transfer/SWIFT Transfer. Our bank details are as under:  <b>1.Name and address of Bank :</b> State Bank of India, Patel Nagar, Hissar-125001, Haryana, India <b>2.Beneficiary account name :</b> Gaurav Publications <b>3.Beneficiary account number :</b> 00000037014428711 <b>4.Swift Code :</b> SBININBB189 <b>5.IFSC Code :</b> SBIN0009280  <i>Send scan copy of the payment voucher/document through email for our record.</i>			
			<i>Sd/-</i> Managing Editor

**Tanggal 4 November 2022: Transfer Biaya Publikasi pd Journal Research on Crops**

 **Muhammad Anshar Pasigai**  
kepada Gaurav ▾ 22.26 (0 menit yang lalu) ☆ ↶ ⋮


Dear Editor-in-Chief (Dr. Vedpal Singh)

We are grateful for the acceptance of our manuscript entitled: "**Effect of harvest age and storage duration on viability and vigor of shallot (*Allium cepa* L.) tubers** - Muhammad Ansar, Bahrudin, Maemunah and Paiman for publication in the journal RESEARCH ON CROPS. Vol.23, No.4 (December) 2022. For this reason, we send a transfer receipt for payment of publication fees according to the instructions on the invoice.

Thank you  
Best Regard

Muhammad Ansar  
Tadulako University-Indonesia  
\*\*\*

Satu lampiran • Dipindai dengan Gmail ⓘ



November, 04 20...

↶ Balas    ↷ Teruskan

**4 November 2022: Transfer Publication fee**

Receipt of transfer Payment manuscript Entitled "Effect of harvest age and storage duration on viability and vigor of shallot (*Allium cepa* L.) tubers - Muhammad Ansar, Bahrudin, Maemunah and Paiman for publication in the journal RESEARCH ON CROPS. Vol.23, No.4 (December) 2022.

PT. BANK NEGARA INDONESIA (Persero), Tbk  
CABANG : PALU

IBOC - Maintenance (S10)

Teller ID : 88273  
Date : 04/11/2022  
Time : 10:22:50



Sender's Reference:  
:20:S10PLU00018722  
Bank Operation Code:  
:23B:CRED  
Value Date/Currency/Interbank Settled Amount:  
:32A:221104USD600,  
Ordering Customer:  
:50K:/0000000203560938  
BPK MUHAMMAD ANSHAR  
BTN ROVIGA BLOK B3 NO 16 PALU  
SULAWESI TENGAH INDONESIA  
Ordering Institution:  
:52A:BNINIDJAXXX  
Account With Institution:  
:57D:STATE BANK OF INDIA  
PATEL NAGAR, HISSAR-125001  
HARYANA, INDIA  
Beneficiary Customer:  
:59:/00000037014428711  
GAURAV PUBLICATIONS  
INDIA  
Remittance Information:  
:70:PUBLICATION FEE AN MUHAMMAD ANSAR  
UNIVERSITAS TADULAKO  
PALU SULAWESI TENGAH  
Details Of Charges:  
:71A:OUR



PT. Bank Negara Indonesia (Persero) Tbk  
Kantor Cabang Palu  
Jl. Jend. Sudirman No. 58  
Palu 94112 Indonesia  
www.bni.co.id

REFERENCE : S10PLU00018722

NO. TRX. : 88273 959111 96962 TRAN 04/11/2022 10:08:01  
NO. REK. : 000000203560938 Bpk MUHAMMAD ANSHAR  
JUMLAH : IDR 433,125- 1568  
222 - PALU

NO. TRX. : 88273 959111 96962 TRAN 04/11/2022 10:08:01  
NO. REK. : 222360420801001 PENDAPATAN PROPISI KU  
JUMLAH : IDR 35,000 1568  
222 - PALU


NO. TRX. : 88273 959111 96962 TRAN 04/11/2022 10:08:01  
NO. REK. : 222360482010001 Pendapatan Restitusi B  
JUMLAH : IDR 398,125 1568  
222 - PALU

NO. TRX. : 88273 959111 96962 TRAN 04/11/2022 10:08:01  
NO. REK. : 000000203560938 Bpk MUHAMMAD ANSHAR  
JUMLAH : IDR 9,555,000- 1568  
222 - PALU

NO. TRX. : 88273 959111 96962 TRAN 04/11/2022 10:08:01  
NO. REK. : 222840200101001 KU YAKIR  
JUMLAH : USD 600 1568  
222 - PALU



PT. Bank Negara Indonesia (Persero) Tbk  
Kantor Cabang Palu  
Jl. Jend. Sudirman No. 58  
Palu 94112 Indonesia  
www.bni.co.id



Lampiran

*Viability and vigor of shallot tubers*

*Res. Crop.* **23** (4) : (2022)

*With two figures*

*Printed in India*

**Effect of harvest age and storage duration on viability and vigor of shallot (*Allium cepa* L.) tubers**

MUHAMMAD ANSAR<sup>1,\*</sup>, BAHRUDIN<sup>1</sup>, MAEMUNAH<sup>1</sup> AND PAIMAN<sup>2</sup>

<sup>1</sup>*Department of Agrotechnology, Faculty of Agriculture*

*Universitas Tadulako, Palu 94118, Center Sulawesi, Indonesia*

*\*(e-mail: [ansharpasigai@gmail.com](mailto:ansharpasigai@gmail.com))*

(Received: October 04, 2022 / Accepted: November 28, 2022)

**ABSTRACT**

Shallot is one of the tuber crops used in human life worldwide. Lembah Palu variety of shallot (LPVS) is the best raw material for the fried shallot industry. The main problem is that tubers have no standard harvest age and storage duration to support these plants' germination and early growth. Therefore, shallots' growth and productivity are influenced by tubers' viability and vigor. This study aimed to investigate the harvest age and storage duration to give the best viability and vigor of shallot tubers. This research was arranged in a randomized complete block design (RCBD) factorial and three replications. The first factor was the harvest age of shallots, which consisted of four levels: 60, 65, 70, and 75 days after planting (DAP). The second factor was the storage duration of tubers, which consisted of four levels: 30, 40, 50, and 60 days after harvest (DAH). The harvesting age of 60-65 DAP significantly affected the parameter of germination power, tuber weight loss, and seedling dry weight. Likewise, the storage duration significantly affected the parameter of germination time, germination rate, tuber weight loss, and seedling dry weight. These research findings show that the harvest age of 60-65 DAP and the storage duration of 30 DAH give the best viability and vigor of shallot tuber. Therefore, we suggest that it is necessary to pay attention to tubers' harvest age and storage duration to find their maximal growth and yield of shallots.

**Key words:** Germination, shallot, tuber, viability, vigor

---

<sup>2</sup>Department of Agrotechnology, Faculty of Agriculture, Universitas PGRI Yogyakarta, Yogyakarta 55182, Indonesia.

## INTRODUCTION

Shallot (*Allium cepa* L.) is generally one of the oldest tuber crops used in human life worldwide. The Lembah Palu variety of shallots (LPVS) only exists in the Central Sulawesi Province of Indonesia and has been developed by farming communities for 30 years ago. LPVS tubers are well-known as the main raw material of the highest quality for the fried shallot industry.

LPVS generally can only be developed in lowlands < 400 m above sea level (ASL). In the Palu Valley area of Central Sulawesi Province, Indonesia. LPVS can grow up to 800 m ASL; if the microenvironment was modified, using a plastic hood and mulch to increase the growth and yields were not significantly different at 100 and 400 m ASL (Pasigai *et al.*, 2016).

Fried shallots produced from LPVS raw materials are one of the leading horticultural products that have the opportunity to become an export commodity. Market demand for fried shallot products is quite high but has not yet been fulfilled due to limited raw materials for LPVS (Bahrudin *et al.*, 2019). The productivity of LPVS is still very low, only 3.5-4.5 tons/ha, while its productivity can reach 9.7 tons/ha (Anonymous, 2011).

One of the problems in increasing LPVS production is using tubers for seedlings that do not yet have a standard harvest age and storage duration for tubers before planting to obtain high tuber viability and vigor. Seed vigor is important because several factors can influence it: genetic composition, environmental conditions during production, parent plant nutrition, ripening and harvesting stages, seed size and mass, seed age, mechanical integrity, and environmental conditions during storage (Ignácio *et al.*, 2015). Tubers play an important role as an early determinant of plant growth and development and an important factor in increasing crop yields. Saving tubers is an important crop production process to avoid unfavorable environmental conditions and accelerated damage, which starts after harvesting. Storage conditions play an important role in maintaining high seed quality, which is directly related to environmental conditions (especially temperature and relative humidity) (Suriyong *et al.*, 2015).

Seed size is an important physical indicator of seed quality that influences vegetative growth and is often linked to yields, market quality factors, and crop efficiency (Ambika *et al.*, 2014). Seed damage was associated with genotype, seed history, and physiological and chemical composition (Copeland and McDonald, 1999). The shelf life of shallot bulbs is a genetic trait that can be improved by efficient harvesting, post-harvest management, and good tuber storage conditions. The storage quality of shallot bulbs can be influenced by several factors, such as thiosulfonic acid, pyruvic acid, dissolved solids, sugars, and many other biological compounds (Sekara *et al.*, 2017). The power of seeds plays a role in producing agronomic and horticultural crops (Khan *et al.*, 2017). The use of high-quality seeds is one of the essential elements for increasing agricultural production in any agricultural system (Elias, 2018).

Seed viability and viability can predict seed performance under various environmental conditions and strategies to increase seed viability to allow normal growth. However, the sensitivity of seeds to high temperatures depends on the moisture content; the high of water content, the lower the viability (Suriyong *et al.*, 2015). The high physiological potential of seeds with rapid and uniform germination is a fundamental requirement for forming a good plant, especially in adverse environmental conditions. Windauer *et al.* (2012) showed that germination was influenced by temperature and high sensitivity to water deficiency.



Shallots are slow-growing, shallow-rooted plants with no shade habitus. Therefore, their productivity highly depends on water availability in the soil, proper fertilization, and weed control. Storage of shallot nutrient requirements is highest during vegetative growth (Sekara *et al.*, 2017). During the transition from endo-dormancy to ecto dormancy and subsequent growth, shallot bulbs transition from storage to source organs to maintain cell division in meristematic tissue. The biochemical and physiological components were analyzed by identifying the ratio of monosaccharides (fructose and glucose) to disaccharides (sucrose) and the concentration of zeatin riboside to differentiate between germinated and non-germinated tubers. However, this has not been widely used (Chope *et al.*, 2012).

The growth and productivity of shallots are influenced by the viability and strength of the tubers. These characteristic changes are most likely related to the respiration and remobilization of carbohydrates to provide energy for growth. As a result, the tuber receives all the nutrients needed for sprout growth (Chope *et al.*, 2012). Harvest age and storage time determine tuber viability and strength before planting. However, the exact age of harvest and storage of tubers is unknown before planting to obtain higher growth and productivity of shallots. The germination of shallot tubers is strongly influenced by the age of harvest, storage time, and storage area of shallot bulbs. The speed of germination of shallots is determined by the size of the tubers, the content of carbohydrate reserves, and natural growth hormones (Priya *et al.*, 2014).

Handling tubers is different from handling them for consumption. Handling tubers starts from production, harvesting, processing, drying, cleaning, packaging, and storage. One postharvest action of shallots is drying to remove excess moisture from the harvested shallot bulb tissue's outer skin, roots, and neck. It aimed to improve and maintain the quality and reduce the possibility of infection by organisms that cause tuber disease in the warehouse. Tubers production were an agronomic study considering genetic, environmental, and technological aspects of land agroecosystem conditions. Therefore, it is important to focus on vigor and viability to produce shallot tubers.

Vigor is the ability of tubers to grow and develop into ordinary in optimal growth conditions or powerful that produce above-normal products if planted in optimal health conditions. It was further argued that two pieces of information about viability reflect tubers vigors, such as growth strength and storage capacity. tubers viability are determined by pre-harvest conditions, including soil fertility, method and age of harvest, and post-harvest, including drying, seed care, packaging and storage. Ideally all seeds should have high growth strength so that if planted in various field conditions they will still grow healthy and strong and produce high quality with good quality. This study aimed to investigate the harvest age and storage duration to give the best viability and vigor of shallot tubers.

## **MATERIALS AND METHODS**

### **Study Site**

The research was conducted in April-September 2017. Observation of the harvest age of LPVS tubers was carried out on farmers' land in Bulupontou Jaya Village, Sigi-Biromaru District, Sigi Regency, Central Sulawesi Province, Indonesia. Central Sulawesi located between 2°22 North Latitude and 3°48 South Latitude and between 119°22' and 124°22 East Longitude. The study was conducted on dry land at 120 m above sea level (ASL). The average daily temperature is 29-30 °C and the humidity was 60-65%. The storage place for tubers was at the Central Sulawesi Horticultural Seed Center (CSSCH) in Sidera Village, Sigi-Biromaru District, Sigi Regency. The storage time for tubers was carried out in the

storage room of shallots at 100 m ASL; the average temperature of the daily storage room was between 28 - 29 °C, and humidity ranged from 60-70%.

### Experimental Design

This research was arranged in RCBD factorial and three replications. The first factor was the harvest age of shallots, which consisted of four levels: 60, 65, 70, and 75 DAP. The second factor was the storage duration of the tubers, which consisted of four levels: 30, 40, 50, and 60 DAH. The effect of harvesting age was carried out by harvesting LPVS tubers from the experimental field in stages according to the predetermined harvest age treatment and followed by the treatment of the storage duration of tubers in a special room for storing shallot bulbs at the CSSCH location. The length of time to store the tubers is adjusted to each predetermined treatment.

### Research Procedures

Each experimental plot was given organic fertilizer (bokashi from goat manure) at a dose of 15 tons/ha. The organic fertilizer was evenly mixed on the top of the bed. The experimental plot was 1.05 m (width) × 2.55 m (length) × 0.25 m (height). The distance between treatment beds was 50 cm, while between replicates was 75 cm. Plant spacing was 15 cm × 15 cm. Each hole was planted with one shallot bulb sterilized with a fungicide solution (Dithane M-45), a concentration of 0.02%.

Plant maintenance included providing irrigation water every three days and using a pinwheel (sprinkles) for 1-1.5 hours or until it reaches field capacity. Inorganic fertilizers were given after planting. The first fertilizer was applied at the age of 7 DAP, 100 kg/ha N, 100 kg/ha ZA, 150 kg/ha P, and 100 kg/ha K. At 30 DAP, additional fertilizer was given, 100 kg N/ha. Fertilizer application through a hole of about 5-7 cm on the side of the plant with a depth of 7-10 cm. Pest and disease control used insecticides and fungicides to spray on plants if there were symptoms of the attack. In addition, physical pest control was also carried out by killing pests, especially leaf worms that attack shallots.

### Parameters

The observation parameters of this experiment were germination power (GP), germination time (GT), germination rate (GR), tuber weight loss (TW), and seedling dry weight can be seen in Eq. 1, 2, 3 and 4.

$$GP (\%) = \frac{\text{the number of seeds germinated}}{\text{the number of seeds germinated}} \times 100\%$$

(1)

$$GT = \frac{N_1 T_1 + N_2 T_2 + \dots + N_n T_n}{\text{the number of seeds germinated}} \times 100\%$$

(2)

$$GR = \frac{\% NS_1}{etmal_1} + \frac{\% NS_2}{etmal_2} + \dots + \frac{\% NS_n}{etmal_n}$$

(3)

$$TW (\%) = \frac{\text{Weights before storage}}{\text{Weight after storage}} \times 100\%$$

(4)

The seedling dry weight (g) biomass was dried in an oven at a temperature of 70-80 °C for 24 hours or until its weight was constant. N = Number of seeds germinated; T = Time (days); NS = Normal Sprouts.

### Statistical Analysis

The observed data were analyzed using analysis of variance (ANOVA) (Gomez and Gomez, 1984). The treatment with a significantly different response was followed by Duncan's new multiple range tests (DMRT) at 5% significant level.

## RESULTS AND DISCUSSION

The treatment of harvest age significantly affected germination power, tuber weight loss, and seedlings dry weight of LPVS, but not significantly on germination time and rate. The DMRT at 5% significant level on germination power, tuber weight loss, and seedlings dry weight can be seen in Table 1.

### Germination Power

The results showed that the harvest age significantly affected the germination of LPVS tubers. Harvest age 60-65 DAP resulted in higher germination and significantly differed from the harvest age of 75 DAP (Table 1). This was because the tubers that were slow to harvest had passed physiological maturity. High-quality seeds were inseparable from high viability and vigor. Therefore, one seed viability and vigor determinant was the right harvest time to reach physiological maturity (Suriyong *et al.*, 2015). This condition allowed for rapid and uniform seed germination and better seed formation and development (Cardoso *et al.*, 2015). Differently treated seeds become an effective technique to improve seedling growth and survival percentage compared to conventional seed germination (Kamlesh *et al.*, 2022).

### Tuber Weight Loss

The analysis results showed that the tubers' harvest age and storage time had a major impact on the reduction in tuber weight during storage. Tuber harvesting age of 60-65 DAP resulted in higher tuber weight loss values, namely 14.00% and 13.70%, respectively, significantly different from the 70 and 75 DAP harvest ages. Furthermore, the smallest tuber weight reduction was obtained at the harvest age of 75 DAP (5.60%) (Table 1).

### Seedlings Dry Weight

The analysis showed that the harvesting age and tuber storage time significantly differed from the dry weight of LPVS seedlings. Harvest age 60 DAP produced higher seedling dry weight (3.02 g/plant), significantly different from the harvest age of 75 DAP. Still, it was not significantly different from the harvest age of 65 and 70 DAP (Table 1). It shows that the longer the tuber harvest age will reduce the initial growth of shallot seeds. The dry matter content increased during the period of tuber development. Tubers harvested at 70 DAP will lose most of their leaves because they have withered and died. Harvesting more than 80-90% wilted leaves reduced tuber dry matter content and storage capacity.

The results of the quadratic regression analysis showed that the effect of harvesting age on germination was obtained through the equation:  $y = -0.0195 x^2 + 2.0115 x - 36.03$  and the reduction in tuber weight with the equation:  $y = -0.0066 x^2 + 0.8834 x - 23.462$ . Therefore, based on quadratic regression can be represented in Fig. 1.

Fig. 1 shows that tuber harvesting age affected germination time and weight loss value. After 65 DAS, the harvest age caused the tuber shrinkage value to decrease, and the lowest occurred at the harvest age of 75 DAS. On the other hand, the longer the harvesting age tended to increase the germination time of shallot bulbs, but the increase was relatively small or not significantly different. The reduction in tuber weight at harvest age after 65 DAS from (13.7%) to the lowest at 75 DAS (5.6%) was caused by the enlargement and filling of

the tubers with maximum photosynthetic carbohydrates so that the tubers were denser and had a slight decrease in tuber weight. There were many changes in physiological and biochemical characteristics during the storage of shallot bulbs, including moisture content and concentrations of flavoring compounds, carbohydrates, minerals, and growth regulators (Chope *et al.*, 2012).

The treatment of storage duration significantly affected germination time, germination rate, tuber weight loss, and seedlings dry weight of LPVS, but not significantly on germination power. The DMRT at 5% significant level on germination time, germination rate, tuber weight loss, and seedlings dry weight can be seen in Table 2.

#### **Germination Time**

The results showed that the treatment of tuber storage time significantly affected the germination time of LPVS tubers. The 60 DAH storage time resulted in a faster germination time (3.97 days), significantly different from the storage time of 30, 40, and 50 DAH. Tubers stored for 30-40 days required longer germination time, namely 7.10 and 7.85 days, respectively (Table 2).

#### **Germination Rate**

The analysis showed that the storage duration for LPVS tubers significantly affected the germination rate. For example, the storage duration for tubers of 60 DAH resulted in the highest germination rate (26.57%/etmal) and significantly differed from the storage duration for 30, 40, and 50 DAH tubers. On the other hand, storage duration of 30-40 DAH resulted in the lowest germination rates of 15.73 and 13.88%/etmal, respectively (Table 2). These results were supported by Grohs *et al.* (2017), stating that environmental conditions greatly determine the storage capacity of seeds during the production process to store seeds.

The storability of seeds was a quantitative trait influenced by environmental factors during seed formation, harvesting, and storage. Sekara *et al.* (2017) stated that the process of seed maturity included morphological and physiological changes that occur from the beginning of planting until the seeds mature into seeds ready for harvest. During the seed ripening process, certain changes occur in the characteristics of the seeds and ovules, including changes in seed size, moisture content, dry weight, and seed vigor. Chemical energy is synthesized during photosynthesis when carbohydrates, lipids, and proteins accumulate in the seeds, providing carbohydrate reserves for germination. The strength of shallot seeds was also greatly influenced by the dormancy period and the tuber storage length. Kapoor *et al.* (2011) stated that there was an increase in gibberellin, auxins, cytokinins, and abscisic acid at the end of tuber dormancy.

#### **Tuber Weight Loss**

Water loss leads to an increase in tuber weight during the storage process. In addition, the length of storage for tubers significantly affected the reduction in tuber weight during storage. The storage time for tubers was up to 60 DAP, resulting in the highest reduction in tuber weight (13.55%) and was significantly different from the storage time for tubers of 30, 40, and 50 DAP (Table 2).

#### **Seedlings Dry Weight**

Furthermore, the storage duration of 30 DAH tubers resulted in higher seedling dry weight (3.14 g/plant) and significantly differed from the storage duration of 60 DAH 1.89 g/plant. However, it was not significantly different from the storage duration for tubers of 40 and 50 DAH (Table 2). These results also indicated that the longer the storage duration would reduce the initial growth of shallot seeds. Germination and initial growth of plants

were influenced by environmental factors as well as carbohydrate content, dry weight, and growth hormones in the tubers. Tuber moisture and carbohydrate content were in ideal conditions to support faster germination with higher dry weight growth. The high dry weight of the seeds indicated that the vigor and viability of the seeds were increasing.

Shallot tubers with high dry weight were more suitable for long-term storage and tended to contain higher fructan concentrations (Chope *et al.*, 2012). However, low seed vigor causes low seed emergence, especially in less-than-ideal soil conditions. Therefore, seeds should be stored in a conducive environment (optimal temperature) so that seed quality remains high until the end of storage. Germination can be increased by adjusting the substrate for higher surface contact and the presence of light and temperature between 25-35 °C (Cardoso *et al.*, 2015).

The result of the quadratic regression analysis showed that the effect of storage duration on germination time was obtained equation:  $y = -0.0085 x^2 + 1.6005 x - 58.752$  and on tuber weight loss with equation:  $y = -0.0197 x^2 + 2.4185 x - 66.848$ . Based on quadratic regression can be represented in Fig. 2.

Fig. 2 shows that the storage time for tubers will affect the germination time and the reduction in tuber weight. The length of time it takes for shallot bulbs to germinate was shorter (3.97 days) at 60 DAH. On the other hand, the reduction in tuber weight increased with a longer tuber storage duration (13.55%) at 60 DAH. The increase in tuber weight loss during the longer storage time of the tubers was caused by the ongoing process of tuber respiration, which caused changes in the carbohydrate reserves found in shallot tubers. The results were consistent with the statement of Eshel *et al.* (2014) that the tubers storage for a long time, the tubers began to sprout, the roots began to form, and the weight of the tubers decreased. The results of the analysis showed that there was no significant interaction between harvesting age and storage duration on the vigor and viability of LPVS tubers. Harvesting the tubers at the right level of maturity (physiological maturity) was very important to get a high level of tuber quality. The right time to store the tubers will also result in high vigor and viability. Seeds that have high viability and vigor are shown to have the ability to grow above 80%. The germination percentage increased linearly with the seed age (Ruiz and Parera, 2017).

## CONCLUSION

Based on the analysis and discussion of the result, it can be concluded that the harvesting age of 60-65 DAP significantly affected the parameter of germination power, tuber weight loss, and seedling dry weight. Likewise, the storage duration significantly affected the parameter of germination time, germination rate, tuber weight loss, and seedling dry weight. These research findings that the harvest age of 60-65 DAP and the storage duration of 30 DAH give the best viability and vigor of shallot tuber. Therefore, we suggest that it is necessary to pay attention to tubers' harvest age and storage duration to find their maximal growth and yield of shallots.

## ACKNOWLEDGEMENTS

This research was carried out with financial support from the Higher Education Excellence Research Scheme for Fiscal Year 2017. For this reason, the authors would like to thank the Directorate General of Higher Education, Ministry of Research, Technology and Higher Education for all the facilities provided so that this research could be carried out properly and this scientific article could be published in scientific journals.

## REFERENCES

- Ambika, S., Manonmani, V. and Somasundaram, G. (2014). Review on effect of seed size on seedling vigour and seed yield. *Res J Seed Sci.* **7** : 31-38.
- Anonymous (2011). *Lampiran Keputusan Menteri Pertanian Nomor: 1843/Kpts/SR.120/4/2011 tentang deskripsi bawang merah varietas Lembah Palu.* Kementerian Pertanian, Indonesia. pp. 1–1.
- Bahrudin, B., Ansar, M. and Thaha, A. R. (2019). Organic fertilizers are effective in increasing growth and productivity of shallot Lembah Palu varieties. *Agrol. Agric. Sci. J.* **6** : 13-19.
- Cardoso, A. Á., Obolari, A. D. M. M., Borges, E. E. D. L. E, Silva, C. J. D. and Rodrigues, H. S. (2015). Environmental factors on seed germination, seedling survival and initial growth of sacha inchi (*Plukenetia volubilis* L.). *J. Seed Sci.* **37** : 111–16.
- Chope, G. A., Cools, K., Hammond, J. P., Thompson, A. J. and Terry, L. A. (2012). Physiological, biochemical and transcriptional analysis of onion bulbs during storage. *Ann. Bot.* **109** : 819–31.
- Copeland, L. O. and McDonald, M. B. (1999). Seed viability and viability testing. In: Principles of Seed Science and Technology (3<sup>rd</sup> ed.). Springer Nature Switzerland AG. Part of Springer Nature. Pp. 124-39.
- Elias, S. G. (2018). The importance of using high quality seeds in agriculture systems. *Agric Res Technol Open Access J.* **15** : 1-2.
- Eshel, D., Teper-Bamnlker, P., Vinokur, Y., Saad, I., Zutahy, Y. and Rodov, V. (2014). Fast curing: A method to improve postharvest quality of onions in hot climate harvest. *Postharvest Biol. Technol.* **88** : 34-39.
- Gomez, A. G. and Gomez, K. A. (1984). Statistical procedures for agricultural research (Sec. Ed.). John Wiley & Sons, Los Bannos, Phillipines. Pp. 680.
- Grohs, M., Marchesan, E., Tomazetti, M. B., Roso, R. and Formentini, T. C. (2017). Physiological potential of *Oryza sativa* seeds treated with growth regulators at low temperatures. *Rev. Cienc. Agron.* **48** : 473-79.
- Ignácio, V. L., Nava, I. A., Malavasi, M. de M. and Gris, E. P. (2015). Influence of foliar fertilization with manganese on germination, vigor and storage time of RR soybean seeds. *Rev. Ceres.* **62** : 446-52.
- Kamlesh, J., Pratima, K., Akansha, S., Gaurav, S. and Esha, R. (2022). Effect of different pre-treatments on seed germination and seedling growth of *Adenanthera pavonina*. *Res. Crop.* **23** : 229-34.
- Kapoor, N., Arya, A., Siddiqui, M. A., Kumar, H. and Amir, A. (2011). Physiological and biochemical changes during seed deterioration in aged seeds of rice (*Oryza sativa* L.). *Am. J. Plant Physiol.* **6** : 28-35.
- Khan, F. A., Bhat, S. A., Narayan, S., Maqbool, R., Murtuza, I. and Khan, F. U. (2017). Seed deterioration and priming - An Overview. *SKUAST J. Res.* **19** : 12-21.
- Pasigai, M. A., Thaha, A. R., Nasir, B., Lasmini, S. A., Maemunah, and Bahrudin. (2016).

Teknologi budidaya bawang merah varietas Lembah Palu. Untad Press, Palu, Indonesia. Pp. 130.

- Priya, E. P. B., Sinja, V. R., Alice, R. P. J. S., Shanmugasundaram, S. and Alagusundaram, K. (2014). Storage of onions - A review. *Agric. Rev.* **35** : 239-49.
- Ruiz, M. B. and Parera, C. A. (2017). Effect of harvesting time on seed quality of two bell pepper cultivars (*Capsicum annuum*). *Rev. la Fac. Ciencias Agrar.* **49** : 67-77.
- Sekara, A., Pokluda, R., Del Vacchio, L., Somma, S. and Caruso, G. (2017). Interactions among genotype, environment and agronomic practices on production and quality of storage onion (*Allium cepa* L.) - A review. *Hort. Sci.* **44** : 21-42.
- Suriyong, S., Krittigamas, N., Pinmanee, S., Punyalue, A. and Vearasilp, S. (2015). Influence of storage conditions on change of hemp seed quality. *Agric. Sci. Procedia.* **5** : 170-76.
- Windauer, L. B., Martinez, J., Rapoport, D., Wassner, D. and Benech-Arnold, R. (2012). Germination responses to temperature and water potential in *Jatropha curcas* seeds: A hydrotime model explains the difference between dormancy expression and dormancy induction at different incubation temperatures. *Ann. Bot.* **109** : 265-73.

Table 1. The effect of harvest age on germination power, tuber weight loss, and seedlings dry weight.

Harvest age (DAP)	Germination power (%)	Tuber weight loss (%)	Seedling dry weight (g/plant)
60	100.00 a	14.00 a	3.02 a
65	99.17 a	13.70 a	2.82 ab
70	97.50 ab	7.85 b	2.15 ab
75	94.17 b	5.60 c	2.02 b

Note: The average number followed by the same letter in the same column shows no significant difference based on the DMRT at 5% significance level.

Table 2. The effect of storage duration on germination time, germination rate, tuber weight loss, and seedling dry weight.

Bulb storage duration (DAH)	Germination time (days)	Germination rate (%/etmal)	Tuber weight loss (%)	Seedling dry weight (g/plant)
30	7.10 a	15.73 c	6.60 d	3.14 a
40	7.85 a	13.88 c	9.60 c	2.77 ab
50	5.19 b	20.23 b	11.40 b	2.21 ab
60	3.97 c	26.57 a	13.55 a	1.89 b

Note: The average number followed by the same letter in the same column shows no significant difference based on the DMRT at 5% significance level.




Fig. 1. Effect of harvest age on germination time and the tuber weight loss.






Fig. 2. The effect of storage duration on germination time and tuber weight loss.





### 17 Desember 2022: online Publication



<https://www.gauravpublications.com/journal/research-on-crops/volume-23/issue-4-december/ROC-909>

gauravpublications.com/onlineFirstArticles

2.bp.blogspot.com

Effect of harvest age and storage duration on viability and vigor of shallot (*Allium cepa* L.) tubers  
MUHAMMAD ANSAR, BAHRUDIN, MAEMUNAH AND PAIMAN  
DOI:10.31830/2348-7542.2022.ROC-909 | Research on Crops


Profitable eco-friendly tomato (*Solanum lycopersicum*) farming through the use of eco-enzymes in Indonesia  
ARDIYANTA, CIGILIA TRI KUSUMASTUTI, OKTI PURWANINGSIH AND PAIMAN  
DOI:10.31830/2348-7542.2022.ROC-908 | Research on Crops

Role of biochar amendments in improving the properties of acid sulphate soil  
AGUSALIM MASULILI, SUTIKARINI, RINI SURYANI, IDA AYU SUCI, ISMAIL ASTAR, HARDI DOMINIKUS BANCIN AND PAIMAN  
DOI:10.31830/2348-7542.2022.ROC-907 | Research on Crops

Effect of addition of biopolymer coated DAP on phosphorus release kinetics over an incubation time  
M. DHANALAKSHMI, S. MEENA, P. JANAKI, S. KARTHIKEYAN AND L. NALINA  
DOI:10.31830/2348-7542.2022.ROC-906 | Research on Crops

Influence of basic tillage and coultis strips on the weed infestation and yield of spring wheat (*Triticum aestivum*) in the Tawang region

Message  
 Saya bukan robot  
Submit



gauravpublications.com/journal/research-on-crops/volume-23/issue-4-december/ROC-909

2.bp.blogspot.com

Gaurav Publications Journals Services Policies LOGIN

Research On Crops  
Home > Research on Crops > VOLUME 23 > ISSUE 4 (DECEMBER) > ROC-909


Effect of harvest age and storage duration on viability and vigor of shallot (*Allium cepa* L.) tubers

DOI: 10.31830/2348-7542.2022.ROC-909 | Article Id: ROC-909

Authors and affiliations  
MUHAMMAD ANSAR, BAHRUDIN, MAEMUNAH AND PAIMAN  
ansharpasigai@gmail.com  
Address : Department of Agrotechnology, Faculty of Agriculture, Universitas Tadulako, Palu 94118, Center Sulawesi, Indonesia

Online Published: 17-12-2022

Abstract



Online Published: 17-12-2022

### Abstract

Shallot is one of the tuber crops used in human life worldwide. Lembah Palu variety of shallot (LPVS) is the best raw material for the fried shallot industry. The main problem is that tubers have no standard harvest age and storage duration to support these plants' germination and early growth. Therefore, shallot's growth and productivity are influenced by tubers' viability and vigor. This study aimed to investigate the harvest age and storage duration to give the best viability and vigor of shallot tubers. This research was arranged in a randomized complete block design (RCBD) factorial and three replications. The first factor was the harvest age of shallots, which consisted of four levels: 60, 65, 70, and 75 days after planting (DAP). The second factor was the storage duration of tubers, which consisted of four levels: 30, 40, 50, and 60 days after harvest (DAH). The harvesting age of 60-65 DAP significantly affected the parameter of germination power, tuber weight loss, and seedling dry weight. Likewise, the storage duration significantly affected the parameter of germination time, germination rate, tuber weight loss, and seedling dry weight. These research findings show that the harvest age of 60-65 DAP and the storage duration of 30 DAH give the best viability and vigor of shallot tuber. Therefore, we suggest that it is necessary to pay attention to tubers' harvest age and storage duration to find their maximal growth and yield of shallots.

### Keywords

Germination shallot tuber viability vigor

### References

Ambika, S., Manonmani, V. and Somasundaram, G. (2014). Review on effect of seed size on seedling vigour and seed yield. *Res J Seed Sci.* 7 : 31-38.

Anonymous (2011). *Lampiran Keputusan Menteri Pertanian Nomor: 1843/Kpts/SR.120/4/2011 tentang deskripsi bawang merah varietas Lembah Palu*. Kementerian Pertanian, Indonesia. pp. 1-1.

Bahrudin, B., Ansar, M. and Thaba, A. R. (2019). Organic fertilizers are effective in increasing growth and productivity of shallot Lembah Palu varieties. *Agron. Agric. Sci. J.* 6 : 13-19.

Cardoso, A. A., Obolari, A. D. M. M., Borges, E. E. D. L. E., Silva, C. J. D. and Rodrigues, H. S. (2015). Environmental factors on seed germination, seedling survival and initial growth of sachá inchí (*Plukenetia volubilis* L.). *J. Seed Sci.* 37 : 111-116.

Chepe, G. A., Cook, K., Hammond, J. P., Thompson, A. J. and Terry, L. A. (2012). Physiological, biochemical and transcriptional analysis of onion bulbs during storage. *Ann. Bot.* 109 : 819-831.

Copeland, L. O. and McDonald, M. B. (1999). *Seed viability and viability testing*. In: Principles of Seed Science and Technology (3<sup>rd</sup> ed.). Springer Nature Switzerland AG. Part of Springer Nature. Pp. 124-39.

Elias, S. G. (2018). The importance of using high quality seeds in agriculture systems. *Agric Res Technol Open Access J.* 15 : 1-2.

Eshel, D., Teper-Bammolker, P., Vinokur, Y., Saad, I., Zutahy, Y. and Rodov, V. (2014). Fast curing: A method to improve postharvest quality of onions in hot climate harvest. *Postharvest Biol. Technol.* 88 : 34-39.

Gomez, A. G. and Gomez, K. A. (1984). *Statistical procedures for agricultural research* (Sec. Ed.). John Wiley & Sons, Los Bannos, Philippines. Pp. 680.

Grohs, M., Marchesan, E., Tomazetti, M. B., Roso, R. and Formentini, T. C. (2017). Physiological potential of *Oryza sativa* seeds treated with growth regulators at low temperatures. *Rev. Cienc. Agron.* 48 : 473-79.

Ignácio, V. L., Nava, I. A., Malavasi, M. de M. and Gris, E. P. (2015). Influence of foliar fertilization with manganese on germination, vigor and storage time of RR soybean seeds. *Rev. Cienc.* 62 : 146-52.

Kamlesh, J., Pratima, K., Akansha, S., Gaurav, S. and Esha, R. (2022). Effect of different pre-treatments on seed germination and seedling growth of *Adenanthera pavonina*. *Res. Crop.* 93 : 206-214.



- Editorial Board
- Author guidelines
- Aims and Scope
- Role of Reviewers
- Editorial Policy
- Role of Authors
- Peer-Review Article Process
- For Readers
- Subject Covered
- Subscribe
- Terms and Conditions

### References

- Subscribe
- Terms and Conditions
- Publish**  
your articles with us  
Submit Article Now
- Article Status**  
Please enter your article no. to get the current status  
Search Article
- Contact Us

Effect of harvest age and storage duration on viability and vigor of shallot (*Allium cepa* L.) tubers \*  
[gauravpublications.com/journal/research-on-crops/volume-23/issue-4-december/ROC-909](https://gauravpublications.com/journal/research-on-crops/volume-23/issue-4-december/ROC-909)

2.bp.blogspot.com

Gaurav Publications  
 Journals Services Policies LOGIN

Kamlesh, J., Pratima, K., Akanksha, S., Gaurav, S. and Esha, R. (2022). Effect of different pre-treatments on seed germination and seedling growth of *Adenanthera pavonina*. *Res. Crop.* **23** : 229-244.  
 Kapoor, N., Arya, A., Siddiqui, M. A., Kumar, H. and Amir, A. (2011). Physiological and biochemical changes during seed deterioration in aged seeds of rice (*Oryza sativa* L.). *Am. J. Plant Physiol.* **6** : 28-35.  
 Khan, F. A., Bhat, S. A., Narayan, S., Maqbool, R., Muhtuza, I. and Khan, F. U. (2017). Seed deterioration and priming - An Overview. *SKUAST J. Res.* **10** : 12-21.  
 Pasigai, M. A., Thaha, A. R., Nasir, B., Lasmuni, S. A., Maemunah, and Bahrodin. (2016). Teknologi budidaya bawang merah varietas Lembah Pala. Utad Press, Palu, Indonesia. Pp. 130.  
 Priya, E. P. B., Sija, V. R., Alice, R. P. J. S., Shanmugasundaram, S. and Alagusundaram, K. (2014). Storage of onions - A review. *Agric. Res.* **33** : 239-49.  
 Ruiz, M. B. and Parera, C. A. (2017). Effect of harvesting time on seed quality of two bell pepper cultivars (*Capsicum annuum*). *Rev. la Fac. Ciencias Agrar.* **49** : 67-77.  
 Sekara, A., Pokhuda, R., Dal Vaehio, L., Somma, S. and Caruso, G. (2017). Interactions among genotype, environment and agronomic practices on production and quality of storage onion (*Allium cepa* L.) - A review. *Hort. Sci.* **44** : 21-42.  
 Surivong, S., Krittigamas, N., Pimmanee, S., Panvane, A. and Veasatip, S. (2015). Influence of storage conditions on survival of hemp seed quality. *Agric. Sci. Procedia.* **5** : 170-76.  
 Windauer, L. B., Martinez, J., Rapoport, D., Wassner, D. and Besech-Arnold, R. (2012). Germination responses to temperature and water potential in *Jatropha curcas* seeds: A hydrotime model explains the difference between dormancy expression and dormancy induction at different incubation temperatures. *Ann. Bot.* **109** : 265-73.

Contact Us  
 Enter Name  
 Enter Email  
 Select Country  
 Message  
 Saya bukan robot  
 Submit

27°C Barawan  
 Search  
 8:54 AM 12/18/2022

Gmail  
 Telusuri email  
 12 dari 982

Online Publication of manuscript ROC-909 \* Effect of harvest age and storage duration on viability and vigor of shallot (*Allium cepa* L.) tubers \*  
 Kotak Masuk

Gaurav Publications  
 kepada saya

Sab, 17 Des 15:32 (5 hari yang lalu)

Dear Dr. MUHAMMAD ANSAR,

We are glad to inform you that your article No. ROC-909 entitled "Effect of harvest age and storage duration on viability and vigor of shallot (*Allium cepa* L.) tubers" has been assigned DOI Number and published online. Article is attached with DOI number. You can find your article online on <https://gauravpublications.com/journal/research-on-crops/ROC-909>.

Do not hesitate to contact us if you have any questions or concerns.

Best Regards  
 Managing Editor  
 Gaurav Publications |  
 # 1314 (GF), Housing Board, Sector-15A, | Hisar - 125001 | Haryana (INDIA) |  
 Web: [www.gauravpublications.com](http://www.gauravpublications.com) | E-mail: [info@gauravpublications.com](mailto:info@gauravpublications.com)

Muhammad Anshar Pasigai - ansharpsigai@gmail.com  
 kepada Gaurav



Sel, 20 Des 20:31 (2 hari yang lalu)

Thank you, our manuscript has been published


Regards  
 Muhammad Ansar

Balas Teruskan

**20 Desember 2022: pdf copy of final published paper (ID 11146)**



Link: <https://www.indianjournals.com/ijor.aspx?target=ijor.rcr&volume=23&issue=4&type=toc>



Online published: 7 Januari 2023

